

1. Administrative & Study Identification

Full Study Title: A Phase II Trial to Evaluate the Clinical Efficacy, Safety and Tolerability of MAS825 in Pediatric and Adult Participants With Still's Disease **Short Title/Acronym:** MAS825 in Still's Disease **Protocol Number:** CMAS825J12201 **NCT Number:** NCT07203001 **Sponsor Name:** Novartis Pharmaceuticals **Investigational Product:** MAS825 (bispecific anti-IL1 β and IL-18 monoclonal antibody) **Phase of Development:** Phase II **Medical Monitor:** Novartis Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective * **Primary Objective:** To evaluate the clinical efficacy, safety, and tolerability of MAS825 in pediatric and adult participants with active Still's Disease. * **Secondary Objectives:** To assess the proportion of participants achieving inactive disease, evaluate changes in disease activity and inflammatory biomarkers (such as CRP and ferritin), and assess the overall pharmacokinetic and safety profile of the treatment.

2.2 Background & Rationale Still's Disease (including systemic Juvenile Idiopathic Arthritis and Adult-Onset Still's Disease) is a multifaceted autoinflammatory disorder that can be complicated by life-threatening conditions such as macrophage activation syndrome (MAS) and interstitial lung disease. IL-1 β and IL-18 play a pivotal role in driving this disease. MAS825, a bispecific anti-IL1 β and IL-18 monoclonal antibody, is being evaluated as an innovative approach to safely reduce inflammasome-mediated inflammation and improve clinical outcomes in drug-resistant or active populations.

3. Study Design & Methodology

3.1 Design Framework * **Study Type:** Interventional * **Control Type:** Single-arm * **Blinding:** Open-label * **Randomization:** N/A

3.2 Planned Enrollment & Duration * **Target \$\$\$:** 20 participants * **Number of Sites:** Multicenter (Including investigative sites in Italy, Spain, Netherlands, and Turkey) * **Estimated Study Start:** 03-Nov-2025 * **Estimated Primary Completion:** 24-May-2029

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Age ≥ 1 year with a diagnosis of Still's Disease. 2. Active disease defined as: CRP or ferritin levels greater than Upper Limit of Normal (ULN), and any of the following: Fever $\geq 38^{\circ}\text{C}$ attributed to disease activity; Rash attributed to Still's Disease; Musculoskeletal involvement (arthritis in a number of joints); Serositis; or Macrophage activation syndrome (MAS) activity. 3. Need for glucocorticoids (prednisone or equivalent). *(Note: Still's disease patients with evidence of MAS or interstitial lung disease requiring supplemental oxygen therapy are permitted in the study)*

4.2 Exclusion Criteria 1. Patients out of the specified weight range. 2. Ongoing or previous treatment with certain immunomodulatory drugs. 3. Glucocorticoid dose exceeding a set protocol limit. 4. History of ongoing, chronic, or possibly recurrent infection (e.g., HIV, TB, HCV, HBV) and/or symptoms and signs of clinically significant active bacterial, fungal, or viral infections. 5. Administration of live vaccinations within a set time prior to MAS825 treatment.

5. Intervention & Treatment Plan

5.1 Investigational Regimen * **Dosage/Frequency:** MAS825 administered per protocol-specified dosing schedule. * **Route of Administration:** Parenteral (Intravenous / Subcutaneous injection/infusion). * **Duration of Treatment:** Ongoing until disease progression, unacceptable toxicity, or study completion (May 2029).

5.2 Concomitant & Prohibited Therapy * **Permitted:** Standard of care therapies and glucocorticoids (within protocol limits). * **Prohibited:** Ongoing or previous treatment with prohibited immunomodulatory drugs and live vaccinations.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening	Day -14 to 0 Informed Consent, Medical History, Vital Signs, \$CRP\$ and Ferritin Labs
Baseline	Day 0	First Dose, Safety Labs, Disease Activity Evaluation	Interim Regular Intervals Safety Labs, Efficacy Assessment (Inactive disease criteria), Biomarker monitoring
End of Study	May 2029	Final Vital Signs, Exit Interview, IP Accountability	

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint * **Definition:** Proportion of participants achieving inactive disease. Disease activity is evaluated by the investigator based on physical examination and laboratory parameters. * **Measurement Method:** Protocol-specific criteria which include clinical signs, laboratory values (e.g., CRP, Ferritin), and physician assessment.

7.2 Secondary & Exploratory Endpoints * Pharmacokinetics and safety tolerability (Frequency and severity of AEs and SAEs). * Evaluation of changes in biomarkers associated with IL-1 β and IL-18 pathway engagement.

8. Safety & Adverse Event Reporting

- **Adverse Event (AE) Monitoring:** Standard clinical monitoring for AEs, with particular attention to invasive bacterial infections and immunosuppression-related complications.
 - **Serious Adverse Events (SAEs):** Routine reporting timeline, typically 24 hours of site awareness.
 - **Data Monitoring Committee (DMC):** Internal/External safety review committees according to Novartis standards.
-

9. Statistical Analysis Plan (SAP) Summary

- **Analysis Populations:** Intent-to-Treat (ITT) for primary clinical outcomes, Safety Set for adverse event tracking.
 - **Sample Size Justification:** $N = 20$, designed to assess early clinical efficacy signals and safety in a small, targeted open-label population.
 - **Handling of Missing Data:** Last Observation Carried Forward (LOCF) or Multiple Imputation depending on the exact continuous variable.
-

10. Quality Assurance & Regulatory Compliance

- **GCP Compliance:** This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.
 - **Monitoring Plan:** Standardized onsite and remote monitoring visits to ensure safety and protocol fidelity.
 - **Audit Trail:** Details on eCRF locking and data clarification forms via standard quality audit systems.
-

11. Study Identifiers & Governance

- **Primary ID:** CMAS825J12201

- **Registry Number:** NCT07203001
 - **Sponsor/Lead Organization:** Novartis Pharmaceuticals
 - **Study Title:** A Phase II Trial to Evaluate the Clinical Efficacy, Safety and Tolerability of MAS825 in Pediatric and Adult Participants With Still's Disease
 - **Official Regulatory Title:** A Phase II Trial to Evaluate the Clinical Efficacy, Safety and Tolerability of MAS825 in Pediatric and Adult Participants With Still's Disease
-

12. Clinical Framework (Still's Disease Specific)

- **Primary Condition:** Still's Disease (Juvenile Idiopathic Arthritis / Adult-Onset Still's Disease)
 - **Diagnosis Threshold:** Active disease (CRP or ferritin > ULN, Fever $\geq 38^{\circ}\text{C}$)
 - **Medical Methodology:** Bispecific anti-IL1 β and IL-18 monoclonal antibody therapy
 - **Optimization Target:** Achievement of inactive disease status and inflammasome mediation
-

13. Study Procedures & Methodology

- **Management Pathway:** Standard clinical assessment for Still's disease and Macrophage Activation Syndrome (MAS).
 - **Education Protocol (HCP):** Investigator training on disease activity evaluation criteria.
 - **Education Protocol (Patient):** Trial onboarding, safety, and immunosuppression risk awareness.
 - **Quality Audit Mechanism:** Clinical signs review, laboratory value tracking, and physician assessment tracking.
-

14. Observation & Follow-Up Schedule

- **Total Duration:** Ongoing observation up to study completion (May 2029)
 - **Onsite Visit Cadence:** Regular intervals for biologic administration and clinical assessment.
 - **Remote Monitoring Frequency:** Remote checks as mandated by institutional protocols.
 - **Checklist Review Interval:** Every clinical visit.
-

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	____	[DD-MMM-YYYY]				Lead
Statistician	[Name]	____	[DD-MMM-YYYY]				