

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 02-Apr-2026

1. Administrative & Study Identification

Full Study Title A Study to Evaluate the Safety of Empliciti® (Elotuzumab) When Treating Patients With Multiple Myeloma in Taiwan **Short Title/Acronym** Empliciti PMS Taiwan **Protocol Number** CA204-219 **NCT Number** NCT06163040 **Posting ID** 414968705837422117 **Sponsor Name** Bristol Myers Squibb **Investigational Product** Empliciti® (elotuzumab) **Phase of Development** Not Applicable **Trial Status** COMPLETED (Study has been completed) **Medical Monitor** Bristol-Myers Squibb Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To collect safety data (Adverse Events [AEs], Serious Adverse Events [SAEs], and AEs of Special Interest [AESI]) for a period of 25 weeks or up to 30 days after the discontinuation of Empliciti (elotuzumab) treatment in combination with pomalidomide and dexamethasone (or lenalidomide and dexamethasone) in patients with relapsed or refractory multiple myeloma (RRMM) in Taiwan. **Secondary Objectives:** To assess the real-world clinical characteristics and ongoing safety profile of the elotuzumab combination regimens.

2.2 Background & Rationale To evaluate the safety and clinical applicability of elotuzumab in a real-world setting. This observational study aims to assess the safety of elotuzumab when used in combination with pomalidomide and dexamethasone (EPd) for the treatment of RRMM in participants who had received at least two prior therapies, including lenalidomide and a proteasome inhibitor. It also evaluates elotuzumab in combination with lenalidomide and dexamethasone (ELd) in RRMM participants who received one to three prior therapies.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational **Status:** COMPLETED **Control Type:** Single-arm **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: 7 participants **Number of Sites:** Multiple centers in Taiwan **Estimated Study Start:** 22-Dec-2023 **Estimated Primary Completion:** 17-Mar-2025

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Age $\geq$ 18 years. 2. Confirmed diagnosis of Relapsed or Refractory Multiple Myeloma (RRMM). 3. Received $\geq$ 2 prior therapies including lenalidomide and a proteasome inhibitor (planning to receive EPd) OR received 1 to 3 prior therapies (planning to receive ELd) at the physician's medical judgment.

4.2 Exclusion Criteria 1. Participants with therapeutic indications for which elotuzumab in combination with pomalidomide/lenalidomide and dexamethasone has not been approved in Taiwan. 2. Participants who are contraindicated for treatment with elotuzumab in combination with pomalidomide/lenalidomide and dexamethasone (as described in the Taiwan label). 3. Participants who participate in other interventional clinical trials.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Drug Name: elotuzumab Trade Name: Empliciti® ATC Code: L01FX08 Dosage/Frequency: Elotuzumab administered in combination with either pomalidomide and dexamethasone (EPd) or lenalidomide and dexamethasone (ELd), as per the approved local product label and the treating physician's discretion. Route of Administration: Intravenous (elotuzumab) and Oral (pomalidomide/lenalidomide and dexamethasone). Duration of Treatment: 25 weeks or until disease progression, unacceptable toxicity, or patient withdrawal. EMA URL: https://www.ema.europa.eu/en/documents/product-information/empliciti-epar-product-information_en.pdf

5.2 Concomitant & Prohibited Therapy Permitted: Routine concomitant medications and supportive care therapies as required for the management of multiple myeloma comorbidities. **Prohibited:** Participation in any other interventional clinical trials while receiving study treatment.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -14 to 0	Informed Consent, Medical History, Confirmation of RRMM Diagnosis and Prior Therapies
Baseline	Day 0	Initiation of Elotuzumab combination therapy
Interim	Up to Week 25	Standard-of-care clinical visits; Collection of safety data (AEs, SAEs, AESIs)
End of Study	+30 Days	30 days post-discontinuation of Empliciti for final safety follow-up

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: 1. Number of participants with adverse events: Assessed according to NCI CTCAE v5.0. 2. Number of participants with adverse events of special interest: Opportunistic infections including viral hepatitis and tuberculosis reactivation. **Measurement Method:** Clinical assessment and physician reporting as per routine observational post-marketing surveillance over the study duration (25 weeks or 30 days post-treatment).

7.2 Secondary & Exploratory Endpoints Secondary Outcomes: 1. Number of participants with fungal infections. 2. Number of participants with reactivation of viral hepatitis. 3. Number of participants with other viral infection/reactivation. **Exploratory Endpoints:** Real-world treatment duration and overall clinical characteristics in the evaluated patient cohort.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Physician-led collection of AE and AESI data continuously through routine clinic visits. **Serious Adverse Events (SAEs):** Routine pharmacovigilance timeline reporting (typically 24 hours from site awareness) to the sponsor and regulatory authorities. **Data Monitoring Committee (DMC):** Internal BMS safety monitoring and post-marketing surveillance registry oversight.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all enrolled participants who received at least one dose of the investigational product). **Sample Size Justification:** As an observational post-marketing surveillance study, sample size is based on real-world patient availability within the surveillance timeframe in Taiwan ($N=75$). **Handling of Missing Data:** Descriptive statistics applied; no formal imputation for missing data in real-world safety tabulations.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study is conducted in accordance with Good Pharmacoepidemiology Practices (GPP) and local Taiwanese regulatory guidelines for post-marketing surveillance. **Monitoring Plan:** Centralized remote data monitoring via site-reported electronic registries. **Audit Trail:** Secure database locking and standard data clarification procedures managed by the sponsor.

11. Study Identifiers & Governance

Primary ID: CA204-219 **Registry Number:** NCT06163040 **Posting ID:** 414968705837422117 **Sponsor/Lead Organization:** Bristol Myers Squibb **Study Title:** A Study to Evaluate the Safety of Empliciti® (Elotuzumab) When Treating Patients With Multiple Myeloma in Taiwan **Official Regulatory Title:** Empliciti® (elotuzumab) Post-marketing Surveillance Study for Patients with Multiple Myeloma in Taiwan

12. Clinical Framework (Adapted for Multiple Myeloma)

Disease Area: Relapsed or Refractory Multiple Myeloma (RRMM) **Name:** Refractory Multiple Myeloma **Preferred Name:** Relapsed/refractory multiple myeloma (Preferred UMLS Name: Relapsed or Refractory Multiple Myeloma) **Semantic Type:** Neoplastic Process **SNOMED ID:** 763261000000108 **SNOMED Hierarchy:** 703197003|Multiple myeloma|Multiple myeloma (disorder) **Disease Definition:** Refractory Multiple Myeloma is a form of multiple myeloma, a cancer of plasma cells in the bone marrow, where the disease progresses during treatment or within a short period (typically 60 days) after the last dose of therapy. This indicates that the cancer has become resistant to the current or recent therapeutic regimens, making it challenging to achieve remission or control disease progression with standard treatments. **Diagnosis Threshold:** Confirmed RRMM requiring 2^{nd} line or later systemic therapy **Medical Methodology:** Monoclonal antibody immunotherapy combined with Immunomodulatory Drugs (IMiDs) and Corticosteroids **Optimization Target:** Disease control and continuous safety/toxicity monitoring

13. Study Procedures & Methodology

Management Pathway: Standard clinical treatment pathway for RRMM in Taiwan. **Education Protocol (HCP):** Post-marketing safety and reporting training for treating physicians. **Education Protocol (Patient):** Routine standard-of-care disease education for MM treatment. **Quality Audit Mechanism:** Ongoing safety surveillance data review and sponsor pharmacovigilance processing.

14. Observation & Follow-Up Schedule

Total Duration: 25 weeks of treatment observation + 30 days safety follow-up. **Onsite Visit Cadence:** Directed by standard-of-care clinical requirements. **Remote Monitoring Frequency:** Routine periodic data extractions from the patient registry. **Checklist Review Interval:** As dictated by standard clinical MM treatment cycles (e.g., 21-day or 28-day cycles).

15. Sign-off & Approval

Role	Name	Signature	Date		:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]					Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]					