

1. Administrative & Study Identification

Full Study Title: A PHASE 1, OPEN-LABEL, 2-PERIOD, FIXED-SEQUENCE STUDY TO INVESTIGATE THE ABSORPTION, DISTRIBUTION, METABOLISM, AND EXCRETION OF [14C]PF-07976016 AND TO ASSESS THE ABSOLUTE BIOAVAILABILITY AND FRACTION ABSORBED OF PF-07976016 IN HEALTHY MALE PARTICIPANTS USING A [14C]-MICROTRACER APPROACH **Short Title/ Acronym:** A Study to Understand How the Study Medicine (PF-07976016) is Processed and Eliminated in Healthy Men **Protocol Number:** EUCT2024-518535-11-00 **NCT Number:** NCT07015918 **Sponsor Name:** Pfizer Inc. **Investigational Product:** PF-07976016 (GIPR antagonist) and [14C]PF-07976016 microtracer **Phase of Development:** Phase 1 **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To investigate the absorption, distribution, metabolism, and excretion (ADME) profile of PF-07976016 in healthy male participants using a [14C]-microtracer approach. **Secondary Objectives:** To assess the absolute bioavailability and fraction absorbed of PF-07976016, and to evaluate the safety and tolerability of the study medicine.

2.2 Background & Rationale PF-07976016 is a novel Gastric Inhibitory Polypeptide Receptor (GIPR) antagonist under development for the treatment of obesity and related metabolic diseases. Understanding its human ADME properties and absolute bioavailability is a foundational regulatory requirement in early clinical development to guide future dosing and formulation strategies.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional (Basic Science) **Control Type:** Single Group Assignment (Fixed-Sequence) **Blinding:** Open-Label **Randomization:** N/A (Fixed-Sequence)

3.2 Planned Enrollment & Duration Target \$N\$: Small cohort of healthy male participants (standard for Phase 1 ADME microtracer studies) **Number of Sites:** 1 (ICON, Groningen, Netherlands) **Estimated Study Start:** Early 2025 **Estimated Primary Completion:** August 2025

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Healthy males, 18 to 65 years of age. 2. BMI 17.5 to 32 kg/m² and a total body weight ≥ 50 kg (110 lb). 3. Willing and able to comply with all study procedures.

4.2 Exclusion Criteria 1. History of irregular bowel movements or lactose intolerance. 2. Positive urine drug test, history of alcohol abuse or repeated binge drinking, or use of tobacco/nicotine products ≥ 5 cigarettes/day. 3. Any medical or psychiatric condition, clinical laboratory abnormality, use of prohibited/concomitant medication(s), or previous administration of an investigational product within 30 days preceding the first dose.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Single doses of PF-07976016 administered with and without a [14C] radiolabel in a 2-period, fixed-sequence crossover design. **Route of Administration:** Oral (PO) and Intravenous (IV). **Duration of Treatment:** Up to 3 study periods separated by a 7-day washout period between doses. Total study participation spans approximately 12 weeks.

5.2 Concomitant & Prohibited Therapy Permitted: Generally, no medications are permitted in healthy volunteer Phase 1 ADME studies to prevent PK interference. **Prohibited:** Use of any prohibited or concomitant medication(s) throughout the duration of the study.

6. Study Timeline & Visit Schedule

Visit ID	Window	Key Activities/Procedures
1	1-2 Weeks prior to Day 1	Screening: Informed Consent, Medical History, ECG, Vital Signs, Urine Drug Test
2	Day 1 (per period)	Treatment Periods: Dosing of PF-07976016 (with/without [14C]), extensive PK/radiolabel blood and excreta sampling
3	7 days between doses	Washout: Safety monitoring and drug clearance
4	Up to 21 Days post-dose	End of Study / Follow-Up: Final PK and radioactivity sampling, Safety Labs, Exit Interview

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Total recovery of radioactivity expressed as a percent of the total oral radioactive dose administered, monitored from Day 1 and up to 21 days thereafter. **Measurement Method:** Scintillation counting or accelerator mass

spectrometry (AMS) of blood, urine, and feces samples to map absorption, distribution, metabolism, and excretion.

7.2 Secondary & Exploratory Endpoints 1. Absolute bioavailability and fraction absorbed of PF-07976016. 2. Incidence and severity of Adverse Events (AEs) to assess safety and tolerability.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring of AEs, vital signs, clinical laboratory tests, and ECGs throughout the in-house stays and follow-up period. **Serious Adverse Events (SAEs):** Reported within 24 hours to the Sponsor and regulatory bodies as per standard Phase 1 safety protocols. **Data Monitoring Committee (DMC):** Internal safety review by the Sponsor and Principal Investigator prior to advancing between sequences.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Pharmacokinetic (PK) Set for ADME parameters and the Safety Set for all participants receiving at least one dose. **Sample Size Justification:** Sample size is based on practical and historical considerations for human radiolabeled microtracer studies rather than formal statistical power, which is standard for profiling ADME metrics. **Handling of Missing Data:** Standard PK conventions will be used; values below the lower limit of quantification (LLOQ) will be handled per the statistical analysis plan.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Dedicated onsite Phase 1 clinic monitoring at the designated unit (ICON). **Audit Trail:** Robust tracking of all radioactive materials, eCRF locking, and rigorous quality control for AMS and PK bioanalytical assays.

11. Study Identifiers & Governance

Primary ID: EUCT2024-518535-11-00 **Registry Number:** NCT07015918
Sponsor/Lead Organization: Pfizer Inc. **Study Title:** PF-07976016 ADME and Bioavailability Study **Official Regulatory Title:** A PHASE 1, OPEN-LABEL, 2-PERIOD, FIXED-SEQUENCE STUDY TO INVESTIGATE THE ABSORPTION, DISTRIBUTION, METABOLISM, AND EXCRETION OF [14C]PF-07976016 AND TO ASSESS THE ABSOLUTE BIOAVAILABILITY AND FRACTION ABSORBED OF PF-07976016 IN HEALTHY MALE PARTICIPANTS USING A [14C]-MICROTRACER APPROACH

12. Clinical Framework (Metabolic Specific)

Primary Condition: Healthy Volunteers (Investigational agent for Obesity) **Diagnosis Threshold:** N/A (Healthy Males, BMI 17.5 to 32 kg/m²) **Medical Methodology:** Pharmacokinetic and ADME Profiling via [14C]-microtracer **Optimization Target:** Defining the clinical pharmacology, absolute bioavailability, and elimination pathways of the GIPR antagonist PF-07976016.

13. Study Procedures & Methodology

Management Pathway: Intensive inpatient monitoring in a specialized Phase 1 clinical research unit. **Education Protocol (HCP):** Specialized site training on radiolabeled compound administration and microtracer collection. **Education Protocol (Participant):** Comprehensive briefing on inpatient restrictions, dietary constraints, and total excreta collection requirements. **Quality Audit Mechanism:** Bioanalytical lab certifications for radiocarbon analysis and PK quantification.

14. Observation & Follow-Up Schedule

Total Duration: Up to ~12 weeks **Onsite Visit Cadence:** Intensive inpatient stays during the dosing periods, followed by up to 21 days of post-dose sampling and follow-up. **Remote Monitoring Frequency:** N/A (Predominantly clinic-based Phase 1 confinement) **Checklist Review Interval:** Daily during confinement to monitor AEs and strict dietary/medication restrictions.

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				