

Clinical Study Summary Report (CSSR)

Document Status: Draft | Version: 1.0 | Date: 03-Apr-2026

1. Administrative & Study Identification

Full Study Title: A Study to Learn About the Study Medicine Called PF-08046031 in Advanced Melanoma and Other Solid Tumors

Short Title/Acronym: PF-08046031 Phase 1 Study

Protocol Number: C5901001 (Secondary ID: 2024-514745-10-00)

NCT Number: NCT06799533

Sponsor Name: Pfizer Inc.

Official Regulatory Title: AN OPEN-LABEL PHASE 1 STUDY TO INVESTIGATE PF-08046031 IN ADULTS WITH ADVANCED MELANOMA AND OTHER SOLID TUMORS

Investigational Product: PF-08046031 (formerly SGN-CD228A, a CD228-directed antibody-drug conjugate)

Phase of Development: Phase I

Medical Monitor: Pfizer Inc. Clinical Team

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective: To evaluate the safety and tolerability of PF-08046031 and identify the dose-limiting toxicities and optimal dosing level (Optimization Target: Safety profile characterization and optimal dosing level determination) in participants with advanced melanoma and other solid tumors.

Secondary Objectives: To evaluate the side effects, pharmacokinetic profile, and preliminary antitumor efficacy (e.g., Objective Response Rate) of PF-08046031.

2.2 Background & Rationale

To address the unmet medical need in treating advanced solid tumors utilizing targeted antibody-drug conjugates (ADCs) (Medical Methodology: CD228-directed Antibody-Drug Conjugate (ADC) Therapy). PF-08046031 is a CD228-directed ADC containing a monomethyl auristatin E (MMAE) payload, designed to target the CD228 antigen (melanotransferrin), which is highly expressed on melanoma, non-small cell lung cancer (NSCLC), head and neck squamous cell carcinoma (HNSCC), and esophageal cancer cells. The trial aimed to test safety and efficacy; however, clinical development was halted in March 2026 for strategic portfolio prioritization reasons following Pfizer's acquisition of Seagen, with no underlying safety or efficacy concerns reported.

3. Study Design & Methodology

3.1 Design Framework

Study Type: Interventional

Control Type: Single-arm

Blinding: Open-label (None)

Randomization: N/A (Sequential Assignment)

3.2 Planned Enrollment & Duration

Target \$N\$: 11 participants (trial terminated early)

Number of Sites: Multiple clinical sites globally (e.g., UCSF)

Estimated Study Start: 2024

Estimated Primary Completion: March 2026 (Terminated)

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

1. Age $\geq$ 18 years.
2. Participants in Part 1 (dose escalation) and Part 2 (dose optimization) must have histologically- or cytologically-confirmed metastatic or unresectable cutaneous melanoma (Diagnosis Threshold).
3. Progressive disease following at least 1 prior anti-PD-1/PD-L1 immunotherapy containing regimen, with no appropriate standard therapy available.
4. For Part 3 (dose expansion): Must have histologically- or cytologically-confirmed metastatic or unresectable solid malignancy from one of the following: cutaneous melanoma, NSCLC, HNSCC, or esophageal cancer.

4.2 Exclusion Criteria

1. Active cerebral/meningeal disease related to the underlying malignancy.
2. Previous exposure to CD228-targeted therapy, vedotin or an MMAE-containing agent, or any taxane containing regimen for advanced disease.
3. Grade 3 or higher pulmonary disease unrelated to underlying malignancy, previous history of non-infectious interstitial lung disease (ILD) or pneumonitis that required steroids, or suspected ILD that cannot be ruled out by screening imaging.

5. Intervention & Treatment Plan

5.1 Investigational Regimen

Dosage/Frequency: Ascending doses of PF-08046031 to determine maximum tolerated dose (MTD) in Part 1 and optimal biological dose in Part 2, followed by expansion in Part 3.

Route of Administration: Intravenous (IV) infusion (Management Pathway: Intravenous administration of experimental ADC PF-08046031).

Duration of Treatment: Until disease progression, unacceptable toxicity, withdrawal of consent, or study termination (Total Duration: Study treatment duration plus 30 days safety follow-up).

5.2 Concomitant & Prohibited Therapy

Permitted: Standard supportive care measures for disease symptoms.

Prohibited: Chemotherapy, definitive radiotherapy, biologics, and/or other antitumor treatment with immunotherapy within 4 weeks prior to the first dose of study intervention.

6. Study Timeline & Visit Schedule

Onsite Visit Cadence: As per Phase 1 sequential dosing and assessment schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent (Education Protocol Patient: Informed consent process detailing Phase 1 risks and side effects), Medical History, Tumor Imaging, Baseline Labs
Baseline	Day 0	Enrollment confirmation, First IV Dose
Interim	Per Treatment Cycle	Safety Labs, PK/PD blood draws, Efficacy Assessments (Tumor scans)
End of Study	+30 Days	Final Vital Signs, Exit Interview, Safety follow-up 30 days after last dose

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint

Definition: Number of participants with adverse events (AEs) and laboratory abnormalities evaluated through 30 days after the last study treatment (approximately 6 months).

Measurement Method: Clinical assessment and central/local laboratory analysis grading AEs based on CTCAE criteria.

7.2 Secondary & Exploratory Endpoints

Objective Response Rate (ORR) and other preliminary antitumor activity markers.

Pharmacokinetics (PK) parameters for PF-08046031.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring of any untoward medical occurrence through clinical visits and laboratory findings up to 30 days after the last treatment (Checklist Review Interval: Continuous evaluation during the treatment period).

Serious Adverse Events (SAEs): Standard Phase 1 reporting timelines (typically 24 hours to the sponsor).

Data Monitoring Committee (DMC): Internal safety review committee assessing dose-escalation cohorts.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Analysis Set (all patients receiving at least one dose of the investigational product).

Sample Size Justification: Study was designed as a Phase 1 dose-escalation/expansion without formal powered hypothesis testing; however, terminated early after 11 patients.

Handling of Missing Data: Standard handling for early-phase oncology studies.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: High-frequency onsite clinical monitoring typical of Phase 1 First-in-Human or early-phase dose-escalation trials. Remote monitoring will occur as needed for ongoing safety evaluation.

Audit Trail: eCRF locking, query management, and standard sponsor auditing. Routine clinical site monitoring and safety data review (Quality Audit Mechanism).

Training/Education Protocol (HCP): Site initiation and protocol training for Phase 1 ADC toxicities.

11. Study Identifiers & Governance

Primary ID: {{C5901001}}

Registry Number: {{NCT06799533}}

Sponsor/Lead Organization: {{Pfizer Inc.}}

Study Title: {{A Study to Learn About the Study Medicine Called PF-08046031 in Advanced Melanoma and Other Solid Tumors}}

Official Regulatory Title: {{AN OPEN-LABEL PHASE 1 STUDY TO INVESTIGATE PF-08046031 IN ADULTS WITH ADVANCED MELANOMA AND OTHER SOLID TUMORS}}

12. Clinical Framework (Oncology Specific)

Primary Condition: {{Advanced Melanoma and Other Solid Tumors}}

Diagnosis Threshold: {{Histologically- or cytologically-confirmed metastatic or unresectable disease}}

Medical Methodology: {{CD228-directed Antibody-Drug Conjugate (ADC) Therapy}}

Optimization Target: {{Safety profile characterization and optimal dosing level determination}}

13. Study Procedures & Methodology

Management Pathway: {{Intravenous administration of experimental ADC PF-08046031}}

Education Protocol (HCP): {{Site initiation and protocol training for Phase 1 ADC toxicities}}

Education Protocol (Patient): {{Informed consent process detailing Phase 1 risks and side effects}}

Quality Audit Mechanism: {{Routine clinical site monitoring and safety data review}}

14. Observation & Follow-Up Schedule

Total Duration: {{Study treatment duration plus 30 days safety follow-up}}

Onsite Visit Cadence: {{As per Phase 1 sequential dosing and assessment schedule}}

Remote Monitoring Frequency: {{As needed for ongoing safety evaluation}}

Checklist Review Interval: {{Continuous evaluation during the treatment period}}

15. Sign-off & Approval

Role	Name	Signature	Date		:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]					Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]					