

1. Administrative & Study Identification

Full Study Title: A Prospective, Open-Label, Multicenter Randomized Phase III Trial to Compare The Efficacy and Safety of A Combined Regimen of Obinutuzumab and Venetoclax (GDC-0199/ABT-199) Versus Obinutuzumab and Chlorambucil in Previously Untreated Patients With CLL and Coexisting Medical Conditions
Short Title/Acronym: CLL14 / B025323 **Protocol Number:** B025323 **NCT Number:** NCT02242942 **Sponsor Name:** F. Hoffmann-La Roche Ltd / Roche **Investigational Product:** Venetoclax (GDC-0199/ABT-199) + Obinutuzumab; Chlorambucil + Obinutuzumab **Phase of Development:** Phase III **Medical Monitor:** F. Hoffmann-La Roche Ltd Clinical Operations / [Name and Contact Information]

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To compare the efficacy and safety of a combined targeted regimen of obinutuzumab and venetoclax versus the standard chemoimmunotherapy regimen of obinutuzumab and chlorambucil in patients with chronic lymphocytic leukemia (CLL) and coexisting medical conditions.
Secondary Objectives: To evaluate Progression Free Survival (PFS) based on Institutional Review Committee (IRC) assessments, Overall Response (OR) at completion of treatment, and Complete Response Rate (CRR) at completion of treatment.

2.2 Background & Rationale To provide a targeted therapy combination with a definite, fixed duration of treatment (12 cycles) in previously untreated CLL patients with coexisting conditions. This fixed-duration approach aims to limit the toxicity, resistance risks, and costs associated with continuous/indefinite treatments while achieving deep responses and superior progression-free survival (PFS). The time on study treatment is approximately one year and the follow-up period will be up to 9 years.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Study Status:** Completed (Study has been completed) **Control Type:** Active Comparator (Obinutuzumab + Chlorambucil) **Blinding:** Open-label **Randomization:** 1:1, Randomized

3.2 Planned Enrollment & Duration Target **\$\$\$:** 445 patients **Number of Sites:** Multicenter (Global) **Estimated Study Start:** August 2015 **Estimated Primary Completion:** May 2019

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Documented previously untreated CLL requiring treatment according to the International Workshop on Chronic Lymphocytic Leukemia (IWCLL) criteria. 2. Total Cumulative Illness Rating Scale (CIRS) score > 6, indicating coexisting medical conditions. 3. Adequate marrow function independent of growth factor or transfusion support within 2 weeks of screening, unless cytopenia is due to marrow involvement of CLL. 4. Adequate liver function and a life expectancy > 6 months. 5. Agreement to use highly effective contraceptive methods per protocol.

4.2 Exclusion Criteria 1. Transformation of CLL to aggressive Non-Hodgkin's lymphoma (Richter's transformation or pro-lymphocytic leukemia). 2. Known central nervous system involvement or history of confirmed progressive multifocal leukoencephalopathy (PML). 3. Inadequate renal function, uncontrolled autoimmune hemolytic anemia, or immune thrombocytopenia. 4. An individual organ/system impairment score of 4 as assessed by CIRS limiting ability to receive treatment (exception: eyes, ears, nose, throat). 5. History of prior malignancy (with exceptions per protocol), or use of investigational agents/concurrent anti-cancer treatment within the last 4 weeks. 6. Active bacterial, viral, or fungal infection requiring systemic treatment within the last two months; known infection with HIV, HTLV-1, or positive tests for HBV (HBsAg) or HCV. 7. History of severe allergic or anaphylactic reactions to humanized/murine monoclonal antibodies, or hypersensitivity to chlorambucil, obinutuzumab, venetoclax, or any excipients. 8. Pregnant women and nursing mothers. 9. Requires the use of warfarin, marcumar, or phenprocoumon, or received strong/moderate CYP3A inhibitors or inducers within 7 days prior to first dose.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: * **Experimental Arm (Venetoclax + Obinutuzumab):** Venetoclax oral tablet initiated with a 5-week ramp-up (20 mg to 400 mg daily) and maintained at 400 mg daily for 12 cycles. Obinutuzumab administered intravenously (IV) at 1000 mg per cycle for the first 6 cycles. * **Active Comparator Arm (Chlorambucil + Obinutuzumab):** Chlorambucil (Trade name: Leukeran; ATC Code: L01AA02) 0.5 mg/kg orally on Day 1 and Day 15 of each 28-day cycle for 12 cycles. Obinutuzumab administered IV for the first 6 cycles. **Route of Administration:** Oral (Venetoclax, Chlorambucil) and IV (Obinutuzumab) **Duration of Treatment:** Fixed-duration of 12 cycles (28 days per cycle).

5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care and premedication for infusion-related reactions. **Prohibited:** Strong and moderate CYP3A inhibitors or inducers

within 7 days prior to study drug; use of warfarin, marcumar, or phenprocoumon.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, CIRS Scoring, Labs
Baseline	Day 0 (Cycle 1, Day 1)	Randomization, First Dose Obinutuzumab, Safety Labs
Interim	Cycles 1 - 12	Venetoclax Ramp-Up (C1-C2), Obinutuzumab infusions (C1-C6), Safety Labs, Efficacy Assessment
End of Treatment	End of Cycle 12	Final Vital Signs, Exit Interview, Efficacy Review, IP Accountability
Follow-Up	Post-Cycle 12	Ongoing PFS tracking and long-term evaluation

(Note: Cycles comprise 28 days)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Progression-Free Survival (PFS), defined as the time from randomization to the first occurrence of disease progression or death from any cause. **Measurement Method:** Investigator-assessed disease progression according to IWCLL 2008 criteria.

7.2 Secondary Endpoints

- 1. Progression Free Survival (PFS) (IRC-Assessed):** Based on Institutional Review Committee (IRC) assessments according to IWCLL criteria.
- 2. Overall Response (OR):** Percentage of participants with an OR at completion of treatment, as determined by the investigator according to IWCLL criteria.
- 3. Complete Response Rate (CRR):** Percentage of participants with a complete response rate at the completion of treatment assessment, as determined by the investigator according to IWCLL criteria.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: AEs collected up to 28 days after the last dose of study treatment. **Serious Adverse Events (SAEs):** Specific tracking for Tumor Lysis Syndrome (TLS), neutropenia, and infusion-related reactions. Grade ≥ 3 AEs reported up to 6 months post-treatment; Grade ≥ 3 infections reported for 2 years after the last dose. **Data Monitoring Committee (DMC):** External/Internal committee details reviewing safety run-in phase and ongoing safety profiles.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) for primary efficacy; Safety Set for all patients receiving at least one dose of the study drug. **Sample Size Justification:** Target enrollment of 445 participants, adequately powered to demonstrate a statistically significant improvement in PFS over the active comparator. **Handling of Missing Data:** Standard survival analysis methodologies (Kaplan-Meier estimates) with right-censoring for missing long-term follow-up data.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Routine onsite and remote monitoring for adherence to the protocol and accurate toxicity tracking. **Audit Trail:** Independent review and data clarification maintained securely via eCRF.

11. Study Identifiers & Governance

Primary ID: {{B025323}} **Posting ID:** {{551814129434349837}}
Registry Number: {{NCT02242942}} **Sponsor/Lead Organization:** {{F. Hoffmann-La Roche Ltd / Roche}} **Study Title:** {{Comparison of the Treatments of Obinutuzumab + Venetoclax Versus Obinutuzumab + Chlorambucil in Patients With Chronic Lymphocytic Leukemia}}
Official Regulatory Title: {{A Prospective, Open-Label, Multicenter Randomized Phase III Trial to Compare The Efficacy and Safety of A Combined Regimen of Obinutuzumab and Venetoclax Versus Obinutuzumab and Chlorambucil in Previously Untreated Patients With CLL and Coexisting Medical Conditions}}

12. Clinical Framework

Primary Condition: {{Chronic Lymphocytic Leukemia (CLL) / Leukemia}} **Condition Semantic Type:** {{Neoplastic Process}}
Condition Definition: {{A cancer of the blood and bone marrow characterized by an abnormal proliferation of leukocytes.}}
Diagnosis Threshold: {{IWCLL diagnostic criteria requiring treatment and CIRS > 6}} **Medical Methodology:** {{Fixed-Duration Targeted Therapy}} **Optimization Target:** {{Progression-Free Survival}} **Terminology Codes:** {{MeSH: D007938 | SNOMED: 93143009}}

13. Study Procedures & Methodology

Management Pathway: {{Randomized active-controlled protocol}}
Education Protocol (HCP): {{TLS risk assessment and Venetoclax ramp-up protocol training}} **Education Protocol (Patient):** {{Onsite training for daily oral adherence and hydration for TLS prevention}} **Quality Audit Mechanism:** {{Central safety monitoring and response verification}}

14. Observation & Follow-Up Schedule

Total Duration: {{Treatment up to 12 cycles (~1 year) + Long-term follow-up (up to 9 years)}} **Onsite Visit Cadence:** {{Days 1, 8, 15, and 22 of Cycle 1 and 2; Day 1 of subsequent cycles}} **Remote Monitoring Frequency:** {{Regular intervals during long-term follow-up}} **Checklist Review Interval:** {{Every cycle during active treatment}}

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]			Lead	
Statistician	[Name]	_____	[DD-MMM-YYYY]				