

1. Administrative & Study Identification

Full Study Title: A Phase 1, Open-label, Fixed Sequence Study to Evaluate The Effect of Steady-State Dosing of PF-08049820 on the Single Dose Pharmacokinetics of Oral Contraceptives, Midazolam, and Dabigatran in Healthy Adult Female Participants **Brief Title:** A Study to Learn If the Study Medicine Called PF-08049820 Changes How the Body Processes the Other Study Medicines Called Oral Contraceptives, Midazolam, and Dabigatran in Healthy Adult Female Participants **Short Title/Acronym:** PF-08049820 with Midazolam, Dabigatran, Birth Control **Protocol Number:** C6231004 **NCT Number:** NCT07190430 **Trial Status:** Recruiting **Sponsor Name:** Pfizer **Investigational Product:** PF-08049820 (co-administered with Midazolam, Dabigatran Etexilate, and Portia / Oral Contraceptives) **Phase of Development:** Phase I **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the effect of steady-state dosing of PF-08049820 on the single-dose pharmacokinetics of oral contraceptives, midazolam, and dabigatran in healthy adult female participants. **Secondary Objectives:** To assess the overall safety, tolerability, and clinical response of PF-08049820 when administered alone and in combination with the specified medications.

2.2 Background & Rationale The purpose of this study is to see how a medicine called PF-08049820 (in development for Atopic Dermatitis) affects how other medicines move through the body (drug-drug interactions). This information is critical to understanding the PK profile of PF-08049820 in relation to common medications (birth control pills, midazolam for relaxation/sleep, and dabigatran etexilate for blood clot prevention) and will help safely plan future Phase 2/3 clinical studies.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm (Fixed Sequence) **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: 16 Healthy adult female participants (Standard Phase 1 cohort) **Number of Sites:** United States (Clinic Confinement) **Estimated Study Start:** 03-

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Female participants ≥ 18 years of age at screening who are overtly healthy as determined by medical evaluation including medical history, physical examination, laboratory tests, and cardiac monitoring. 2. BMI of 16 to 32 kg/m²; and a total body weight >50 kg (110 lb). 3. Participants who are willing and able to comply with all scheduled visits, treatment plans, laboratory tests, and other study procedures.

4.2 Exclusion Criteria 1. Evidence or history of clinically significant hematological, renal, endocrine, pulmonary, gastrointestinal, cardiovascular, hepatic, psychiatric, neurological, or allergic disease (including drug allergies). 2. History of myocardial infarction, unstable angina, arterial revascularization, mechanical prosthetic heart valve, stroke, Class II-IV heart failure, or transient ischemic attack. 3. Any known allergy or intolerance to midazolam or other drugs in the benzodiazepine class, and/or to dabigatran etexilate, and/or to Oral Contraceptives. 4. Recent exposure to live or attenuated vaccines within 28 days of the screening visit; or use of tobacco/nicotine in excess of 5 cigarettes/day.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: The study consists of 6 sequential parts. In the first 3 parts, participants take a single dose of each medicine (midazolam, dabigatran, and birth control pills) alone. In the last 3 parts, participants take PF-08049820 twice a day, along with one of the other medicines, to assess PK interaction. **Route of Administration:** Oral **Duration of Treatment:** The entire study takes approximately 12 to 13 weeks, including an overnight clinic confinement of about 25 days.

5.2 Concomitant & Prohibited Therapy Permitted: Standard standardized diet and fluids as directed during clinical confinement. **Prohibited:** Use of certain prescription or nonprescription drugs, dietary supplements (e.g., grapefruit), and herbal supplements within up to 14 days or 5 half-lives (whichever is longer) prior to the first dose of study intervention.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening	Day -28 to Day -1 Informed

Consent, Medical History, Physical Exam, Labs, Cardiac Monitoring | | **Baseline & Confinement** | Day 1 to ~Day 25 | Clinic admission, Baseline PK Labs, First Doses | | **Interim** | Over 25 Days | Sequential administration of Midazolam, Dabigatran, Portia alone, followed by PF-08049820 BID co-administration. Continuous PK blood draws and Safety Labs | | **End of Study** | Week 12 to 13 | Final Vital Signs, Exit Interview, IP Accountability, Final Safety Follow-up |

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Changes in Single-Dose Pharmacokinetics (PK) parameters (including Maximum Observed Concentration [C_{max}] and Area Under the Curve [AUC] of Midazolam, and Area Under the Curve [AUC] of Dabigatran) when administered alone versus when co-administered with steady-state PF-08049820. **Measurement Method:** Serial Plasma/Blood Sampling and Central Laboratory Bioanalytical Analysis.

7.2 Secondary & Exploratory Endpoints 1. Incidence and severity of Treatment Emergent Adverse Events (AEs) and Serious Adverse Events (SAEs). 2. Changes in clinical laboratory assessments and vital signs from baseline.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Direct observation and collection via clinical staff during the 25-day inpatient clinic stay, followed by outpatient safety assessments. **Serious Adverse Events (SAEs):** Must be reported within 24 hours of investigator awareness. **Data Monitoring Committee (DMC):** Internal safety review team overseen by the Sponsor (Pfizer).

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: PK Concentration Population (participants with sufficient concentration data) and Safety Set (all participants receiving at least one dose of study medication). **Sample Size Justification:** Sample size is based on standard Phase 1 Drug-Drug Interaction (DDI) study guidelines to provide adequate precision for estimating the ratio of geometric means for PK parameters. **Handling of Missing Data:** PK parameters will not be imputed for missing concentration data; standard Pfizer Phase 1 rules for missing data will apply.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Continuous onsite monitoring during the 25-day confinement period, supplemented by clinical research associate (CRA) site visits. **Audit Trail:** Complete electronic Case Report Form (eCRF) entry, laboratory data reconciliation, and data clarification forms.

11. Study Identifiers & Governance

Primary ID: C6231004 **Posting ID:** 576402166824594150 **Registry Number:** NCT07190430 **Sponsor/Lead Organization:** Pfizer **Company:** Pfizer **Study Title:** PF-08049820 with Midazolam, Dabigatran, Birth Control **Official Regulatory Title:** A Phase 1, Open-label, Fixed Sequence Study to Evaluate The Effect of Steady-State Dosing of PF-08049820 on the Single Dose Pharmacokinetics of Oral Contraceptives, Midazolam, and Dabigatran in Healthy Adult Female Participants

12. Clinical Framework (Drug-Drug Interaction Specific)

Primary Condition: Healthy Volunteers (Investigational drug targeted for Atopic Dermatitis) **Diagnosis Threshold:** Overtly healthy; BMI 16-32 kg/m² and ≤ 50 kg **Medical Methodology:** Pharmacokinetic Drug Interaction Study **Optimization Target:** Characterizing CYP and transporter interaction profiles for PF-08049820

13. Study Procedures & Methodology

Management Pathway: 6-part fixed-sequence clinical confinement **Education Protocol (HCP):** Protocol training, rigorous PK sampling schedule adherence, and AE monitoring **Education Protocol (Patient):** Informed consent, clinic confinement instructions, and strict dietary/medication restrictions **Quality Audit Mechanism:** Direct observation of dosing and PK sampling times during the 25-day inpatient stay

14. Observation & Follow-Up Schedule

Total Duration: 12 to 13 weeks **Onsite Visit Cadence:** ~25 consecutive days of overnight inpatient confinement **Remote Monitoring Frequency:** Outpatient safety follow-ups post-discharge **Checklist Review Interval:** Daily review of safety and vital signs during the confinement phase

15. Sign-off & Approval

	Role		Name		Signature		Date			:---		:---		:---		:---	
	Principal Investigator		[Insert Name]		_____		[DD-MMM-YYYY]										
	Clinical Operations Lead		[Insert Name]		_____		[DD-MMM-YYYY]										
		Lead Statistician		[Insert Name]		_____		[DD-MMM-YYYY]									