

1. Administrative & Study Identification

Full Study Title: Long-term Safety and Efficacy Extension Study for Participants With Advanced Tumors Who Are Currently on Treatment or in Follow-up in a Pembrolizumab (MK-3475) Study (MK-3475-587/KEYNOTE-587) **Official Title:** A Multicenter, Open-label, Phase 3 Study to Evaluate the Long-term Safety and Efficacy in Participants Who Are Currently on Treatment or in Follow-up in Studies That Include Pembrolizumab **Short Title/ Acronym:** KEYNOTE-587 / MK-3475-587 **Protocol Number:** MK-3475-587 **NCT Number:** NCT03486873 **Posting ID:** 594153067731503040 **Sponsor / Company Name:** Merck (Merck Sharp & Dohme LLC) **Investigational Product:** Pembrolizumab (MK-3475), pembrolizumab-based combinations, or lenvatinib **Phase of Development:** PHASE3 **Trial Status:** RECRUITING (Currently recruiting participants) **Medical Monitor:** Merck Sharp & Dohme LLC Medical Monitor

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the long-term safety and efficacy of pembrolizumab (MK-3475) in participants from previous Merck pembrolizumab-based parent studies who transition into this extension study.

2.2 Background & Rationale To provide continuous access to pembrolizumab and enable long-term follow-up for patients with advanced solid tumors and hematologic malignancies who participated in previous pembrolizumab parent trials and who are currently on treatment or in a follow-up phase.

This study generates critical real-world and long-term clinical evidence to confirm that immune checkpoint blockade provides durable, long-term survival benefits and manageable safety profiles.

2.3 Study Phases & Crossover The study consists of three phases depending on the participant's status at the parent study's completion: 1. **First Course Phase:** For participants on active treatment in the parent study. 2. **Survival Follow-up Phase:** For participants who were in follow-up (post-treatment or survival) in the parent study. 3. **Second Course Phase:** For participants in the Second Course Phase of their parent study.

Crossover Rules: Any participant originating from a parent trial where crossover to pembrolizumab was permitted upon disease progression may be eligible for 35 doses (Q3W) or 17 doses (Q6W) of pembrolizumab (approximately 2 years) if they progress while

on the control arm and pembrolizumab is approved for the indication in their country.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional (Extension Trial)
Control Type: Single-arm / Open-label Extension (incorporates parent-study treatment arms) **Blinding:** Open-label **Randomization:** N/A (Participants roll over directly from parent studies)

3.2 Planned Enrollment & Duration Targeted Enrollment: 3,500 participants **Number of Sites:** Multicenter (Global) **Disease Areas:** Solid Tumors, Hematologic Malignancies

4. Subject Selection (Eligibility Criteria)

General Exclusion: There are no exclusion criteria to participate in the general MK-3475-587 extension study itself. Specific criteria apply to entering the Second Course Phase or receiving Lenvatinib.

4.1 General Inclusion Criteria 1. Treated on a parent pembrolizumab study established by the Sponsor as MK-3475-587 ready. 2. Currently receiving pembrolizumab, pembrolizumab-based combinations, or lenvatinib from parent studies, or in a follow-up phase.

4.2 Additional Criteria for Second Course Phase Inclusion: 1. Has not received any anticancer systemic treatment since the last dose of pembrolizumab or pembrolizumab-based combination in First Course. 2. Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1. 3. Demonstrates adequate organ function. 4. Resolution of toxic effects of First Course Phase to Grade ≤ 1 (except alopecia). Recovered from toxicity/complications of major surgery or radiation (>30 Gy). 5. Women of childbearing potential (WOCBP) must not be pregnant/breastfeeding and agree to use contraception during the study treatment period and for ≥ 120 days after.

Exclusion: 1. Severe hypersensitivity ($\geq$ Grade 3) to pembrolizumab and/or excipients. 2. Received a live vaccine within 30 days prior. 3. Immunodeficiency or chronic systemic steroid therapy (>10 mg prednisone equivalent) within 7 days prior to Cycle 1 Day 1. 4. Additional progressing malignancy requiring active treatment (exceptions apply). 5. Active CNS metastases and/or carcinomatous meningitis. 6. Active autoimmune disease requiring systemic treatment in the past 2 years (replacement therapy allowed). 7. History of (non-infectious) pneumonitis that required steroids or current pneumonitis (unless it occurred during First Course and did not meet permanent discontinuation criteria). 8. Non-small cell lung cancer (NSCLC)

participants: interstitial lung disease. 9. Active infection requiring systemic therapy, known HIV, active TB, or known active Hepatitis B/C (unless permitted by parent study). 10. Severe cardiovascular disease (NYHA Class III/IV, symptomatic ischemia, arrhythmias requiring chronic treatment). 11. Hepatic decompensation (Child-Pugh score >6 [class B and C]). 12. Uncontrolled thyroid dysfunction or uncontrolled diabetes mellitus. 13. Prior allogeneic tissue/solid organ transplant.

4.3 Additional Criteria for Lenvatinib Dosing Inclusion: 1. Adequately controlled blood pressure (BP <150/90 mmHg). 2. Males must be abstinent or use highly effective contraception during and for 7 days post-treatment. 3. Females must use highly effective contraception or be abstinent during and for $\geq$ 4 days post-treatment.

Exclusion: 1. Major surgery within 3 weeks prior to the first dose. 2. Preexisting $\geq$ Grade 3 gastrointestinal or non-gastrointestinal fistula. 3. Urine protein $\geq$ 1 g/24 hours. 4. LVEF below institutional normal range. 5. Radiographic evidence of encasement/invasion of a major blood vessel or intratumoral cavitation. 6. QTcF interval prolongation >480 ms. 7. Clinically significant cardiovascular disease within 12 months. 8. Gastrointestinal malabsorption. 9. Active hemoptysis (at least 0.5 teaspoon bright red blood) within 3 weeks. 10. History of contraindication/severe hypersensitivity to lenvatinib.

5. Intervention & Treatment Plan

5.1 Investigational Regimen First Course Phase: Pembrolizumab or lenvatinib up to 35 doses or more every 3 weeks (Q3W), or 17 doses or more every 6 weeks (Q6W). **Second Course Phase:** Pembrolizumab up to 17 doses Q3W or 8 doses Q6W. **Route of Administration:** Intravenous (IV) for pembrolizumab; Oral for lenvatinib.

5.2 Concomitant & Prohibited Therapy Permitted: Routine supportive care and replacement therapy (e.g., thyroxine, insulin, physiologic corticosteroid replacement). **Prohibited:** Systemic immunosuppressive steroids (except replacement or managing imAEs), live vaccines.

6. Outcome Measures (Endpoints)

6.1 Primary Outcomes 1. Overall Survival (OS): Defined as the time from randomization or start of study treatment for non-randomized participants (on the parent study) to death due to any cause. Participants without documented death at the time of analysis will be censored at the date of the last known to be alive.

6.2 Secondary Outcomes 1. **Modified Progression Free Survival (PFS)** Per Evaluation Criteria Used in the Parent Trial. 2. **Modified Event Free Survival (EFS)** Per Evaluation Criteria Used in the Parent Trial. 3. **Number of Participants Who Experience Serious Adverse Events (SAEs).**

7. Database Metadata & Disease Classifications

To support cross-trial data aggregation, the following indexing criteria apply to this record: * **Preferred UMLS Name:** Pembrolizumab Therapy Follow-up * **Condition / Disease Name:** Solid Tumor (Preferred Name: Solid Neoplasm) * **Semantic Type:** Neoplastic Process * **Definition:** A benign or malignant neoplasm arising from tissues that do not include fluid areas. Representative examples include epithelial neoplasms (e.g. lung carcinoma, prostate carcinoma, breast carcinoma, colon carcinoma), and neoplasms arising from the soft tissues and bones (e.g. leiomyosarcoma, liposarcoma, chondrosarcoma, osteosarcoma). Neoplasms originating from the blood or bone marrow (leukemias and myeloproliferative disorders) are not considered solid tumors. * **MeSH Code:** C04.588.550 * **SNOMED ID:** 126939001 * **SNOMED Hierarchy:** 93855008|Solid tumor|Neoplasm (disorder) * **Source Level:** 5 * **Criteria Type:** PHASE_REQUIREMENT (Criteria Text: Trial must be in PHASE3; Phase ID: PHASE3) * **Status Name / Description:** RECRUITING / Currently recruiting participants * **Associated Drug Terminology:** (Indexed Drug Record) * **Drug Name / Preferred Name:** olaparib * **Trade Names:** Lynparza * **ATC Code:** L01XK01 * **EMA URL:** https://www.ema.europa.eu/en/documents/product-information/lynparza-epar-product-information_en.pdf

8. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.
Monitoring Plan: Standard site monitoring for oncology extension phases to ensure protocol adherence and safety tracking.

9. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Insert Name]	_____	06-Apr-2026				
Clinical Operations Lead	[Insert Name]	_____	06-Apr-2026				
Lead Statistician	[Insert Name]	_____	06-Apr-2026				