

Clinical Study Summary Report (CSSR)

Document Status: Final | **Version:** 1.0 | **Date:** 07-Apr-2026 | **Trial Status:** COMPLETED

1. Administrative & Study Identification

Full Study Title: A 12-month, Open-label, Prospective, Multicenter, Interventional, Single-arm Study Assessing the Efficacy and Safety of Ofatumumab 20 mg Subcutaneous (s.c.) Injection in Relapsing Multiple Sclerosis (RMS) Patients in China
Short Title/Acronym: OMB157 / Ofatumumab in RMS Patients in China
Protocol Number: COMB157G2402 **NCT Number:** NCT05199571 **Sponsor Name:** Novartis **Investigational Product:** Ofatumumab 20 mg (OMB157)
Phase of Development: Phase 4 **Medical Monitor:** Novartis Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the efficacy and safety of ofatumumab administered subcutaneously (s.c.) in adult participants with relapsing multiple sclerosis (RMS) in China by measuring the annualized relapse rate (ARR). **Secondary Objectives:** To evaluate the safety and tolerability of the treatment, the annualized rate of new or enlarged T2 lesions, the number of gadolinium (Gd)-enhancing T1 lesions on MRI, and disability progression/improvement.

2.2 Background & Rationale To provide real-world and clinical trial evidence on the safety and effectiveness of ofatumumab—a fully human anti-CD20 monoclonal antibody that targets B-cell depletion—in Chinese patients with RMS. This study aims to consolidate therapeutic options and limit disease progression, brain lesions, and physical disability through a convenient, at-home, once-monthly subcutaneous administration.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: 99 participants
Number of Sites: Multicenter (Approximately 10 sites in China, including locations in Tianjin, Guangzhou, Wuhan, and Shanghai)
Estimated Study Start: July 2022 **Trial Status:** COMPLETED

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Male or female Chinese participants aged 18 to 55 years (inclusive) at enrollment. 2. Clinical definite diagnosis of RMS according to the 2017 Revised McDonald criteria. 3. Documentation of two relapses during the past 2 years, OR one documented relapse during the last year, OR a positive Gd-enhancing MRI scan during the year prior to screening. 4. Disability status with an EDSS score of 0 - 5.5 (inclusive) at Screening. 5. Neurologically stable within 1 month prior to both Screening and Baseline (including no MS relapse in this period).

4.2 Exclusion Criteria 1. Participants with primary progressive MS (PPMS) or secondary progressive MS (SPMS) without disease activity, or meeting criteria for neuromyelitis optica spectrum disorder (NMOSD). 2. Pregnant or nursing (lactating) women; women of child-bearing potential unless using effective methods of contraception while taking study treatment and for at least 6 months after stopping medication. 3. Participants with an active chronic disease of the immune system other than MS. 4. Active systemic infections (including but not limited to active COVID-19 infection) or known to have AIDS or test positive for HIV antibody at Screening; active hepatitis B disease. 5. Participants with neurological findings consistent with PML or confirmed PML. 6. Participants at high risk of developing or having reactivation of syphilis or tuberculosis. 7. Receipt of any live or live-attenuated vaccines within 4 weeks prior to first study drug administration, or treated with other specified medications within protocol timeframes.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Ofatumumab 20 mg administered at Baseline/Week 0, Week 1, and Week 2, followed by subsequent monthly dosing starting at Week 4. **Route of Administration:** Subcutaneous (s.c.) injection (first dose administered in the clinic; subsequent doses self-administered at home). **Duration of Treatment:** 12 months (followed by a 6-month post-treatment follow-up).

5.2 Concomitant & Prohibited Therapy Permitted: Standard symptomatic management for MS. **Prohibited:** Live or live-attenuated vaccines within four weeks prior to the first dose; concurrent use of other disease-modifying therapies (DMTs) or immunosuppressants targeting MS during the study period.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening	Up to 30 days (Day -30 to 0)

Informed Consent, Medical History, Baseline MRI, clinical evaluation | | **Baseline** | Day 0 | Clinical evaluation, first ofatumumab 20 mg s.c. injection in clinic | | **Interim Treatment** | Weeks 1, 2, 4, then monthly | 5 clinic visits during the Treatment period for clinical evaluation and lab tests; monthly S.c. injections | | **End of Study / Follow-up** | Month 12 to 18 | Post treatment Follow-Up visits at End of Study plus 3 months (EOS + 3M) and EOS + 6M, unless continuing commercial ofatumumab. |

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Adjusted Annualized Relapse Rate (ARR) Based on Confirmed Relapses. A confirmed MS relapse is defined as one accompanied by an EDSS increase of at least 0.5 points, or an increase of 1 point on two functional scores, or 2 points on one functional score (excluding changes involving bowel/bladder or cerebral functional system) assessed by an Independent EDSS rater. **Measurement Method:** Adjusted ARR obtained from fitting a negative binomial regression model adjusted for number of relapses in the previous year, baseline number of T1 Gd-enhancing lesions and baseline age as continuous covariates (offset: Natural log of time in study in years).

7.2 Secondary & Exploratory Endpoints * Annualized Rate of New or Enlarging T2 Lesions. * Number of Gadolinium (Gd)-Enhancing T1 Lesions Per MRI Scan. * Number of Adverse Events and Serious Adverse Events.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring of general health and medical problems via clinical evaluations during the 5 treatment clinic visits and at EOS + 3M and EOS + 6M. **Serious Adverse Events (SAEs):** Standard 24-hour reporting timeline to the sponsor/pharmacovigilance team. **Data Monitoring Committee (DMC):** Internal Novartis review committee.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Full Analysis Set (FAS), following the Intent-to-Treat (ITT) principle. **Sample Size Justification:** Approximately 99 participants to provide adequate power for estimating ARR and MRI lesion reductions in the Chinese RMS subpopulation. **Handling of Missing Data:** For the primary analysis, Adjusted ARR is estimated using a negative binomial regression model containing individual cumulative confirmed relapse counts as the response variable, with participants' time in the study modeled as an offset variable (Natural log of time in study in years).

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: On-site monitoring during 5 scheduled clinic visits throughout the treatment period, supplemented by safety follow-ups. **Audit Trail:** eCRF locking and imaging data central reading (for MRI evaluations) to ensure data integrity and clarification.

11. Study Identifiers & Governance

Primary ID: COMB157G2402 **Registry Number:** NCT05199571 **Sponsor/Lead Organization:** Novartis **Study Title:** Study of Efficacy and Safety of Ofatumumab in Relapsing Multiple Sclerosis (RMS) Patients in China **Official Regulatory Title:** A 12-month, Open-label, Prospective, Multicenter, Interventional, Single-arm Study Assessing the Efficacy and Safety of Ofatumumab 20 mg Subcutaneous (s.c.) Injection in Relapsing Multiple Sclerosis (RMS) Patients in China

12. Clinical Framework (MS Specific)

Primary Condition: Relapsing Multiple Sclerosis (RMS) **Diagnosis Threshold:** 2017 Revised McDonald Criteria **Medical Methodology:** B-cell targeted therapy (anti-CD20 monoclonal antibody)

Optimization Target: Disease-Modifying Therapy (DMT) efficacy to minimize relapses and MRI lesion burden

13. Study Procedures & Methodology

Management Pathway: Standard RMS care pathway integrated with subcutaneous CD20 inhibition **Education Protocol (HCP):** Site initiation visits and training on the recognition of confirmed MS relapses and EDSS scoring **Education Protocol (Patient):** Onsite training for the first dose injection; self-learning and guided instruction for subsequent at-home self-administration **Quality Audit Mechanism:** Independent EDSS rater evaluations and centralized MRI reading

14. Observation & Follow-Up Schedule

Total Duration: 18 months (12 months treatment + 6 months follow-up) **Onsite Visit Cadence:** Screening, Baseline, 5 clinic visits during treatment, EOS + 3M, and EOS + 6M **Remote Monitoring Frequency:** Routine touchpoints for at-home administration

tracking **Checklist Review Interval:** Clinic visits to review AEs and administration logs

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				