

## **1. Administrative & Study Identification**

**Full Study Title:** A Study to Learn About the Study Medicine Elranatamab Alone and With Daratumumab in People With Multiple Myeloma Who Have Received Other Treatments **Short Title/Acronym:** MagnetisMM-5 **Protocol Number:** C1071005 (PF-06863135) **NCT Number:** NCT05020236 **Trial Status:** RECRUITING **Sponsor/Company Name:** Pfizer **Investigational Product:** Elranatamab (PF-06863135), Daratumumab, Pomalidomide, Dexamethasone **Phase of Development:** Phase 3 **Disease Area:** Multiple Myeloma **Medical Monitor:** Pfizer Clinical Operations Lead

---

## **2. Study Synopsis & Rationale**

**2.1 Study Objective Primary Objective:** To evaluate the efficacy and safety of elranatamab monotherapy and elranatamab in combination with daratumumab compared to the active comparator combination of daratumumab, pomalidomide, and dexamethasone in participants with relapsed/refractory multiple myeloma (RRMM). **Secondary Objectives:** To evaluate overall survival, PFS on next-line treatment, objective response rate (ORR), duration of response, complete response (CR) rate, overall and sustained minimal residual disease (MRD) negativity rates, safety, quality of life, immunogenicity, and pharmacokinetics.

**2.2 Background & Rationale** Patients with relapsed/refractory multiple myeloma who have already received previous standard treatments, including lenalidomide and a proteasome inhibitor, need new, efficacious treatment options. Elranatamab is a humanized BCMA-CD3 bispecific antibody that targets BCMA on myeloma cells and CD3 on T cells. Combining it with daratumumab (an anti-CD38 monoclonal antibody) is supported by proposed dual mechanisms of action that could further enhance anti-tumor responses and provide greater clinical benefit compared to current standard regimens.

---

## **3. Study Design & Methodology**

**3.1 Design Framework Study Type:** Interventional **Control Type:** Active Comparator (Daratumumab + Pomalidomide + Dexamethasone) **Blinding:** Open-label **Randomization:** 1:1:1 (in Part 2 of the study)

**3.2 Planned Enrollment & Duration Targeted Enrollment:** 944 patients **Number of Sites:** Multicenter, global (United States,

## 4. Subject Selection (Eligibility Criteria)

**4.1 Inclusion Criteria** 1. Prior diagnosis of multiple myeloma as defined by IMWG criteria (Rajkumar et al, 2014). 2. Measurable disease based on IMWG criteria as defined by at least 1 of the following: - Serum M-protein  $\geq 0.5$  g/dL. - Urinary M-protein excretion  $\geq 200$  mg/24 hours. - Serum immunoglobulin FLC  $\geq 10$  mg/dL ( $\geq 100$  mg/L) AND abnormal serum immunoglobulin kappa to lambda FLC ratio ( $<0.26$  or  $>1.65$ ). 3. Prior anti-multiple myeloma therapy including treatment with lenalidomide and a proteasome inhibitor. 4. ECOG performance status  $\leq 2$ . 5. Resolved acute effects of any prior therapy to baseline severity or CTCAE Grade  $\leq 1$ . 6. Not pregnant and willing to use contraception.

**4.2 Exclusion Criteria** 1. Smoldering multiple myeloma, plasma cell leukemia, amyloidosis, or POEMS Syndrome. 2. Stem cell transplant within 12 weeks prior to enrolment, or active graft-versus-host disease. 3. Previous treatment with a BCMA-directed therapy. 4. Active HBV, HCV, SARS-CoV2, HIV, or any active, uncontrolled bacterial, fungal, or viral infection. 5. Any other active malignancy within 3 years prior to enrolment, except for adequately treated basal cell or squamous cell skin cancer, or carcinoma in situ. 6. Live attenuated vaccine within 4 weeks of the first dose of study intervention. 7. Administration with an investigational product (e.g., drug or vaccine) concurrent with study intervention or within 30 days preceding the first dose of study intervention used in this study.

---

## 5. Intervention & Treatment Plan

**5.1 Investigational Regimen Dosage/Frequency:** Administered based on cycle schedules until disease progression, unacceptable side effects, or withdrawal of consent. Elranatamab is given as a subcutaneous injection alone or with subcutaneous Daratumumab. The Active Comparator arm receives subcutaneous Daratumumab with oral Pomalidomide and Dexamethasone. **Route of Administration:** Subcutaneous (Elranatamab, Daratumumab); Oral (Pomalidomide, Dexamethasone). **Duration of Treatment:** Until disease progression, unacceptable side effects, or withdrawal of consent.

**5.2 Concomitant & Prohibited Therapy Permitted:** Standard concomitant medications for condition management and mitigation of adverse effects (e.g., infection prophylaxis). **Prohibited:** Live attenuated vaccines within 4 weeks of the first dose; other investigational products within 30 days preceding the first dose; concurrent anti-myeloma therapies outside the assigned arm.

---

## 6. Study Timeline & Visit Schedule

| Visit ID       | Window (Days) | Key Activities/Procedures                                                                    |
|----------------|---------------|----------------------------------------------------------------------------------------------|
| Screening      | Days -28 to 0 | Informed Consent, Medical History, Confirmation of IMWG measurable disease                   |
| Baseline       | Day 0         | Randomization (1:1:1), First Dose Administration, Safety Baseline                            |
| Interim Cycles | Cycle-based   | Administration of Study Drugs, Safety Labs, Efficacy Assessment (IMWG criteria), QoL surveys |
| End of Study   | Variable      | Disease progression verification, Exit Interview, Safety follow-up                           |

---

## 7. Outcome Measures (Endpoints)

**7.1 Primary Endpoints**

- Part 1 Safety Lead-In:** Incidence of dose limiting toxicities.
- Part 2 Randomized:** Progression free survival per International Myeloma Working Group (IMWG) criteria.
- Part 3:** Frequency of treatment-emergent adverse events.

**7.2 Secondary & Exploratory Endpoints**

- Part 1 Safety Lead-In:** Progression free survival per International Myeloma Working Group (IMWG) criteria.
- Overall survival (OS).
- Objective response rate (ORR) per International Myeloma Working Group (IMWG) criteria.
- Duration of Response and PFS on next-line treatment.
- Complete Response (CR) rate and overall and sustained Minimal Residual Disease (MRD) negativity rates.

---

## 8. Safety & Adverse Event Reporting

**Adverse Event (AE) Monitoring:** Continuous monitoring of clinical symptoms, particularly focusing on infection rates, frequency of treatment-emergent adverse events, cytokine release syndrome (CRS), and neurologic toxicities typical of bispecific antibodies.

**Serious Adverse Events (SAEs):** Standard 24-hour reporting protocols for severe toxicities.

**Data Monitoring Committee (DMC):** Independent safety monitoring board evaluating dose-escalation toxicities (Part 1) and ongoing Phase 3 safety metrics (Parts 2 & 3).

---

## 9. Statistical Analysis Plan (SAP) Summary

**Analysis Populations:** Intent-to-Treat (ITT) Population for primary efficacy endpoints; Safety Analysis Set for patients who received at least one dose of study drug.

**Sample Size Justification:** Total targeted enrollment of 944 patients powered to detect statistically significant improvements in PFS for elranatamab-containing arms versus the DPd standard-of-care arm.

**Handling of Missing Data:** Standard right-censoring for time-to-

event endpoints (PFS/OS); non-responder imputation for categorical response endpoints (ORR/CR).

---

## 10. Quality Assurance & Regulatory Compliance

**GCP Compliance:** This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

**Monitoring Plan:** Standard risk-based onsite and remote monitoring of participating global centers. **Audit Trail:** Validated Electronic Data Capture (EDC) system utilization with comprehensive data clarification and audit tracking.

---

## 11. Study Identifiers & Governance

**Primary ID:** C1071005 **Registry Number:** NCT05020236 **Posting ID:** 693398384424854136 **Sponsor/Lead Organization:** Pfizer **Study Title:** MagnetisMM-5 **Official Regulatory Title:** AN OPEN-LABEL, 3-ARM, MULTICENTER, RANDOMIZED PHASE 3 STUDY TO EVALUATE THE EFFICACY AND SAFETY OF ELRANATAMAB (PF-06863135) MONOTHERAPY AND ELRANATAMAB + DARATUMUMAB VERSUS DARATUMUMAB + POMALIDOMIDE + DEXAMETHASONE IN PARTICIPANTS WITH RELAPSED/REFRACTORY MULTIPLE MYELOMA WHO HAVE RECEIVED AT LEAST 1 PRIOR LINE OF THERAPY INCLUDING LENALIDOMIDE AND A PROTEASOME INHIBITOR

---

## 12. Clinical Framework (Multiple Myeloma Specific)

**Primary Condition:** Relapsed/Refractory Multiple Myeloma (RRMM)

**Diagnosis Threshold:** IMWG criteria with measurable disease (e.g., Serum M-protein  $\geq 0.5$  g/dL) **Medical Methodology:** BCMA-CD3 bispecific antibody therapy alone or combined with CD38-directed monoclonal antibody **Optimization Target:** Progression-Free Survival (PFS) and Disease Response optimization

---

## 13. Study Procedures & Methodology

**Management Pathway:** Continuous active systemic therapy until disease progression or unacceptable toxicity

**Education Protocol (HCP):** Specialized training on the administration of subcutaneous bispecifics and the management of distinct immune-related adverse events **Education Protocol (Patient):** Comprehensive instruction on identifying symptoms of infection, CRS, and routine self-monitoring procedures **Quality Audit Mechanism:** Blinded independent central review (BICR) for primary efficacy assessments and stringent clinical site monitoring

---

## 14. Observation & Follow-Up Schedule

**Total Duration:** Follow-up continues until disease progression, death, or study withdrawal **Onsite Visit Cadence:** Standardized according to typical 28-day treatment cycles **Remote Monitoring Frequency:** Inter-cycle continuous safety tracking **Checklist Review Interval:** Regular disease staging and serologic testing aligned with IMWG response criteria

## 15. Sign-off & Approval

| Role                     | Name   | Signature | Date          |  |  | :--- | :--- | :--- | :--- |
|--------------------------|--------|-----------|---------------|--|--|------|------|------|------|
| Principal Investigator   | [Name] | _____     | [DD-MMM-YYYY] |  |  |      |      |      |      |
| Clinical Operations Lead | [Name] | _____     | [DD-MMM-YYYY] |  |  |      |      |      | Lead |
| Statistician             | [Name] | _____     | [DD-MMM-YYYY] |  |  |      |      |      |      |