

1. Administrative & Study Identification

Full Study Title: An Observational Study of Etrasimod in Adult Patients With Moderate to Severe Ulcerative Colitis **Short Title/ Acronym:** ENDEAVOUR-UC **Protocol Number:** C5041047 **NCT Number:** NCT06398626 **Sponsor Name:** Pfizer **Investigational Product:** Etrasimod (Velsipity®) **Phase of Development:** Phase IV / Observational (Real-World Effectiveness) **Medical Monitor:** Pfizer Clinical Operations / Lead Principal Investigator

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To observe the real-world effectiveness of etrasimod as a treatment for adult patients with moderate to severe ulcerative colitis (UC) by evaluating the proportion of patients achieving symptomatic remission at Week 12. **Secondary Objectives:** To evaluate symptomatic remission, steroid-free symptomatic remission, and steroid-free clinical remission at Weeks 24, 36, and 52 using the partial modified Mayo score (pMMS). To assess symptomatic response, clinical remission, and changes in fatigue, bowel urgency, and abdominal pain at Weeks 12, 24, 36, and 52.

2.2 Background & Rationale High-quality observational studies remain vital to understand the real-world (RW) experience of etrasimod outside of a controlled trial setting. The study aims to provide robust prospective effectiveness data to see how this S1P receptor modulator performs when prescribed to adult patients in routine clinical care.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational (Cohort) **Control Type:** Single-arm (Non-interventional) **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: ~250-300 estimated patients **Number of Sites:** Approximately 30 US sites **Estimated Study Start:** 17-Sep-2024 **Estimated Primary Completion:** 01-Oct-2027

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Age ≥ 18 and < 65 years at baseline. 2. Confirmed diagnosis of ulcerative colitis (UC) with

moderately to severely active disease. 3. Prescribed etrasimod as part of routine clinical care, independent from the decision to enroll the patient in this study.

4.2 Exclusion Criteria 1. The presence of clinical findings suggestive of Crohn's disease. 2. Any previous exposure to etrasimod, including participation in the etrasimod clinical program. 3. Concomitant use of conventional therapies (e.g., methotrexate, or thiopurines such as azathioprine, mercaptopurine, tioguanine) or advanced therapies (biologics, JAK inhibitors like filgotinib, tofacitinib, or upadacitinib, or any other S1P receptor modulator).

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Prescribed etrasimod according to the standard of care instructions. **Route of Administration:** Oral **Duration of Treatment:** 52 weeks of follow-up from etrasimod initiation.

5.2 Concomitant & Prohibited Therapy Permitted: Standard of care concurrent medications for UC that do not violate exclusion criteria. **Prohibited:** Co-medication with methotrexate, thiopurines, biologics (TNF, integrin or cytokine antagonists), JAK inhibitors, or other S1P receptor modulators.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
1	Screening/Baseline	Day 0 Informed Consent, Patient & Disease Characteristics, ePRO setup
2	Daily Phase	Days 1-14 Daily PRO data collection via web-based application
3	Interim	Weeks 4, 8, 12, 24, 36 Symptom Questionnaires via web app; Endoscopy, Fecal Calprotectin, & CBC (if part of standard of care)
4	End of Study	Week 52 Final Questionnaires, Endpoint Analysis, followed by a 28-day safety follow-up

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Proportion of patients with symptomatic remission at Week 12. Symptomatic remission is defined as a partial Modified Mayo Score (pMMS) stool frequency subscore (SFS) = 0 (or = 1 with a ≥ 1 -point decrease from baseline) and rectal bleeding subscore (RBS) = 0. **Measurement Method:** Patient-reported outcome (PRO) data collected via a web-based application.

7.2 Secondary & Exploratory Endpoints * Symptomatic remission, steroid-free symptomatic remission, and steroid-free clinical

remission at Weeks 24, 36, and 52 utilizing the pMMS. *
Symptomatic response, clinical remission, and changes in fatigue, bowel urgency, and abdominal pain at Weeks 12, 24, 36, and 52.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Conducted under standard of care clinical observations across the 52 weeks and evaluated during an additional 28-day safety follow-up period. **Serious Adverse Events (SAEs):** Standard reporting timelines within 24 hours of site awareness. **Data Monitoring Committee (DMC):** Governed by internal sponsor safety monitoring parameters for observational registries.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Real-world cohort based on treatment exposure lines (AT-naïve vs. AT-exposed). **Sample Size Justification:** The sample size of 250–300 patients was informed by ELEVATE UC phase 3 randomized controlled trial data, assuming a 40% Week 12 symptomatic remission rate to achieve a 6.1% precision level (half-width of 95% confidence interval). **Handling of Missing Data:** Standard approaches (e.g., LOCF or Multiple Imputation) utilized for missing patient-reported data points where appropriate.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP), real-world evidence guidelines, and the Declaration of Helsinki. **Monitoring Plan:** Remote data monitoring of the web-based patient-reported outcome application; additional metrics (endoscopy, fecal calprotectin) collected only if performed as standard of care. **Audit Trail:** Robust electronic records and data clarification forms supporting the web-based PRO application entries.

11. Study Identifiers & Governance

Primary ID: C5041047 **Registry Number:** NCT06398626 **Sponsor/Lead Organization:** Pfizer **Study Title:** ENDEAVOUR-UC **Official Regulatory Title:** An Observational Study of Etrasimod in Adult Patients With Moderate to Severe Ulcerative Colitis

Primary Condition: Moderately to Severely Active Ulcerative Colitis (UC) **Diagnosis Threshold:** Confirmed diagnosis and clinically prescribed etrasimod **Medical Methodology:** Routine clinical care and real-world standard of care observation **Optimization Target:** Symptomatic and steroid-free clinical remission

Management Pathway: Standard clinical pathway for UC utilizing etrasimod **Education Protocol (HCP):** N/A (Non-interventional observational design) **Education Protocol (Patient):** Self-reporting methodology using a web-based ePRO application **Quality Audit Mechanism:** Real-time completion tracking of the daily and scheduled digital questionnaires

Total Duration: 52 follow-up weeks + 28-day safety follow-up
Onsite Visit Cadence: Dictated entirely by patient's standard of care routine
Remote Monitoring Frequency: Web-app questionnaire entries daily for the first 2 weeks, then at Weeks 4, 8, 12, 24, 36, and 52
Checklist Review Interval: Continuous digital review corresponding with interim weeks

Role	Name	Signature	Date	_____	_____	_____	_____
Principal Investigator	[Insert Name]	_____	[DD-MMM-YYYY]	_____	_____	_____	_____
Clinical Operations Lead	[Insert Name]	_____	[DD-MMM-YYYY]	_____	_____	_____	_____
Lead Statistician	[Insert Name]	_____	[DD-MMM-YYYY]	_____	_____	_____	_____