

1. Administrative & Study Identification

Full Study Title: Long-term, Interventional, Open Label Extension Study Evaluating the Safety of Tocilizumab Treatment in Patients With Polyarticular-course Juvenile Idiopathic Arthritis From France Who Completed the Global, Multinational Trial (WA19977)
Short Title/Acronym: A Long-Term Extension Study of RoActemra/Actemra (Tocilizumab) in Patients With Juvenile Idiopathic Arthritis From France Who Completed WA19977 Core Study
Protocol Number: ML28166 (EudraCT Number: 2011-005515-90) **NCT Number:** NCT01673919 **Sponsor Name:** Roche **Investigational Product:** Tocilizumab [RoActemra/Actemra] **Phase of Development:** Phase 3
Trial Status: COMPLETED **Medical Monitor:** Roche Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the long-term safety of RoActemra/Actemra (tocilizumab) in patients with polyarticular-course juvenile idiopathic arthritis (pJIA) who completed the WA19977 core study. **Secondary Objectives:** To evaluate the long-term efficacy, disease activity, and maintenance of clinical responses over the extended treatment period.

2.2 Background & Rationale To provide continued access to tocilizumab and assess the long-term safety and efficacy of the treatment in patients with polyarticular-course JIA who successfully completed the WA19977 core study and continued to derive clinical benefit from the therapy, thereby generating real-world, long-term extension data.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm (Open-label extension) **Blinding:** Open-label
Randomization: N/A

3.2 Planned Enrollment & Duration Target Enrollment: 7 participants **Number of Sites:** Participating sites in France **Trial Status:** COMPLETED **Estimated Study Start:** February 2012 **Estimated Primary Completion:** January 2014

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Patients who completed visit 33 (week 104) of WA19977 study and who may benefit from study drug treatment according to the investigator's assessment. 2. Patients have to receive the first RoActemra/Actemra infusion in this study at the Week 8 visit at the latest. 3. Females of child-bearing potential and males with female partners of child-bearing potential must agree to use effective contraception as defined by protocol.

4.2 Exclusion Criteria 1. Patients with, according to investigator judgment, not satisfactory benefit from RoActemra/Actemra therapy within WA19977. 2. Treatment with any investigational agent since the last administration of study drug in the core study WA19977. 3. Patient developed any other autoimmune rheumatic disease or overlap syndrome other than the permitted polyarticular-course Juvenile Idiopathic Arthritis (JIA) subsets: rheumatoid factor positive or negative JIA or extended oligoarticular JIA. 4. Patient is pregnant, lactating, or intending to become pregnant during the study and up to 12 weeks after the last administration of study drug. 5. Any significant concomitant disease or medical or surgical condition. 6. History of significant allergic or infusion reactions to prior biologic therapy. 7. Currently active primary or secondary immunodeficiency. 8. Any infections with contraindications to RoActemra/Actemra therapy according to investigator judgment. 9. Inadequate hepatic, renal or bone marrow function.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Tocilizumab 8 mg/kg administered every 4 weeks. **Route of Administration:** Intravenous (IV) infusion. **Duration of Treatment:** 104 weeks of anticipated extended study treatment.

5.2 Concomitant & Prohibited Therapy Permitted: Stable doses of concomitant JIA medications (e.g., nonsteroidal anti-inflammatory drugs [NSAIDs], oral corticosteroids, or methotrexate) per standard of care. **Prohibited:** Use of any other investigational agents or non-permitted biologic disease-modifying antirheumatic drugs (bDMARDs) during the extension period.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Baseline	Day 0	Roll-over from WA19977 Core Study, Informed Consent, Vital Signs
Interim	Every 4 Weeks	IV Infusion of Tocilizumab, Safety Labs, Efficacy Assessment (JIA ACR criteria)
End of Study	Week 104	Final Efficacy & Safety Evaluation, IP Accountability, Exit Interview
Follow-		

up | +12 Weeks | Off-treatment safety follow-up for monitoring of SAEs/AEs |

(Note: Standard cadence aligns with the 4-week dosing schedule of the investigational product).

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints 1. Number of Participants With Any Adverse Events and Any Serious Adverse Events:

An adverse event (AE) is defined as any untoward medical occurrence in a participant or clinical investigation participant, temporally associated with the use of a medicinal product, whether or not considered to be related to the medicinal product. A serious adverse event (SAE) is any untoward medical occurrence that, at any dose, results in death, is life threatening, requires hospitalization or prolongation of existing hospitalization, results in disability/incapacity, or results in a congenital anomaly/birth defect.

2. **Number of Participants With Adverse Events of Special Interest:** The AEs of special interests included gingival bleeding, tooth abscess, acarodermatitis, ear infection, gastroenteritis, herpes zoster ophthalmic, lice infestation, nasopharyngitis, oral fungal infection, oral herpes, pharyngitis, rhinitis, sinusitis, tonsillitis, tracheitis, tracheobronchitis, urinary tract infection, menorrhagia, asthma, epistaxis, and hematoma.

3. **Number of Participants With Adverse Events Related to Tocilizumab:** Relatedness of any AEs was reported as possibly related, probably related, or remotely related to TCZ.

7.2 **Secondary Endpoints** 1. Mean Exposure to Study Treatment. 2. Mean Duration of Study Follow-Up. 3. Number of Participants With AEs Leading to TCZ Modification, AEs Leading to Death, Anaphylaxis or Serious Hypersensitivity and Deaths.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: AEs monitored continuously, with special interest events including systemic infections, infusion reactions, neutropenia, elevated transaminases, and gastrointestinal complications. **Serious Adverse Events (SAEs):** Reported within 24 hours of site awareness, defined as events resulting in death, life-threatening conditions, or hospitalizations. **Data Monitoring Committee (DMC):** Internal safety monitoring managed by the sponsor (Roche).

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all enrolled participants who received at least one dose of the investigational product during the extension phase). **Sample Size Justification:** Target $N=7\$$;

driven by the specific cohort of eligible French patients rolling over from the completed WA19977 global core study. **Handling of Missing Data:** Non-responder imputation (NRI) for binary efficacy endpoints and Last Observation Carried Forward (LOCF) for continuous variables where applicable.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Regular clinical site monitoring visits to ensure protocol adherence, accurate data transcription, and stringent AE/SAE reporting. **Audit Trail:** Data securely captured with standard eCRF systems incorporating complete audit trails and data clarification form mechanisms.

11. Study Identifiers & Governance

Primary ID: ML28166 **Registry Number:** NCT01673919 **Sponsor/Lead Organization:** Roche **Study Title:** A Long-Term Extension Study of RoActemra/Actemra (Tocilizumab) in Patients With Juvenile Idiopathic Arthritis From France Who Completed WA19977 Core Study **Official Regulatory Title:** Long-term, Interventional, Open Label Extension Study Evaluating the Safety of Tocilizumab Treatment in Patients With Polyarticular-course Juvenile Idiopathic Arthritis From France Who Completed the Global, Multinational Trial (WA19977)

12. Clinical Framework (Disease Specific)

Primary Condition: Juvenile Idiopathic Arthritis **Diagnosis Threshold:** Patients who completed the 104-week core trial WA19977 **Medical Methodology:** Interleukin-6 (IL-6) receptor antagonist therapy **Optimization Target:** Maintenance of clinical response and management of adverse events

13. Study Procedures & Methodology

Management Pathway: Long-term open-label biologic therapy extension **Education Protocol (HCP):** Infusion site monitoring, biologic drug administration guidelines, and specialized immunomodulatory safety management. **Education Protocol (Patient):** Education on reporting symptoms of infection, hypersensitivity, and recording JIA activity indicators. **Quality Audit Mechanism:** Routine assessment of drug accountability and strict oversight of laboratory parameter shifts (hepatic/hematologic).

14. Observation & Follow-Up Schedule

Total Duration: 104 Weeks (Treatment) + 12 Weeks (Follow-up)

Onsite Visit Cadence: Every 4 weeks (coinciding with the 8 mg/kg IV infusion) **Remote Monitoring Frequency:** Supplemental check-ins as required for AE follow-up. **Checklist Review Interval:**

Evaluated at each 4-week infusion cycle for disease flares.

15. Sign-off & Approval

Role	Name	Signature	Date		----	----	----	----
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]					Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]					