

1. Administrative & Study Identification

Full Study Title: AN INTERVENTIONAL, PHASE 1, OPEN-LABEL, TWO-COHORT, TWO-PERIOD, FIXED SEQUENCE STUDY TO ESTIMATE THE EFFECT OF MULTIPLE DOSES OF ITRACONAZOLE AND PROBENECID ON THE PHARMACOKINETICS OF SINGLE DOSE PF-07220060 ADMINISTERED UNDER THE FED CONDITION TO HEALTHY PARTICIPANTS **Short Title/Acronym:** A Study to Learn About How the Study Medicines Called Itraconazole and Probenecid Change How the Body Processes PF-07220060 **Protocol Number:** PF-07220060 **NCT Number:** NCT07160738 **Posting ID:** 738761055350026152 **Trial Status:** Not Yet Recruiting (Not yet recruiting participants) **Sponsor Name:** Pfizer **Investigational Product:** PF-07220060 (Atirmociclib), Itraconazole, Probenecid (ATC Code: M04AB01; Trade Names: Benemid, Polycillin-PRB) **Phase of Development:** Phase 1 **Medical Monitor:** Pfizer CT.gov Call Center

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To estimate the effect of multiple doses of a strong CYP3A4 inhibitor (itraconazole) and a potent UGT2B7 inhibitor (probenecid) on the pharmacokinetics (PK) of a single oral dose of PF-07220060 administered under fed conditions. **Secondary Objectives:** To evaluate the safety and tolerability of a single oral dose of PF-07220060 when administered alone and in combination with itraconazole or probenecid.

2.2 Background & Rationale PF-07220060 (Atirmociclib) is a highly selective CDK4 inhibitor currently in development for cancer. Because oncology patients frequently take concomitant medications, it is critical to understand the pharmacokinetic profile of PF-07220060 when co-administered with CYP3A4 and UGT2B7 inhibitors (such as itraconazole and probenecid). This Drug-Drug Interaction (DDI) study is medically necessary to characterize metabolic pathways, avoid dose-limiting toxicities, and inform safe dosing recommendations in clinical practice.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm within fixed-sequence crossover (Period 1 vs. Period 2) **Blinding:** Open-label **Randomization:** Non-Randomized (Fixed sequence assignment)

3.2 Planned Enrollment & Duration Target **\$N\$:** 28 evaluable healthy participants **Number of Sites:** United States (US)
Estimated Study Start: 29-Aug-2025 (Based on First Submitted record) **Estimated Primary Completion:** Q4 2025 / Q1 2026 (approximate timeline for healthy volunteer study)

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Male and female aged 18 to 65 years who are overtly healthy as confirmed by medical evaluation, physical examination, and laboratory tests. 2. Body mass index (BMI) of 17.5-30.5 kg/m²; and a total body weight >50 kg (110 lb). 3. Evidence of a personally signed and dated Informed Consent Document (ICD).

4.2 Exclusion Criteria 1. Evidence or history of blood dyscrasias, uric acid kidney stones, peptic ulcer, or gout (applicable to probenecid cohort). 2. Evidence or history of genetic deficiency of glucose-6-phosphate dehydrogenase in red blood cells (applicable to probenecid cohort). 3. Concomitant use of methotrexate, salicylates, penicillin, sulfonamide, or strong inducers/inhibitors of CYP3A4 or UGT2B7 within 5 half-lives plus 14 days prior to the first dose. History of sensitivity to azole antifungals or probenecid.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: * **Cohort 1:** Period 1 involves a single dose of PF-07220060. Period 2 involves oral itraconazole once daily on Days 1 to 9, with a single dose of PF-07220060 given on Day 4 immediately after itraconazole. * **Cohort 2:** Period 1 involves a single dose of PF-07220060. Period 2 involves oral probenecid four times a day on Days 1 to 9, with a single dose of PF-07220060 given on Day 3 immediately after probenecid. **Route of Administration:** Oral (tablets for PF-07220060 and probenecid; capsules for itraconazole). All PF-07220060 doses are administered ~30 minutes after a moderate fat standard calorie meal. **Duration of Treatment:** Up to 78 days for Cohort 1 and 77 days for Cohort 2.

5.2 Concomitant & Prohibited Therapy Permitted: Standard meals during the clinical confinement. **Prohibited:** Strong CYP3A4 or UGT2B7 inhibitors/inducers, salicylates, penicillin, sulfonamide, ketamine, indomethacin, naproxen, lorazepam, and other interacting therapies during the protocol-defined washout windows.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, Physical Exam, Safety Labs
Baseline / Period 1	Day 1	Confinement, Standard Meal, First Dose PF-07220060, Baseline PK Sampling
Interim / Period 2	Post Washout	PK Sampling up to 120-144h post-dose; Itraconazole or Probenecid loading; Co-administration with PF-07220060
End of Study	Up to Day 78	Final Vital Signs, Exit Medical Evaluation, Discharged from Research Unit

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints 1. Plasma PF-07220060 Maximum Observed Plasma Concentration (\$C_{max}\$) 2. Plasma PF-07220060 Area under the plasma concentration versus time curve (\$AUC\$) **Measurement Method:** Serial blood sampling (pre-dose, and sequentially up to 120-144 hours post-dose) analyzed via validated bioanalytical laboratory methods.

7.2 Secondary & Exploratory Endpoints 1. Number of Participants With Laboratory Abnormalities 2. Number of Participants With Clinically Significant Change From Baseline in Vital Signs 3. Number of Participants With Treatment-Emergent Adverse Events (AEs)

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous observation by clinical staff during in-patient confinement, physical examinations, and serial laboratory monitoring. **Serious Adverse Events (SAEs):** Standard Phase 1 reporting timeline to the sponsor (typically within 24 hours of investigator awareness). **Data Monitoring Committee (DMC):** Internal sponsor medical/safety review team.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: PK Concentration Set (all enrolled participants receiving \$\ge 1\$ dose of study drug with \$\ge 1\$ measurable concentration) and Safety Analysis Set. **Sample Size Justification:** A target of 28 healthy subjects is adequate to estimate the PK parameters and establish safety based on standard intra-subject variance for crossover DDI studies. **Handling of Missing Data:** Pharmacokinetic parameters calculated using non-compartmental analysis (NCA). Missing PK values are excluded from derived parameter calculations per standard conventions.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Standard sponsor clinical site monitoring for Phase 1 research units. **Audit Trail:** Electronic Case Report Form (eCRF) tracking, locked clinical databases, and rigorous sample chain-of-custody for bioanalytical analysis.

11. Study Identifiers & Governance

Primary ID: PF-07220060 **Registry Number:** NCT07160738 **Posting ID:** 738761055350026152 **Sponsor/Lead Organization:** Pfizer **Company:** Pfizer **Study Title:** A Study to Learn About How the Study Medicines Called Itraconazole and Probenecid Change How the Body Processes PF-07220060 **Official Regulatory Title:** AN INTERVENTIONAL, PHASE 1, OPEN-LABEL, TWO-COHORT, TWO-PERIOD, FIXED SEQUENCE STUDY TO ESTIMATE THE EFFECT OF MULTIPLE DOSES OF ITRACONAZOLE AND PROBENECID ON THE PHARMACOKINETICS OF SINGLE DOSE PF-07220060 ADMINISTERED UNDER THE FED CONDITION TO HEALTHY PARTICIPANTS

12. Clinical Framework (Pharmacokinetic Specific)

Primary Condition: Healthy Adults (Well adult) **Diagnosis Threshold:** Overtly healthy, BMI 17.5-30.5 kg/m², weight > 50kg **Medical Methodology:** Pharmacokinetic Drug-Drug Interaction (DDI) Evaluation **Optimization Target:** Measurement of PF-07220060 Plasma \$C_{max}\$ and \$AUC\$

13. Study Procedures & Methodology

Management Pathway: In-patient clinical research unit (CRU) monitoring and PK blood sampling **Education Protocol (HCP):** Site initiation visit and clinical protocol training **Education Protocol (Patient):** Informed consent process and CRU guidelines orientation **Quality Audit Mechanism:** Source data verification, clinical monitoring, and bioanalytical lab validation

14. Observation & Follow-Up Schedule

Total Duration: Up to 78 days **Onsite Visit Cadence:** In-clinic confinement during Period 1 and Period 2 with intensive blood sampling up to 120-144 hours post-dose **Remote Monitoring Frequency:** N/A (Inpatient Phase 1 clinical setting) **Checklist Review Interval:** Continuous/Daily assessment during in-clinic stay

15. Sign-off & Approval

Role	Name	Signature	Date		:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]					Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]					