

1. Administrative & Study Identification

Public Study Title: TACTIVE-U: A Study to Learn About the Study Medicine (Vepdegestrant) When Given With Other Medicines in People With Advanced or Metastatic Breast Cancer (Sub-Study A)

Full/Official Study Title: TACTIVE-U: An Interventional Safety and Efficacy Phase 1b/2, Open-label Umbrella Study to Investigate Tolerability, pk, and Antitumor Activity of Vepdegestrant (ARV-471/PF-07850327), an Oral Proteolysis Targeting Chimera, in Combination With Other Anticancer Treatments in Participants Aged 18 Years and Over With ER+ Advanced or Metastatic Breast Cancer, Sub-study A (ARV-471 in Combination With Abemaciclib) **Short Title/Acronym:** TACTIVE-U Sub-Study A **Protocol Number:** C4891006

NCT Number: NCT05548127 **Trial Status:** Active, not recruiting (ACTIVE_NOT_RECRUITING) **Sponsor/Company Name:** Pfizer

Investigational Product: Vepdegestrant (ARV-471) and Abemaciclib (Trade Name: Verzenio, ATC Code: L01EF03) **Phase of Development:** Phase 1b/2 **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: Phase 1b: To evaluate the Dose-Limiting Toxicities (DLTs) to determine the Recommended Phase 2 Dose (RP2D) of vepdegestrant when administered in combination with abemaciclib. Phase 2: To evaluate the preliminary antitumor activity and safety of the combination RP2D. **Secondary Objectives:** To evaluate additional antitumor activity parameters including Clinical Benefit Rate (CBR), Duration of Response (DOR), Progression-Free Survival (PFS), overall survival, safety profile (AEs, SAEs), lab abnormalities, and pharmacokinetics (PK).

2.2 Background & Rationale To evaluate the safety, tolerability, and efficacy of vepdegestrant (an oral PROTAC estrogen receptor degrader) combined with abemaciclib (a CDK4/6 inhibitor). This study addresses the medical necessity of providing an effective treatment for patients with Estrogen Receptor Positive (ER+) and Human Epidermal Growth Factor Receptor 2 Negative (HER2-) Advanced or Metastatic Breast Cancer (A/MBC) whose disease has progressed following prior lines of therapy, including prior CDK4/6 inhibitors.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm (within Sub-Study A) **Blinding:** Open-label
Randomization: N/A (Escalation/de-escalation approach)

3.2 Planned Enrollment & Duration Target N\$: Approximately 37 patients **Number of Sites:** Global multicenter **Estimated Study Start:** 23-Feb-2023 **Estimated Primary Completion:** 01-Nov-2024

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Histological or cytological diagnosis of ER+ and HER2- advanced/metastatic breast cancer that is not amenable to surgical resection with curative intent ($\geq 1\%$ ER+ stained cells on the most recent tumor biopsy). 2. Prior anticancer therapies: At least 1 and no more than 2 lines of prior therapies for advanced/metastatic disease; 1 line of any CDK4/6 inhibitor-based regimen is required (independent of the setting e.g., adjuvant or advanced/metastatic). 3. At least 1 measurable lesion as defined by RECIST v1.1. 4. ECOG PS ≤ 1 .

4.2 Exclusion Criteria 1. Visceral crisis at risk of life-threatening complications in the short term. 2. Known history of drug-induced pneumonitis or other significant symptomatic deterioration of lung functions. 3. Newly diagnosed brain metastases, or symptomatic CNS metastases, carcinomatous meningitis, or leptomeningeal disease. Participants with a history of CNS metastases or cord compression are eligible if they have been definitively treated, clinically stable and discontinued anti-seizure medications and corticosteroids for at least 14 days prior to enrollment in the study. 4. History of any other tumor malignancies within the past 3 years, except for the following: (1) adequately treated basal or squamous cell carcinoma of the skin; (2) curatively treated in situ carcinoma of the cervix. 5. Inflammatory breast cancer. 6. Impaired cardiovascular function or clinically significant cardiovascular diseases. 7. Concurrent administration of medications, food, or herb supplements that are strong inhibitors and strong/moderate inducers of CYP3A and drugs known to predispose to Torsade de Pointes or QT interval prolongation. 8. Renal impairment, not adequate liver function and/or bone marrow function. 9. Known active infection.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Vepdegestrant administered 1 time a day; Abemaciclib administered 2 times a day. **Route of Administration:** Oral (both medications), taken at home. **Duration of Treatment:** Treatment continues until disease

progression, unacceptable toxicity (side effects become too severe), or withdrawal of consent.

5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care medications per investigator discretion.

Prohibited: Concurrent administration of medications, foods, or herbal supplements that are strong inhibitors and strong/moderate inducers of CYP3A, and drugs known to predispose to Torsade de Pointes or QT interval prolongation.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, Baseline Tumor Imaging, Laboratory Tests
Baseline / Cycle 1	Day 1 to 28	First Dose of Study Medications, PK sampling, DLT observation period for Phase 1b
Interim	Every 4 weeks	Safety Labs, Efficacy Assessment (RECIST v1.1), Dispense Study Drug
End of Study	At Progression	Final Vital Signs, Exit Interview, IP Accountability, Final Safety Review

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints * **Phase 1b:** Number of participants with Dose Limiting Toxicities (DLTs) rate during the DLT observation period (Cycle 1 [28 days]), estimated based on data from DLT-evaluable participants. * **Phase 2:** Percentage of participants with objective response (OR) by investigator assessment. Objective response refers to confirmed complete response (CR) or partial response (PR) according to RECIST version 1.1.

7.2 Secondary & Exploratory Endpoints * **Safety (Phase 1b & 2):** Number of participants experiencing any Adverse Event (AE), Serious Adverse Event (SAE), and Treatment-Related SAE. * **Laboratory Abnormalities (Phase 1b & 2):** Number of participants with laboratory abnormalities in hematology, coagulation parameters, and chemistry parameters. * **Efficacy:** Clinical Benefit Rate (CBR), Duration of Response (DOR), and Progression-Free Survival (PFS).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Untoward medical occurrences evaluated throughout the study, focusing heavily on Treatment-Emergent Adverse Events (TEAEs) and DLTs. **Serious Adverse Events (SAEs):** Reported within standard protocol timelines (typically 24 hours). Special focus on Treatment-Related SAEs. **Data Monitoring Committee (DMC):** Safety profile and dose escalation/de-escalation

decisions are governed by a safety review committee utilizing the mTPI-2 decision criteria.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Phase 1b safety analysis is based on DLT-evaluable participants. Efficacy uses the Intent-to-Treat (ITT) or response-evaluable sets. **Sample Size Justification:** The escalation/de-escalation uses mTPI-2 decision criteria based on the number of DLT-evaluable participants and the number of DLTs observed. **Handling of Missing Data:** Standard handling for oncology studies; censored at the last evaluable time point for time-to-event outcomes (e.g., PFS, DOR).

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Standard routine onsite and remote monitoring to ensure patient safety and data integrity. **Audit Trail:** Data entered into validated electronic Case Report Forms (eCRFs) with comprehensive audit trails.

11. Study Identifiers & Governance

Primary ID (Protocol Number): C4891006 **Registry Number:** NCT05548127 **Posting ID:** 783213531361932075 **Sponsor/Lead Organization:** Pfizer **Study Title:** TACTIVE-U Sub-Study A **Official Regulatory Title:** TACTIVE-U: An Interventional Safety and Efficacy Phase 1b/2, Open-label Umbrella Study to Investigate Tolerability, pk, and Antitumor Activity of Vepdegestrant (ARV-471/PF-07850327), an Oral Proteolysis Targeting Chimera, in Combination With Other Anticancer Treatments in Participants Aged 18 Years and Over With ER+ Advanced or Metastatic Breast Cancer, Sub-study A

12. Clinical Framework (Breast Cancer Specific)

Primary Condition: Breast Cancer (Malignant neoplasm of breast) - Advanced or Metastatic **Diagnosis Threshold:** ER+ ($\geq 1\%$ stained cells) and HER2- **Medical Methodology:** Targeted protein degradation (Vepdegestrant) combined with CDK4/6 inhibition (Abemaciclib) **Optimization Target:** Dose optimization (RP2D) and efficacy evaluation for vepdegestrant + abemaciclib

13. Study Procedures & Methodology

Management Pathway: Daily oral administration of vepdegestrant and twice-daily oral administration of abemaciclib. **Education Protocol (HCP):** Standard protocol training for toxicity management and DLT evaluation. **Education Protocol (Patient):** Education on oral administration schedules, food interactions, and recognizing potential AEs. **Quality Audit Mechanism:** Clinical monitoring, RECIST v1.1 central/local imaging review, and real-time AE tracking.

14. Observation & Follow-Up Schedule

Total Duration: Until disease progression or unacceptable toxicity. **Onsite Visit Cadence:** Clinic visits occur approximately every 4 weeks. **Remote Monitoring Frequency:** As indicated by sponsor data management plans. **Checklist Review Interval:** Routine safety evaluations at every 28-day cycle.

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				