

1. Administrative & Study Identification

Full Study Title: A Phase II, Single-Arm Study of Giredestrant in Patients With Grade 1 Endometrial Cancer **Short Title/Acronym:** EndomERA **Protocol Number:** C044195 **NCT Number:** NCT05634499 **Sponsor Name:** Hoffmann-La Roche **Investigational Product:** Giredestrant (R07197597 / GDC-9545 / RG6171) **Phase of Development:** Phase II **Medical Monitor:** [Name and Contact Information]

2. Study Synopsis & Rationale

2.1 Study Objective **Primary Objective:** To evaluate the efficacy of giredestrant monotherapy in participants with Stage 1, Grade 1 endometrioid endometrial cancer. **Secondary Objectives:** To evaluate the safety, tolerability, and pharmacokinetics of giredestrant monotherapy.

2.2 Background & Rationale Endometrial cancer is a type of cancer that starts in the lining of the uterus. For Stage 1, Grade 1 endometrial cancer, the standard treatment usually includes surgery to remove the uterus (hysterectomy). However, some women may choose hormone medicine instead if they want to preserve their fertility or are not suitable candidates for surgery. There is a significant unmet medical need for better, more effective hormone treatments for these patients. This study evaluates giredestrant as a potential targeted hormonal therapy for this patient population.

3. Study Design & Methodology

3.1 Design Framework **Study Type:** Interventional **Control Type:** Single-arm **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration **Target N\$:** 30 participants **Number of Sites:** 12 global locations **Estimated Study Start:** 27-Jun-2023 **Estimated Primary Completion:** 30-Jun-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Female participants \$\\ge\$ 18\$ years of age. 2. Confirmed Stage 1, Grade 1 endometrial cancer (EC) of endometrioid histology for which participants are willing to receive 6 cycles of study therapy. 3. Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 or 1. 4. Magnetic resonance imaging (MRI)-confirmation of non-deeply invasive tumor

(<50% myometrial invasion) and MRI/CT confirmation of no extrauterine disease. 5. No prior treatment for endometrial cancer.

4.2 Exclusion Criteria 1. Pregnancy or breastfeeding, or intention of becoming pregnant during the study or within 30 days after the final dose. 2. Participants with non-endometrioid histologies, such as serous, clear cell, and mixed. 3. Treatment with investigational therapy, chemotherapy, immunotherapy, CDK4/6 inhibitors, endocrine therapy, biologic therapy, or herbal therapy within 28 days prior to initiation of study enrollment. 4. Any gastrointestinal condition causing malabsorption or obstruction (e.g., celiac sprue, gastric bypass surgery).

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Giredestrant 30 mg taken once a day (QD) on Days 1 to 28 of each 28-day cycle. **Route of Administration:** Oral (PO) **Duration of Treatment:** 6 treatment cycles (approximately 6 months). After completion of 6 cycles, the participant and investigator can choose to continue study treatment for an additional 18 cycles.

5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care medications per investigator discretion. **Prohibited:** Strong CYP3A4 inhibitors or inducers within 14 days or 5 drug elimination half-lives prior to initiation of study treatment.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, Imaging (MRI/CT), Endometrial Biopsy
Baseline	Day 0	Eligibility Confirmation, First Dose
Interim	Day 1 of each Cycle	Safety/Efficacy Assessments, Endometrial biopsies at Month 3 and Month 6
End of Study		
Follow-up	1 month after completing study treatment, and then every 6 months	

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Efficacy of giredestrant monotherapy in treating Grade 1 endometrioid endometrial cancer. **Measurement Method:** Clinical and radiographic evaluations.

7.2 Secondary & Exploratory Endpoints * Safety and tolerability of giredestrant (Adverse Event monitoring). * Pharmacokinetics (PK) of giredestrant in the body.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring at each clinical visit (Start of every treatment cycle). **Serious Adverse Events (SAEs):** Standard global reporting timelines (typically 24 hours). **Data Monitoring Committee (DMC):** Internal/External committee details specified per Sponsor (Hoffmann-La Roche) guidelines.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) for primary efficacy analysis; Safety Set for all participants receiving at least one dose of giredestrant. **Sample Size Justification:** Target enrollment of $N=30$ is determined to be sufficient to estimate the preliminary efficacy and safety profiles for this single-arm Phase II signal-finding study. **Handling of Missing Data:** Last Observation Carried Forward (LOCF) or Multiple Imputation.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Routine site monitoring to ensure protocol compliance and data integrity. **Audit Trail:** Standard electronic Case Report Form (eCRF) tracking and data clarification processes.

11. Study Identifiers & Governance

Primary ID: C044195 **Registry Number:** NCT05634499 **Sponsor/Lead Organization:** Hoffmann-La Roche **Study Title:** EndomERA **Official Regulatory Title:** A Phase II, Single-Arm Study of Giredestrant in Patients With Grade 1 Endometrial Cancer

12. Clinical Framework (Hyperkalemia Specific)

(Note: These fields were originally tailored to the HK-QIP study. Below are the equivalent fields mapped to EndomERA) **Primary Condition:** Stage 1, Grade 1 Endometrioid Endometrial Cancer **Diagnosis Threshold:** Confirmed histology with MRI evidence of $\leq 50\%$ myometrial invasion. **Medical Methodology:** Targeted Endocrine Therapy (investigational SERD). **Optimization Target:** Neoplastic response/tumor shrinkage prior to surgical decision.

13. Study Procedures & Methodology

Management Pathway: Oral administration of 30 mg giredestrant daily for an initial 6 months to evaluate tumor response and delay necessity for hysterectomy. **Education Protocol (HCP):** Protocol training, biopsy sample collection guidelines. **Education Protocol (Patient):** Oral therapy adherence instructions and symptom tracking. **Quality Audit Mechanism:** Central laboratory confirmation of histology.

14. Observation & Follow-Up Schedule

Total Duration: Minimum 6 cycles (approx. 6 months); optional extension up to 24 cycles, followed by long-term monitoring. **Onsite Visit Cadence:** Monthly (Day 1 of each 28-day cycle). **Remote Monitoring Frequency:** Follow-ups conducted at 1 month post-treatment, then every 6 months. **Checklist Review Interval:** Routine adverse event and adherence review every cycle.

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				