

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 02-Apr-2026

1. Administrative & Study Identification

Full Study Title: Korean Post-marketing Surveillance for Xeljanz XR (Registered) **Short Title/Acronym:** Korean PMS for Xeljanz XR
Protocol Number: A3921369 **NCT Number:** NCT04876781 **Posting ID:** 85643817693988424 **Sponsor/Company Name:** Pfizer **Preferred UMLS Name:** Postmarketing Drug Surveillance **Phase of Development:** Not Applicable (Phase ID: Not Applicable) **Trial Status:** RECRUITING (Status Name: RECRUITING | Status Description: Currently recruiting participants) **Source Level:** 5 **Medical Monitor:** Pfizer Pharmacovigilance / Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the safety of Xeljanz XR under routine clinical practice based on adverse events (AEs) occurring during the 6 months from the first dose. **Secondary Objectives:** To evaluate the effectiveness of Xeljanz XR using the modified Disease Activity Score (DAS28) change from baseline, EULAR response, ACR20 response, and ASAS 20 response.

2.2 Background & Rationale To fulfill the risk management plan in accordance with Article 4-1-11 of the "Regulation on Safety of Medicinal Products, etc." in Korea. As part of an additional pharmacovigilance activity, this Post-marketing Surveillance (PMS) was planned to evaluate the safety and effectiveness of Xeljanz XR under routine clinical practice.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational (Open-label, non-comparative, non-interventional, prospective, multi-center) **Control Type:** Single-arm **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Targeted Enrollment: At least 200 patients **Number of Sites:** Multi-center (South Korea) **Estimated Study Start:** 12-Jan-2022 **Estimated Primary Completion:** ~12-Jan-2026 (Approximately four years from study start)

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Treatment of moderate to severe active Rheumatoid Arthritis (RA) in adult patients (18 years of age or older) who have had an inadequate response or are intolerant to methotrexate. 2. In combination with methotrexate, treatment of

active psoriatic arthritis (PsA) in adult patients (18 years of age or older) who have had an inadequate response or who have been intolerant to a prior DMARD therapy. 3. Treatment of active ankylosing spondylitis (AS) in adult patients (18 years of age or older) who have responded inadequately to conventional therapy. 4. **Criteria Type:** PHASE_REQUIREMENT - Criteria Text: Trial must be in Not Applicable.

4.2 Exclusion Criteria 1. Patients who do not meet the approved local prescribing and labeling information for Xeljanz XR in Korea. 2. Patients with a history of hypersensitivity to any ingredients of this product. 3. Patients with conditions contraindicated for Tofacitinib, such as active serious infections (e.g., tuberculosis, sepsis). 4. Patients with severe hepatic function disorder, or abnormal lab values (ANC <1,000 cells/mm3, ALC <500 cells/mm3, Hemoglobin <9 g/dL). 5. Pregnant or possibly pregnant women.

5. Intervention & Treatment Plan

5.1 Investigational Regimen **Drug Name:** tofacitinib (Drug Preferred Name: tofacitinib) **Trade Name:** Xeljanz (Xeljanz XR) **ATC Code:** L04AA29 **Dosage/Frequency:** Xeljanz XR 11 mg (Tofacitinib citrate) once daily as per routine clinical practice. **Route of Administration:** Oral (extended-release tablets). **Duration of Treatment:** 6 months of observation from the first dose. **EMA URL:** https://www.ema.europa.eu/en/documents/product-information/xeljanz-epar-product-information_en.pdf

5.2 Concomitant & Prohibited Therapy Permitted: Methotrexate (required for PsA in this context, permitted for RA) and other routine conventional therapies. **Prohibited:** Concurrent use with biologic DMARDs or potent immunosuppressants (as per standard prescribing guidelines for Xeljanz XR).

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening/Baseline	Day 0	Informed Consent, Medical History, First Dose of Xeljanz XR, Baseline Disease Activity (DAS28)
End of Study	Month 6	Final Safety Evaluation, Adverse Event Assessment, Efficacy Assessment (DAS28, EULAR, ACR20, ASAS20)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Incidence and severity of treatment-emergent adverse events (AEs) occurring during the 6 months from the first dose of Xeljanz XR. **Measurement Method:** Evaluated via investigator assessment of AEs and their

relatedness to Xeljanz XR during routine clinical practice (Yes/No).

7.2 Secondary & Exploratory Endpoints * Change from baseline in the modified Disease Activity Score using 28 joint counts (DAS28). * Percentage of participants with European League Against Rheumatism (EULAR) response based on DAS28. * Number of participants with an American College of Rheumatology 20% improvement criteria (ACR20) response after treatment. * ASAS 20 Response Criteria (improvement of at least 20% and absolute improvement of at least 10 units on a 0-100 scale in at least three domains).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Assessed during the 6 months from the first dose. Relatedness to Xeljanz XR is assessed directly by the investigator (Yes/No). **Serious Adverse Events (SAEs):** Reported per local Korean pharmacovigilance regulations (typically within 24 hours of site awareness). **Data Monitoring Committee (DMC):** Managed via the Sponsor's Pharmacovigilance and Safety Risk Management teams.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all enrolled patients who received at least one dose of Xeljanz XR and had safety follow-up). **Sample Size Justification:** Target of at least 200 patients to ensure sufficient pharmacovigilance data as mandated by local regulatory standards. **Handling of Missing Data:** Last Observation Carried Forward (LOCF) for efficacy endpoints or observed case analysis as per standard observational protocols.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP), the Declaration of Helsinki, and Article 4-1-11 of the "Regulation on Safety of Medicinal Products, etc." in Korea. **Monitoring Plan:** Routine pharmacovigilance data collection, risk management plan tracking, and source data verification as applicable for PMS. **Audit Trail:** eCRF locking and standard safety database tracking.

11. Study Identifiers & Governance

Primary ID: A3921369 **Registry Number:** NCT04876781 **Sponsor/Lead Organization:** Pfizer **Study Title:** Korean Post-marketing

12. Clinical Framework

Disease Areas: Active Moderate to Severe Rheumatoid Arthritis, Active Ankylosing Spondylitis, Active Psoriatic Arthritis
Condition Name: Arthritis (Preferred Name: Arthritis) **Condition Definition:** Acute or chronic inflammation of JOINTS. **Semantic Type:** Disease or Syndrome **Medical Methodology:** Routine clinical practice / Pharmacovigilance observation **Optimization Target:** Disease Activity Score (DAS28) improvement, ACR20, and ASAS20 response thresholds

Condition Ontology Codes: * **MeSH Code:** D001168 * **HPO Code:** HP: 0001369 * **SNOMED ID:** 3723001 * **SNOMED Hierarchy:** 399269003 | Arthropathy | Arthropathy (disorder), 785875003 | Disorder of joint region | Disorder of joint region (disorder), 64572001 | Disease | Disease (disorder), 88230002 | Disorder of skeletal system | Disorder of skeletal system (disorder), 928000 | Disorder of musculoskeletal system | Disorder of musculoskeletal system (disorder), 362965005 | Disorder of body system | Disorder of body system (disorder), 363169009 | Inflammation of specific body organs | Inflammation of specific body organs (disorder), 363170005 | Inflammation of specific body structures or tissue | Inflammation of specific body structures or tissue (disorder), 128139000 | Inflammatory disorder | Inflammatory disorder (disorder), 363179006 | Inflammatory disorder of musculoskeletal system | Inflammatory disorder of musculoskeletal system (disorder), 363171009 | Inflammation of specific body systems | Inflammation of specific body systems (disorder), 897444001 | Structural abnormality of skeleton | Structural abnormality of skeleton (disorder).

13. Study Procedures & Methodology

Management Pathway: Routine clinical treatment pathway for RA, PsA, and AS. **Education Protocol (HCP):** Training on the Risk Management Plan and safety reporting obligations for Xeljanz XR in the Korean market. **Education Protocol (Patient):** Provision of patient information on drug safety and AE reporting. **Quality Audit Mechanism:** Pharmacovigilance tracking, AE incidence reporting compliance, and site audits.

14. Observation & Follow-Up Schedule

Total Duration: 6 months per patient **Onsite Visit Cadence:** Baseline and Month 6 (or aligned with routine care standard practice). **Remote Monitoring Frequency:** Continuous AE reporting via standard pharmacovigilance channels. **Checklist Review**

Interval: Safety and effectiveness evaluation at the 6-month study endpoint.

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Insert Name]	____	02-Apr-2026				
Clinical Operations Lead	[Insert Name]	____	02-Apr-2026				
Lead Statistician	[Insert Name]	____	02-Apr-2026				