

1. Administrative & Study Identification

Full Study Title: Observational Study in Japanese Patients With Peripheral T-Cell Lymphoma Who Received Second-Line Therapy
Short Title/Acronym: R/R PTCL Observational Study Japan (CA073-1019)
Protocol Number: CA073-1019
NCT Number: NCT06422247
Sponsor Name: Bristol-Myers Squibb
Investigational Product: Systemic Therapy (Various 2nd-line single agents such as brentuximab vedotin, romidepsin, tucidinostat, etc.)
Phase of Development: Observational
Medical Monitor: BMS Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To elucidate the outcomes of 2nd-line therapy and overall survival in Japanese patients with relapsed or refractory (R/R) mature T/NK-cell lymphomas (MTNKL) in the era of novel agents.
Secondary Objectives: To evaluate time to next treatment (TTNT) and analyze real-world treatment patterns.

2.2 Background & Rationale Mature T-cell and natural killer-cell lymphomas (MTNKLs) are often refractory to conventional treatment and associated with a poor prognosis. Between March 2014 and March 2024, nine novel agents were approved in Japan as single agents for R/R MTNKL. However, the real-world treatment patterns and prognosis of patients in this new treatment era remain unclear. This nationwide retrospective study generates real-world evidence regarding clinical outcomes for patients receiving these novel agents as second-line therapies.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational (Nationwide, retrospective cohort study)
Control Type: N/A (Single-arm)
Blinding: Open-label / N/A (Retrospective)
Randomization: N/A

3.2 Planned Enrollment & Duration Overall Status: Completed
Target \$N\$: 256 patients analyzed
Number of Sites: 23 institutions in Japan
Study Start Date: 05-Apr-2024

Primary Completion Date: 05-Feb-2025 (Actual)

Study Completion Date: 05-Feb-2025 (Actual)

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

1. Male or female participants, aged ≥ 18 years of age at diagnosis of PTCL.
2. Confirmed diagnosis of specific subtypes of peripheral T-cell lymphoma (PTCL) or mature T-cell and NK-cell lymphomas (MTNKL) according to The World Health Organization (WHO) classification of lymphoid neoplasm, 4th edition defined by WHO and International Agency for Research on Cancer (IARC).
3. Participants who have been treated with a systemic therapy for PTCL and have initiated a systemic therapy as a second-line therapy for relapsed or refractory PTCL between April 1, 2018 and March 31, 2023.

4.2 Exclusion Criteria

1. Participants who have a medical history of PTCL treatment by an unapproved drug in Japan as of 31 March 2024 or an off-label drug for PTCL in Japan as of 31 March 2024.
 2. Participants who have a medical history of participation in a separately defined registration study for regulatory approval in PTCL.
 3. Participants who have a medical history of PTCL treatment in a separately defined clinical study with intervention by an on-label regimen for PTCL.
 4. Participants judged to be inappropriate for enrollment in this study by the site investigator.
-

5. Intervention & Treatment Plan

5.1 Investigational Regimen

Dosage/Frequency: Real-world administration of systemic 2nd-line therapies (e.g., brentuximab vedotin, tucidinostat, romidepsin, pralatrexate) dosed according to local prescribing information.

Route of Administration: Intravenous or Oral, depending on the specific therapy prescribed.

Duration of Treatment: Varied based on real-world clinical practice, continued until disease progression, unacceptable toxicity, or initiation of the next line of treatment.

5.2 Concomitant & Prohibited Therapy

Permitted: Concomitant supportive care standard in real-world Japanese clinical practice.

Prohibited: Unapproved or off-label experimental therapies for PTCL.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening	N/A Retrospective chart review of the eligible index period (April 2018 - March 2023) Baseline
Day 0	Initiation of 2nd-line systemic therapy	Interim	
N/A	Retrospective EMR data extraction of treatment duration and survival milestones	End of Study	N/A End of patient follow-up or date of death

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint

Definition: Overall Survival (OS) after 2nd-line therapy initiation (OS-2L), defined as the time from the start of 2nd-line therapy to death from any cause.

Measurement Method: Extraction and calculation from Electronic Medical Records (EMR).

7.2 Secondary & Exploratory Endpoints

1. Time to next treatment (TTNT).
 2. Treatment patterns (e.g., types of single agents used, frequency of hematopoietic stem cell transplantation [HSCT]).
-

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: N/A (Retrospective survival study; proactive AE collection is not the primary focus).

Serious Adverse Events (SAEs): N/A.

Data Monitoring Committee (DMC): N/A.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Retrospective observational cohort (Eligible patients treated with 2nd-line systemic therapy, \$N=256\$).

Sample Size Justification: Convenience sample of all eligible patients meeting the inclusion/exclusion criteria at the 23 participating institutions during the specified timeframe.

Handling of Missing Data: Standard handling for observational EMR data; Kaplan-Meier methodology used to estimate median OS and TTNT.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Pharmacoepidemiology Practices (GPP) and Japanese ethical guidelines for medical and health research involving human

subjects.

Monitoring Plan: Centralized data quality review of extracted EMR data.

Audit Trail: Secured clinical database with standard audit trails for retrospective electronic data capture.

11. Study Identifiers & Governance

Primary ID: CA073-1019

Registry Number: NCT06422247

Sponsor/Lead Organization: Bristol-Myers Squibb

Study Title: Observational Study in Japanese Patients With Peripheral T-Cell Lymphoma Who Received Second-Line Therapy

Official Regulatory Title: Observational Study in Japanese Patients With Peripheral T-Cell Lymphoma Who Received Second-Line Therapy

12. Clinical Framework (PTCL Specific)

Primary Condition: Peripheral T-cell Lymphoma (PTCL) / Relapsed or Refractory Mature T/NK-cell Lymphoma

Diagnosis Threshold: Confirmed histological diagnosis per WHO 4th edition classification defined by WHO and IARC

Medical Methodology: Real-world 2nd-line systemic therapy (Novel single agents)

Optimization Target: Extension of Overall Survival (OS-2L)

13. Study Procedures & Methodology

Management Pathway: Routine real-world clinical care for R/R PTCL in Japan

Education Protocol (HCP): N/A (Retrospective observational study)

Education Protocol (Patient): N/A (Retrospective observational study)

Quality Audit Mechanism: Retrospective data verification and validation of EMR abstraction

14. Observation & Follow-Up Schedule

Total Duration: Variable retrospective follow-up (analyzing patients treated between April 2018 and March 2023)

Onsite Visit Cadence: N/A (Retrospective medical chart review)

Remote Monitoring Frequency: N/A

Checklist Review Interval: N/A

15. Sign-off & Approval

	Role		Name		Signature		Date			:---		:---		:---		:---	
	Principal Investigator		[Name]		_____		02-Apr-2026										
	Clinical Operations Lead		[Name]		_____		02-Apr-2026										
	Statistician		[Name]		_____		02-Apr-2026										