

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 02-Apr-2026

1. Administrative & Study Identification

Full Study Title: Treatment Patterns and Outcomes in Patients With Lower-risk Myelodysplastic Syndromes Treated With Luspatercept in China **Short Title/Acronym:** REACH Study **Protocol Number:** CA056-1166 **NCT Number:** NCT07073690 **Posting ID:** 866882088164425056 **Sponsor Name:** Bristol Myers Squibb **Company:** Bristol Myers Squibb **Investigational Product:** Luspatercept (Trade Name: Reblozyl) **Phase of Development:** Not Applicable (Observational, Real-world retrospective study) **Trial Status:** ACTIVE_NOT_RECRUITING **Medical Monitor:** Bristol Myers Squibb Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate real-world treatment patterns and clinical outcomes in adults with lower-risk Myelodysplastic Syndromes (LR-MDS) who were erythropoiesis-stimulating agent (ESA)-naïve and received luspatercept treatment in China. **Secondary Objectives:** To assess the duration of luspatercept monotherapy treatment, Health Care Resource Utilization (HCRU), and participant transfusion independence and increase in hemoglobin.

2.2 Background & Rationale This study focuses on understanding how patients with lower-risk Myelodysplastic Syndromes (LR-MDS) in China are treated with luspatercept. Myelodysplastic Syndrome is defined as a clonal hematopoietic disorder characterized by dysplasia and ineffective hematopoiesis in one or more of the hematopoietic cell lines. The dysplasia may be accompanied by an increase in myeloblasts, but the number is less than 20%, which, according to the WHO guidelines, is the requisite threshold for the diagnosis of acute myeloid leukemia. It may occur de novo or as a result of exposure to alkylating agents and/or radiotherapy (WHO, 2001). These patients have not previously received erythropoiesis-stimulating agents (ESAs). This research is necessary to gather real-world data on the effectiveness of luspatercept in managing LR-MDS symptoms, offering a new approach to assessing the real-world impact of the care provided.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational, Retrospective Cohort Study **Control Type:** Single-arm (Non-Probability Sample) **Blinding:** Open-label / Unblinded **Randomization:** N/A

3.2 Planned Enrollment & Duration Target **\$N\$:** 80 evaluable patients **Number of Sites:** Multiple sites in China **Estimated Study Start:** 18-Apr-2025 **Estimated Primary Completion:** 26-Nov-2025

4. Subject Selection (Eligibility Criteria)

- 4.1 Inclusion Criteria** 1. Participants \$\\ge\$ 18\$ years of age. 2. Documented diagnosis of no del(5q) MDS that meets lower-risk Myelodysplastic Syndromes (MDS) classification. (Diagnosis criteria and risk classification criteria are based on clinical practice). 3. Patients who were erythropoiesis-stimulating agent (ESA)-naïve have received luspatercept monotherapy at least 2 cycles regardless of RS status. 4. Historical documentation of anemia before luspatercept treatment.
- 4.2 Exclusion Criteria** 1. Had a history of acute myeloid leukemia (AML) prior to MDS diagnosis. 2. Received stem cell transplant prior to luspatercept treatment initiation.
-

5. Intervention & Treatment Plan

- 5.1 Investigational Regimen** **Product Name:** Luspatercept (Trade Name: Reblozyl) **ATC Code:** B03XA06 **Dosage/Frequency:** Luspatercept monotherapy (standard of care dosing, minimum of 2 cycles). **Route of Administration:** Subcutaneous injection. **Duration of Treatment:** Retrospectively assessed across the duration of monotherapy. **EMA Reference:** [Reblozyl EPAR Product Information](#)
- 5.2 Concomitant & Prohibited Therapy Permitted:** Routine care and medications per standard clinical practice for MDS. **Prohibited:** Prior use of Erythropoiesis-stimulating agents (ESA) or prior stem cell transplant.
-

6. Study Timeline & Visit Schedule

(Note: As this is a retrospective observational study, the standard visit schedule is substituted with data abstraction milestones)

Visit ID	Window (Days)	Key Activities/Procedures
1	Screening/Index	N/A Identification of eligible patients, historical documentation of anemia
2	Baseline	N/A Extraction of patient characteristics, diagnosis, and pretreatment lab values
3	Observation Period	N/A Data extraction of luspatercept dosing, hemoglobin levels, and transfusion records
4	End of Abstraction	N/A Final calculation of Health Care Resource Utilization (HCRU)

7. Outcome Measures (Endpoints)

7.1 Primary Outcomes 1. Participant red blood cell transfusion independence. 2. Participants mean hemoglobin (Hb) increase ≥ 1.5 g/dL.

7.2 Secondary Outcomes 1. Participant transfusion independence and increase in hemoglobin (Hb). 2. Duration of luspatercept monotherapy treatment. 3. Health Care Resource Utilization (HCRU).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Retrospective extraction of documented AEs from medical records. **Serious Adverse Events (SAEs):** Captured as documented in real-world clinical practice. **Data Monitoring Committee (DMC):** N/A (Retrospective observational study).

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: All eligible patients who received at least 2 cycles of luspatercept. **Sample Size Justification:** $N = 80$ is targeted to provide a descriptive real-world evidence cohort for Chinese patients with LR-MDS. **Handling of Missing Data:** Standard handling/exclusion methods applicable to retrospective real-world data analysis.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Pharmacovigilance Practices (GPP) and applicable real-world data collection guidelines. **Monitoring Plan:** Source data verification from medical records to ensure integrity and accuracy of the extracted data. **Audit Trail:** Secure database tracking for eCRFs during the retrospective extraction process.

11. Study Identifiers & Governance

Primary ID: {{CA056-1166}} **Posting ID:** {{866882088164425056}}
Registry Number: {{NCT07073690}} **Sponsor/Lead Organization:** {{Bristol Myers Squibb}} **Study Title:** {{Treatment Patterns and Outcomes in Patients With Lower-risk Myelodysplastic Syndromes Treated With Luspatercept in China}} **Official Regulatory Title:** {{Real-World Treatment Patterns and Outcomes of Luspatercept in Patients With Lower-risk Myelodysplastic Syndromes (LR-MDS) in China ('REACH' Study)}}

12. Clinical Framework (Myelodysplastic Syndromes Specific)

Primary Condition: {{Lower-risk Myelodysplastic Syndromes (LR-MDS) / Myelodysplastic Syndrome}} **Semantic Type:** {{Neoplastic Process}} **Clinical Codes:** {{MeSH: D009190 | HPO: HP:0002863 | SNOMED: 109995007}} **Diagnosis Threshold:** {{no del(5q) MDS, lower-risk classification with documented anemia}} **Medical Methodology:** {{Luspatercept Monotherapy in ESA-naïve patients}} **Optimization Target:** {{Red blood cell transfusion independence and Hb increase ≥ 1.5 g/dL}}

13. Study Procedures & Methodology

Management Pathway: {{Standard real-world clinical care for LR-MDS}} **Education Protocol (HCP):** {{N/A - Retrospective Study}} **Education Protocol (Patient):** {{N/A - Retrospective Study}} **Quality Audit Mechanism:** {{Electronic medical record data verification}}

14. Observation & Follow-Up Schedule

Total Duration: {{Retrospective observation window based on historical records}} **Onsite Visit Cadence:** {{N/A - Retrospective Data Extraction}} **Remote Monitoring Frequency:** {{N/A}} **Checklist Review Interval:** {{N/A}}

15. Sign-off & Approval

Role	Name	Signature	Date		----	----	----	----
Principal Investigator	[Insert Name]	_____	02-Apr-2026					
Clinical Operations Lead	[Insert Name]	_____	02-Apr-2026					
Lead Statistician	[Insert Name]	_____	02-Apr-2026					