

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 03-Apr-2026

1. Administrative & Study Identification

Full Study Title: A Study of Felmetatug Vedotin/SGN-B7H4V in Advanced Solid Tumors **Official Regulatory Title:** A Phase 1 Study of Felmetatug Vedotin/SGN-B7H4V in Advanced Solid Tumors **Short Title/Acronym:** SGNB7H4V-001 **Protocol Number:** SGNB7H4V-001 (also C5761001) **NCT Number:** NCT05194072 **Sponsor/Company Name:** Pfizer (formerly initiated by Seagen) **Investigational Product:** Felmetatug Vedotin (SGN-B7H4V) **Phase of Development:** Phase 1 **Medical Monitor:** Clinical Operations Lead, Pfizer

2. Study Synopsis & Rationale

2.1 Study Objective **Primary Objective:** To evaluate the safety and tolerability of the medicine Felmetatug Vedotin given alone and in combination with pembrolizumab in participants with solid tumors. This includes identifying the number of participants with dose-limiting toxicities (DLTs), adverse events (AEs), and laboratory abnormalities. **Secondary Objectives:** To evaluate the preliminary anti-tumor activity, including Confirmed Objective Response Rate (ORR) by investigator assessment, Complete Response Rate (CRR), and Duration of Response (DOR).

2.2 Background & Rationale To address the critical unmet medical need for patients with relapsed or refractory advanced solid tumors who have exhausted or are intolerant to standard-of-care (SOC) therapies. B7-H4 is an immune checkpoint ligand that is overexpressed in various solid tumors but has low expression in normal tissues. Felmetatug vedotin is a targeted antibody-drug conjugate (ADC) designed to bind to B7-H4 and deliver a potent monomethyl auristatin E (MMAE) payload directly into tumor cells, leading to targeted cell death.

3. Study Design & Methodology

3.1 Design Framework **Study Type:** Interventional **Control Type:** Single-arm (Dose escalation/expansion) **Blinding:** Open-label **Randomization:** N/A (Parallel group assignments based on tumor type and combination therapies)

3.2 Planned Enrollment & Duration **Targeted Enrollment:** 250 participants **Number of Sites:** Multiple global trial locations **Estimated Study Start:** 11-Jan-2022 **Trial Status:** TERMINATED. *Note: The study was terminated early by Pfizer in early 2025 for strategic reasons, as early data indicated the drug was unlikely*

to achieve a meaningful clinical improvement over standard-of-care therapies.

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Adult participants (≥ 18 years) with histologically or cytologically confirmed locally advanced unresectable or metastatic solid tumor types, including: * High-grade serous epithelial ovarian cancer, primary peritoneal cancer, or fallopian tube cancer * HER2-negative, HR positive breast cancer * Triple-negative breast cancer (TNBC) * Endometrial carcinoma * Non-small cell lung cancer (Squamous cell carcinoma [SqCC], Adenocarcinoma [AC]) * Cholangiocarcinoma or gallbladder carcinoma * Adenoid cystic carcinoma (ACC) 2. ECOG Performance Status of 0 or 1. 3. Participants must have measurable disease per RECIST version 1.1 at baseline (not applicable for E3 participants). 4. **Treatment/Cohort-Specific Criteria:** * **Parts A, B, and C:** Disease must be relapsed or refractory or intolerant to SOC therapies, and should have no appropriate SOC therapeutic option. * **Part D and E1/E2:** Participants must have had no prior treatment for locally advanced unresectable or metastatic TNBC. * **Cohort E1:** TNBC with CPS ≥ 10 by local testing. * **Cohort E2:** TNBC with CPS < 10 by local testing. * **Cohort E3:** Triple negative breast cancer with residual disease following neoadjuvant therapy and definitive surgery; must have completed at least 6 cycles of neoadjuvant therapy.

4.2 Exclusion Criteria 1. History of another malignancy within 3 years before the first dose of study drug. Any evidence of residual disease from a previously diagnosed malignancy. 2. Known active central nervous system (CNS) metastases. Participants with previously treated brain metastases may participate provided they are clinically stable for at least 4 weeks, have no new/enlarging brain metastases, and are off corticosteroids prescribed for symptoms associated with brain metastases for at least 7 days prior to treatment. 3. Carcinomatous meningitis. 4. Previous receipt of an MMAE-containing agent or an agent targeting B7-H4. 5. Pre-existing neuropathy $\geq$ Grade 2 per NCI CTCAE Version 5.0. 6. Corneal disease or injury requiring treatment or active monitoring.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Investigational Drug: Felmetatug Vedotin evaluated across escalating dose levels alone or in combination with **pembrolizumab** (Trade Name: **Keytruda**; ATC Code: **L01FF02**). **Study Structure:** * **Parts A & B:** Find out how much Felmetatug Vedotin should be given. * **Part C:** Use the amount found in A & B to find out safety and efficacy. * **Part D:** Find out if and how much Felmetatug Vedotin can be given with

pembrolizumab. * **Part E:** Use the amount found in Part D to find out safety and efficacy of the combination in TNBC cohorts (E1, E2, E3). **Route of Administration:** Intravenous (IV).

5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care for adverse event management. **Prohibited:** Corticosteroids prescribed for brain metastases symptoms within 7 days of treatment; concurrent experimental anti-cancer therapies.

6. Study Timeline & Visit Schedule

| Visit ID | Window (Days) | Key Activities/Procedures | | :---
| :--- | :--- | | **Screening** | 2 to 4 weeks prior | Informed Consent, Medical History, Tumor Tissue collection, Baseline Imaging | | **Treatment (Parts A-E)** | Every 21-day cycle | IV Infusion, PK/PD Sampling, Safety Labs, AE Monitoring | | **Interim Scans** | Every 8 to 12 weeks | Efficacy Assessment (RECIST 1.1 imaging) | | **End of Study** | Up to 5 years | Post-treatment safety follow-up, Survival tracking |

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints 1. Number of participants with adverse events (AEs). 2. Number of participants with laboratory abnormalities. 3. Number of participants with dose limiting toxicities (DLTs).

7.2 Secondary Endpoints 1. Confirmed objective response rate (ORR) by investigator assessment. 2. Complete response rate (CRR). 3. Duration of response (DOR).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard patient monitoring prior to each cycle dose for any untoward medical occurrence in a clinical investigational participant administered a medicinal product. **Serious Adverse Events (SAEs):** Required continuous monitoring and expedited reporting within 24 hours to the sponsor. **Data Monitoring Committee (DMC):** Internal safety review committee governed dose escalation decisions and toxicity tracking.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Analysis Set (all participants who received at least 1 dose of study drug); Efficacy Evaluable Set (participants with measurable baseline disease and ≥ 1 post-baseline scan). **Sample Size Justification:** Targeted 250

participants, sized according to standard Phase 1 dose-escalation algorithms to identify the Maximum Tolerated Dose (MTD) and evaluate preliminary expansion cohorts. **Handling of Missing Data:** Summarized descriptively; Time-to-event outcomes (DOR) evaluated via Kaplan-Meier methodologies.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study was conducted in accordance with Good Clinical Practice (GCP) guidelines and the Declaration of Helsinki. **Monitoring Plan:** Scheduled onsite and remote clinical site monitoring for source data verification. **Audit Trail:** Secure Electronic Data Capture (EDC) systems with eCRF locking and stringent data clarification mechanisms.

11. Study Identifiers & Governance

Primary ID: SGNB7H4V-001 **NCT Number:** NCT05194072 **Posting ID:** 89661651962581739 **Sponsor/Lead Organization:** Pfizer **Official Regulatory Title:** A Phase 1 Study of Felmetatug Vedotin/SGN-B7H4V in Advanced Solid Tumors

12. Clinical Framework (Advanced Solid Tumors Specific)

Preferred UMLS Name: Advanced Solid Tumor **Disease Areas/Conditions:** Ovarian Neoplasms, Peritoneal Neoplasms, Fallopian Tube Neoplasms, Triple Negative Breast Neoplasms, HER2 Negative Breast Neoplasms, Hormone Receptor Positive Breast Neoplasms, Endometrial Neoplasms, Carcinoma, Non-Small-Cell Lung, Cholangiocarcinoma, Gallbladder Carcinoma, Adenoid Cystic Carcinoma. **Diagnosis Threshold:** Histologically or cytologically confirmed locally advanced unresectable or metastatic disease. **Medical Methodology:** Antibody-Drug Conjugate (ADC) and Immune Checkpoint Inhibitor Combination. **Optimization Target:** Targeted B7-H4 cytotoxicity / T-cell immune checkpoint blockade.

13. Study Procedures & Methodology

Management Pathway: Phase 1 Dose-Escalation and Expansion Oncology Pathway **Education Protocol (HCP):** Site Initiation Visits (SIV) and comprehensive protocol/AE management training **Education Protocol (Patient):** Informed consent process, treatment expectation outlining, and symptom reporting guidelines **Quality Audit Mechanism:** Central and Investigator-led review of RECIST 1.1 tumor assessments

14. Observation & Follow-Up Schedule

Total Duration: Up to 5 years (Follow-up) **Onsite Visit Cadence:** Pre-infusion clearance and drug administration at every cycle
Remote Monitoring Frequency: Routine safety check-ins and post-treatment survival tracking **Checklist Review Interval:** Routine imaging (CT/MRI) assessment every 8 to 12 weeks

15. Sign-off & Approval

Role	Name	Signature	Date	_____	_____	_____	_____
Principal Investigator	[Insert Name]	_____	[DD-MMM-YYYY]	_____	_____	_____	_____
Clinical Operations Lead	[Insert Name]	_____	[DD-MMM-YYYY]	_____	_____	_____	_____
Lead Statistician	[Insert Name]	_____	[DD-MMM-YYYY]	_____	_____	_____	_____