

1. Administrative & Study Identification

Full Study Title: A Randomized, Double-Blind, Placebo- and Open-Label Active Comparator-Controlled, Parallel-Group, Multi-Center Phase II Study to Evaluate the Efficacy, Safety, and Tolerability of Once-Daily R07795081 Administered for 30 Weeks to Participants With Type 2 Diabetes Mellitus **Short Title/Acronym:** A Study to Compare Different Doses of R07795081 With a Placebo or Semaglutide in People With Type 2 Diabetes **Protocol Number:** BP45703 / 2024-520322-11-00 **NCT Number:** NCT07112872 **Posting ID:** 961137043236129959 **Sponsor/Company Name:** Hoffmann-La Roche (Roche) **Trial Status:** Recruiting **Investigational Product:** R07795081 (also known as CT-996 or RG6652) **Active Comparator Drug:** Semaglutide (ATC Code: A10BJ06; Trade Names: Wegovy, Rybelsus, Ozempic; [EMA Product Info \(Ozempic\)](#), [EMA Product Info \(Rybelsus\)](#), [EMA Product Info \(Wegovy\)](#)) **Preferred UMLS Name:** Antidiabetic Agents **Phase of Development:** Phase II **Medical Monitor:** Hoffmann-La Roche Clinical Development Medical Director

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the efficacy, tolerability, and safety of once-daily oral R07795081 for glycemic control in adult participants with Type 2 diabetes mellitus (T2D). **Secondary Objectives:** To compare the efficacy of R07795081 against the active comparator (semaglutide) and assess specific secondary glycemic metrics, including reaching target HbA1c levels and fasting plasma glucose improvements.

2.2 Background & Rationale To address the need for more accessible and tolerable treatments for Type 2 Diabetes and obesity. Small-molecule oral GLP-1 receptor agonists like R07795081 present an advantageous pharmacological approach, offering the convenience of oral administration, potentially improved patient adherence, and cost-effectiveness compared to traditional injectable peptide GLP-1 agonists. This Phase II dose-finding study seeks to establish the optimal dose of R07795081 for glycemic control in humans against placebo and semaglutide.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Placebo-controlled and Open-Label Active Comparator-controlled **Blinding:** Double-blind (for R07795081 vs. Placebo) and Open-label

(for Active Comparator Semaglutide) **Randomization:** Parallel-group, randomized, multi-center, dose-range-finding

3.2 Planned Enrollment & Duration Target Enrollment (\$N\$): 240 participants **Number of Sites:** Multi-center **Current Status:** Recruiting **Estimated Primary Completion:** After conclusion of the 30-week treatment duration

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Have a diagnosis of Type 2 diabetes mellitus (T2D) for at least 6 months before screening. 2. Have an HbA1c $\geq 7\%$ and $\leq 10.5\%$ at screening. 3. Management of T2D with diet and exercise alone or with either a stable dose of metformin or/and sodium-glucose cotransporter-2 (SGLT-2) inhibitors. 4. Body mass index (BMI) ≥ 23.0 kg/m² at screening. 5. A stable body weight within 3 months prior to screening (maximum 5% self-reported body weight gain and/or loss).

4.2 Exclusion Criteria 1. Have Type 1 diabetes (T1D), history of ketosis or hyperosmolar state/coma, or any other types of diabetes except T2D. 2. Have had 1 or more episodes of Level 3 hypoglycemia or has hypoglycemia unawareness within the 6 months prior to screening. 3. History or presence of proliferative diabetic retinopathy, diabetic macular edema, or non-proliferative diabetic retinopathy that requires acute treatment. 4. Evidence of clinically significant/active nephropathy or neuropathy (including resting tachycardia, orthostatic hypotension, and diabetic diarrhea). 5. Current treatment or treatment within 3 months of screening with any other anti-hyperglycemic medication except metformin or SGLT-2 inhibitors. 6. Have obesity induced by other endocrinologic disorders (e.g., Cushing's syndrome) or diagnosed monogenetic or syndromic forms of obesity (e.g., melanocortin-4 receptor deficiency or Prader-Willi Syndrome). 7. Have a known, clinically significant gastric emptying abnormality. 8. Have poorly controlled hypertension at screening, untreated renal artery stenosis, or evidence of labile blood pressure including symptomatic postural hypotension. 9. Have any of the following cardiovascular conditions within 3 months prior to screening: Acute myocardial infarction; Cerebrovascular accident (stroke)/transient ischemic attack; Unstable angina; Hospitalization due to congestive heart failure.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: R07795081 administered at randomized experimental doses; Placebo taken matching the study drug; Active comparator Semaglutide administered at protocol-specified doses. All study drugs are evaluated based on a Once Daily (QD) administration model or

appropriate comparator dosing. **Route of Administration:** Oral.
Duration of Treatment: 30 weeks of continuous treatment.

5.2 Concomitant & Prohibited Therapy Permitted: Stable doses of metformin and/or SGLT-2 inhibitors; standard diet and exercise regimens. **Prohibited:** Use of other anti-hyperglycemic medications or other GLP-1 receptor agonists not supplied by the study.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, BMI, baseline HbA1c Labs
Baseline	Day 0	Randomization, First Dose
Interim	Weeks 4-24	Safety Labs, Efficacy Assessment (Glycemic control), Dose evaluation
End of Study	Week 30	Final Vital Signs, Final HbA1c measures, IP Accountability

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint 1. R07795081 vs. Placebo: Change in Glycated Hemoglobin (HbA1c) from Baseline at Week 30.

7.2 Secondary Endpoints 1. R07795081 vs. Semaglutide: Change in HbA1c from Baseline at Week 30. 2. Percentage of Participants with HbA1c $\leq 5.7\%$, $\leq 6.5\%$, and $\leq 7.0\%$ at Week 30. 3. Change in Fasting Plasma Glucose from Baseline at Week 30.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring of GI tolerability, glycemic events (e.g., potential hypoglycemia), and overall safety parameters via patient reporting and site visits. **Serious Adverse Events (SAEs):** Standard 24-hour reporting timeline to the sponsor. **Data Monitoring Committee (DMC):** Utilized for ongoing safety evaluation given the investigational dose-finding nature of Phase II.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) for primary efficacy; Safety Set for AE/SAE assessment. **Sample Size Justification:** A target enrollment of 240 participants powered to detect statistically significant dose-dependent reductions in HbA1c for R07795081 compared to placebo, and for evaluation against the Semaglutide active control arm. **Handling of Missing Data:** Mixed-effect Model Repeated Measure (MMRM) or Multiple Imputation for metabolic endpoint dropouts.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Routine on-site and remote monitoring visits to ensure protocol adherence and source data verification. **Audit**

Trail: Robust Electronic Data Capture (EDC) system with full audit trails for eCRF lock and data queries.

11. Study Identifiers & Governance

Primary ID: BP45703 / 2024-520322-11-00 **Posting ID:**

961137043236129959 **Registry Number:** NCT07112872 **Sponsor/Lead**

Organization: Hoffmann-La Roche (Roche) **Study Title:** A Study to Compare Different Doses of R07795081 With a Placebo or

Semaglutide in People With Type 2 Diabetes **Official Regulatory**

Title: A Randomized, Double-Blind, Placebo- and Open-Label Active Comparator- Controlled, Parallel-Group, Multi-Center Phase II

Study to Evaluate the Efficacy, Safety, and Tolerability of Once-Daily R07795081 Administered for 30 Weeks to Participants With

Type 2 Diabetes Mellitus

12. Clinical Framework (T2D Specifics)

Disease Area: Type 2 Diabetes Mellitus **Preferred Disease Name:**

Diabetes Mellitus, Non-Insulin-Dependent **Semantic Type:** Disease

or Syndrome **Clinical Definition:** A type of diabetes mellitus that is characterized by insulin resistance or desensitization and

increased blood glucose levels. This is a chronic disease that can develop gradually over the life of a patient and can be

linked to both environmental factors and heredity. **Associated**

Codes/Ontology: - **MeSH Code:** D003924 - **HPO Code:** HP:0005978 -

SNOMED ID: 44054006 (Hierarchy: Disease, Disorder of endocrine system, Metabolic disease, Disorder of glucose metabolism, etc.)

13. Study Procedures & Methodology

Management Pathway: Daily oral administration of study drug alongside standard T2D management (diet, exercise, metformin/

SGLT-2i). **Education Protocol (HCP):** Trial protocol adherence and AE management instructions. **Education Protocol (Patient):**

Nutritional counseling and medication adherence/dosing

instructions. **Quality Audit Mechanism:** Electronic diary tracking, pill counts, and regular site visits.

14. Observation & Follow-Up Schedule

Total Duration: 30 Weeks of Treatment **Onsite Visit Cadence:** Routine intervals defined by the protocol (typically Weeks 4, 8, 12, 20, 30 for metabolic studies). **Remote Monitoring Frequency:** Supplementary virtual check-ins for safety/dosing adherence where applicable. **Checklist Review Interval:** Routine review of concurrent medications and body weight.

15. Sign-off & Approval

Role	Name	Signature	Date		----	----	----	----
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]					Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]					