

1. Administrative & Study Identification

Full Study Title: A Study to Learn About the Effects of Two Study Medicines (Maplirpaccept [PF-07901801] And Glofitamab) When Given Together In People With Relapsed Or Refractory Diffuse Large B Cell Lymphoma **Official Title:** A Phase 1b/2, Open-Label Study of PF-07901801 in Combination with Glofitamab After a Fixed, Single Dose of Obinutuzumab in Participants with Relapsed/Refractory Diffuse Large B Cell Lymphoma Not Eligible for Stem Cell Transplantation **Short Title/Acronym:** Maplirpaccept + Glofitamab + Obinutuzumab in R/R DLBCL **Protocol Number:** C4971006 **NCT Number:** NCT05896163 **Sponsor/Company:** Pfizer **Trial Status:** Active, not recruiting **Investigational Product:** Maplirpaccept (PF-07901801) and Glofitamab, given after Obinutuzumab (Gazyva, ATC: L01FA03) **Phase of Development:** Phase 1b/2 **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: - Phase 1b: To evaluate the safety, tolerability, and determine dose-limiting toxicities (DLTs) to select the optimal doses of maplirpaccept when administered in combination with glofitamab. - Phase 2: To evaluate safety and efficacy (objective response) to determine the recommended Phase 3 dose of maplirpaccept to be administered in combination with glofitamab.

Secondary Objectives: To evaluate frequency of adverse events, frequency of clinical laboratory abnormalities, potential clinical benefits, pharmacokinetics, progression-free survival (PFS), overall response rate, and overall survival.

2.2 Background & Rationale To address the critical unmet medical need in patients with relapsed or refractory diffuse large B-cell lymphoma (R/R DLBCL) who are not eligible for or unwilling to undergo high-dose chemotherapy and subsequent autologous stem cell transplantation (ASCT) or unable to receive approved chimeric antigen receptor T-cell (CAR-T) therapy. Maplirpaccept (PF-07901801) is an anti-CD47 fusion protein designed to enhance phagocytosis and antitumor activity via inhibition of CD47-mediated signaling. Combining maplirpaccept with a CD20xCD3 T-cell-engaging bispecific antibody (glofitamab) following a single pretreatment dose of obinutuzumab is hypothesized to redirect T-cells against malignant cells and enhance the immune response.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm (Phase 1b Dose Escalation) and Randomized Dose-ranging (Phase 2) **Blinding:** Open-label **Randomization:** Sequential assignment in Phase 1b; Randomized (to 1 of 3 maplirpacept dose levels) in Phase 2.

3.2 Planned Enrollment & Duration Target **\$N\$:** 24 participants
Number of Sites: Global Multicenter Study (including United States, Australia, Israel, Japan, Taiwan) **Estimated Study Start:** 30-Aug-2023 **Estimated Primary Completion:** 31-Mar-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Histologically confirmed diagnosis of DLBCL. 2. Relapsed or refractory disease (must have received at least 2 prior lines of treatment for NHL; Phase 2 requires at least 2 and no more than 4 prior systemic regimens). Prior therapy must include an anti-CD20 antibody. 3. Unable or unwilling to undergo a stem cell transplant and/or unable to receive CAR-T cell therapy. 4. Eastern Cooperative Oncology Group (ECOG) performance status ≤ 2 . 5. Adequate bone marrow, hepatic and renal function.

4.2 Exclusion Criteria 1. Prior treatment with anti-CD47 and/or prior glofitamab or anti-CD20xCD3 containing regimen. Refractoriness to an obinutuzumab monotherapy containing regimen. 2. Prior allogeneic stem cell transplantation or autologous stem cell transplantation within 12 weeks prior to enrolment. 3. Known active bacterial, viral, fungal, mycobacterial, parasitic, or other infection. 4. High Grade B-Cell Lymphoma.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Cycle 0 features a single pre-treatment dose of obinutuzumab (trade name: Gazyva) followed by 2 step-up doses of glofitamab. In Cycle 1 Day 1, combination of maplirpacept with full-dose glofitamab begins. Maplirpacept is given weekly for the first 3 cycles, then every 3 weeks. Glofitamab is given every 3 weeks for up to 11 cycles. **Route of Administration:** Intravenous (IV) Infusion **Duration of Treatment:** 21-day cycles. Glofitamab is given for approximately 9 months. Maplirpacept continues as monotherapy thereafter until disease progression or unacceptable toxicity.

5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care for managing infusion-related reactions or cytokine release syndrome (CRS). **Prohibited:** Concurrent investigational agents or other anti-cancer/chemotherapy regimens.

6. Study Timeline & Visit Schedule

| Visit ID | Window (Days) | Key Activities/Procedures | | :---
| :--- | :--- | | **Screening** | Day -28 to Day -1 | Informed
Consent, Medical History, Scans/Imaging, Baseline Labs | | **Cycle**
0 | Pre-Treatment | Obinutuzumab pretreatment, Glofitamab step-up
doses (mitigate CRS) | | **Cycle 1 Day 1** | Week 0 | First dose
Maplirpacept + full dose Glofitamab, PK/safety monitoring | |
Interim | Q3W | 21-day cycles: Safety Labs, Efficacy Assessment,
IV Infusions | | **End of Study** | Follow-up | Final Vital Signs,
Exit Interview, Survival Follow-up |

7. Outcome Measures (Endpoints)

7.1 Primary Outcomes - Phase 1b: Number of participants with Dose limiting toxicities (DLT). DLTs are a predefined set of adverse events that are at least possibly related to any or all of the investigational agents. - **Phase 2:** Objective Response (OR). Defined as proportion of participants with objective response based assessment of complete response (CR) or partial response (PR) according to Lugano Response Classification Criteria 2014.

7.2 Secondary & Exploratory Outcomes - Phase 1b and Phase 2: Frequency of adverse events (AE). - **Phase 1b and Phase 2:** Frequency of clinical laboratory abnormalities. - **Phase 1b:** Objective Response (OR). - **Exploratory:** Progression-Free Survival (PFS), Overall Survival (OS), and characterization of pharmacokinetics (PK) of Maplirpacept.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Patient monitoring prioritizing standard AEs, laboratory abnormalities, as well as immunotherapy-specific toxicities, primarily Cytokine Release Syndrome (CRS) and Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS). **Serious Adverse Events (SAEs):** Reported standardly within 24 hours of site awareness. **Data Monitoring Committee (DMC):** Internal/External safety review committees governing dose escalation and cohort expansions.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) for efficacy endpoints; Safety Set for DLT and AE/SAE monitoring. **Sample Size Justification:** $N = 24$, sized according to the FDA "Project Optimus" initiative for dose optimization and selection in oncology. **Handling of Missing Data:** Standard survival analysis

metrics (e.g., right-censoring for PFS) at the last evaluable tumor assessment.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Regular onsite and remote monitoring corresponding to cycle intervals. **Audit Trail:** eCRF entry tracking, data clarification forms, and standard clinical trial auditing protocols.

11. Study Identifiers & Governance

Posting ID: 962057060083113980 **Primary ID:** C4971006 **Registry Number:** NCT05896163 **Sponsor/Lead Organization:** Pfizer **Study Title:** A Study to Learn About the Effects of Two Study Medicines (Maplirpacept [PF-07901801] And Glofitamab) When Given Together In People With Relapsed Or Refractory Diffuse Large B Cell Lymphoma **Official Regulatory Title:** A Phase 1b/2, Open-Label Study of PF-07901801 in Combination with Glofitamab After a Fixed, Single Dose of Obinutuzumab in Participants with Relapsed/Refractory Diffuse Large B Cell Lymphoma Not Eligible for Stem Cell Transplantation **Trial Status:** Active, not recruiting

12. Clinical Framework (Lymphoma Specific)

Primary Condition: Diffuse Large B-Cell Lymphoma (DLBCL)

Diagnosis Threshold: Histologically confirmed relapsed or refractory DLBCL **Medical Methodology:** Immune targeted therapy and CD20xCD3 T-cell engaging bispecific antibodies **Optimization**

Target: Dose optimization and Phase 3 dose selection (per FDA Project Optimus)

13. Study Procedures & Methodology

Management Pathway: Pretreatment with obinutuzumab and glofitamab step-up dosing to mitigate cytokine release syndrome. **Education**

Protocol (HCP): Site initiation visits and ongoing safety protocol training for CRS and ICANS monitoring. **Education**

Protocol (Patient): Education on signs and symptoms of immune-related adverse events. **Quality Audit Mechanism:** Independent safety reviews to adjudicate DLTs prior to cohort dose escalation.

14. Observation & Follow-Up Schedule

Total Duration: Follow-up for progressive disease or survival (~up to 3 years from enrollment). **Onsite Visit Cadence:** Every 21-day treatment cycle (Day 1 of each cycle and weekly during Cycle 1). **Remote Monitoring Frequency:** Inter-cycle continuous monitoring as required for AE tracking. **Checklist Review Interval:** Disease assessment/scans evaluated at pre-defined cycle intervals.

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]				Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]				