

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A PHASE 1B, OPEN-LABEL STUDY OF ELRANATAMAB IN COMBINATION WITH CARFILZOMIB PLUS DEXAMETHASONE AND ELRANATAMAB IN COMBINATION WITH PF-07901801 IN PARTICIPANTS WITH RELAPSED REFRACTORY MULTIPLE MYELOMA
Short Title:	A Clinical Trial of Four Medicines (Elranatamab Plus Carfilzomib and Dexamethasone or Maplirpacept) in People With Relapsed Refractory Multiple Myeloma
Protocol Number:	1055316693966694070
NCT Number:	NCT05675449
Sponsor Name:	Pfizer
Investigational Product:	carfilzomib
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Part 1 Number of participants with dose limiting toxicity (DLT): Dose limiting toxicity rate based on dose limiting toxicity evaluable participants. 2. Part 2A Number of participants with dose limiting toxicity: Dose limiting toxicity based on dose limiting toxicity evaluable participants. 3. Part 2B Number of participants with dose limiting Toxicity: Dose limiting toxicity rate based on dose limiting toxicity evaluable participants.
Secondary Obj.:	1. Part 1: Number of Participants with Treatment Emergent Adverse Events (TEAE) by Seriousness and Relationship to Treatment 2. Part 1: Number of Participants with Adverse Events (AE) characterized by type, frequency, severity. 3. Part 1: Number of Participants with Clinically Significant Change From Baseline in Laboratory Abnormalities

2.2 Background & Rationale

The main purpose of the study is to evaluate the safety and tolerability of the combination of elranatamab and carfilzomib and dexamethasone or elranatamab and maplirpacept.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, Status: RECRUITING
Target N:	90

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Prior diagnosis of multiple myeloma as defined by IMWG criteria.
- * Measurable disease based on IMWG criteria as defined by at least 1 of the following:
 - * Serum M-protein ≥ 0.5 g/dL.
 - * Urinary M-protein excretion ≥ 200 mg/24 hours.
 - * Serum immunoglobulin FLC ≥ 10 mg/dL (≥ 100 mg/L) AND abnormal serum immunoglobulin kappa to lambda FLC ratio (< 0.26 or > 1.65).
- * Part 1: Received at least 1 but not more than 3 prior lines of therapy for multiple myeloma (induction therapy followed by stem cell transplant and consolidation/maintenance therapy will be considered as 1 line of therapy).
- * Part 2: Received at least 3 prior lines of therapy for multiple myeloma who are refractory to at least one IMiD, one PI and one anti-CD38 antibody.
- * ECOG performance status 0-1.
- * Resolved acute effects of any prior therapy to baseline severity or CTCAE Grade ≤ 1 .
- * Not pregnant or breastfeeding and willing to use contraception.

4.2 Exclusion Criteria

- * Plasma cell leukemia, Smouldering MM, Waldenströms macroglobulinemia, Amyloidosis, POEMS Syndrome, Primary refractory MM
- * Impaired cardiovascular function or clinically significant cardiovascular diseases.
- * Participants with any active, uncontrolled bacterial, fungal, or viral infection.
- * Stem cell transplant within 12 weeks prior to enrollment, or active graft versus host disease.
- * Any other active malignancy within 3 years prior to enrollment, except for adequately treated basal cell or squamous cell skin cancer, or carcinoma in situ.
- * Part 1: Previous treatment with a BCMA-directed therapy.
- * Part 2: Previous treatment with any anti-BCMA directed therapy, with the exception of CAR-T. Previous treatment with a CD47-SIRP alpha-directed therapy.
- * Part 1: Prior treatment with carfilzomib
- * Live attenuated vaccine within 4 weeks of the first dose of study intervention.
- * Administration with an investigational product (e.g. drug or vaccine) concurrent with study intervention or within 30 days preceding the first dose of study intervention used in this study.
- * Any of the following within 3 months of enrollment: erosive esophagitis, treatment resistant peptic ulcer, infectious or inflammatory bowel disease, pulmonary embolism or uncontrolled thromboembolic event.
- * Intolerance to or participants who have had a severe (Grade ≥ 3) allergic or anaphylactic reaction to antibodies or therapeutic proteins

11. Study Identifiers & Governance

Primary ID:	1055316693966694070
Registry Number:	NCT05675449
Sponsor:	Pfizer
Study Title:	A Clinical Trial of Four Medicines (Elranatamab Plus Carfilzomib and Dexamethasone or Maplirpacept) in People With Relapsed Refractory Multiple Myeloma

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____