

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1 Study of Sigvotatug Vedotin in Advanced Solid Tumors
Short Title:	A Study of Sigvotatug Vedotin in Advanced Solid Tumors
Protocol Number:	199880777510017399
NCT Number:	NCT04389632
Sponsor Name:	Pfizer
Investigational Product:	cisplatin
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of participants with adverse events (AEs): Any untoward medical occurrence in a clinical investigational participant administered a medicinal product and which does not necessarily have a causal relationship with this treatment. 2. Number of patients with laboratory abnormalities 3. Number of participants with dose-limiting toxicities (DLTs)
Secondary Obj.:	1. Confirmed objective response rate (ORR) per Response Evaluation Criteria in Solid Tumors (RECIST) v1.1 by investigator assessment 2. Duration of objective response (DOR) per RECIST v1.1 by investigator assessment 3. Progression-free survival (PFS) per RECIST v1.1 by investigator assessment

2.2 Background & Rationale

This trial will look at a drug called sigvotatug vedotin (SGN-B6A) alone and with pembrolizumab, with or without chemotherapy, to find out whether it is safe for people who have solid tumors. It will study sigvotatug vedotin to find out what its side effects are. A side effect is anything the drug does besides treating cancer. It will also study whether sigvotatug vedotin works to treat solid tumors.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, Status: RECRUITING
Target N:	1006

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

* Disease indication

* Participants must have histologically or cytologically confirmed metastatic or unresectable solid malignancy within one of the tumor types listed below (dependent on study part).

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Non-small cell lung cancer (NSCLC)
- * Head and neck squamous cell cancer (HNSCC)
- * Advanced HER2-negative breast cancer
- * Esophageal squamous cell carcinoma (ESCC)
- * Esophageal/Gastro-esophageal junction adenocarcinoma (EAC/GEJ)
- * Cutaneous squamous cell cancer (cSCC)
- * Exocrine pancreatic adenocarcinoma
- * Bladder cancer
- * Cervical cancer
- * Gastric cancer
- * High grade serous ovarian cancer (HGSOC)
- * Part A only: Participants must have disease that is relapsed or refractory or be intolerant to standard-of-care therapies and should have no appropriate standard-of-care therapeutic options.
- * Part B only: Participants must have disease that is relapsed or refractory or be intolerant to standard-of-care therapies. Participants must have received platinum-based therapy and a PD-1/PD-(L)1 inhibitor, if applicable and available.
- * Part C only: For pembrolizumab combination cohorts, participants must be eligible for pembrolizumab per local standard of care. For pembrolizumab with cisplatin or carboplatin, participants must be eligible for both pembrolizumab and the platinum agent per local standard of care. Participants must be treatment naïve for locally advanced or metastatic systemic therapy (prior definitively intended or \[neo\]adjuvant therapy is allowed).
- * Part D only: Participants must be treatment naïve for locally advanced or metastatic systemic therapy.
- * Participants enrolled in the following study parts should have a tumor site accessible for biopsy and agree to biopsy as follows:
 - * Disease-specific expansion cohorts (Part B and Part D): A baseline fresh tumor biopsy is required. An archival biopsy collected within 90 days prior to first dose of study drug may be used.
 - * Biology expansion cohort: pretreatment biopsy and on-treatment (Cycle 1) biopsy
- * An Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 or 1
- * Measurable disease per the RECIST v1.1 at baseline

Exclusion Criteria

- * History of another malignancy within 3 years before first dose of study drug, or any evidence of residual disease from a previously diagnosed malignancy. Exceptions are malignancies with a negligible risk of metastasis or death.
- * Known active central nervous system metastases. Participants with previously treated brain metastases may participate provided they:
 - * are clinically stable for at least 4 weeks prior to study entry after brain metastasis treatment,
 - * have no new or enlarging brain metastases, and

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* are off of corticosteroids prescribed for symptoms associated with brain metastases for at least 7 days prior to first dose of study drug.

* In Part D, participants with untreated, asymptomatic CNS metastases smaller than 1 cm may be enrolled without definitive treatment as long as they have no neurological symptoms, no or minimal surrounding edema, and no requirements for corticosteroids.

* Carcinomatous meningitis

* Previous receipt of an MMAE-containing agent or an agent targeting integrin beta-6

* Pre-existing neuropathy Grade 1 or greater per the National Cancer Institute's Common Terminology Criteria for Adverse Events, version 5.0 (NCI CTCAE v5.0) for Parts C and D cohorts with cisplatin or carboplatin; Grade 2 or greater per the NCI CTCAE v5.0 for all other cohorts

* Any uncontrolled Grade 3 or higher (per NCI CTCAE v5.0) viral, bacterial, or fungal infection within 2 weeks prior to the first dose of sigvotatug vedotin.

* Routine antimicrobial prophylaxis is permitted

* Grade ≥3 pulmonary disease unrelated to underlying malignancy. This includes clinically severe pulmonary function compromise resulting from clinically significant pulmonary illnesses

* Part C and D: Prior therapy with a PD-1 inhibitor, anti-PD-(L)1, or anti PD-L2 agent or with an agent directed to another stimulatory or co-inhibitory T-cell receptor and was discontinued from that treatment due to a Grade 3 or higher immune-mediated adverse event (IMAE).

* History of noninfectious interstitial lung disease (ILD) or pneumonitis that required steroids, current ILD or pneumonitis, or suspected ILD or pneumonitis that cannot be ruled out by imaging at screening

* Known diffusing capacity of the lung for carbon monoxide (DLCO; adjusted for hemoglobin) <50% predicted

* Investigator site staff directly involved in the conduct of the study and their family members, site staff otherwise supervised by the investigator, and sponsor and sponsor delegate employees directly involved in the conduct of the study and their family members.

4.2 Exclusion Criteria

Not Provided

11. Study Identifiers & Governance

Primary ID: 199880777510017399

Registry Number: NCT04389632

Sponsor: Pfizer

Study Title: A Study of Sigvotatug Vedotin in Advanced Solid Tumors

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____