

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1, Open-label, Multicenter Study of the Safety, Pharmacokinetics, and Pharmacodynamics of MK-4464 as Monotherapy and in Combination With Pembrolizumab in Participants With Advanced/Metastatic Solid Tumors
Short Title:	Study of MK-4464 as Monotherapy and in Combination With Pembrolizumab in Participants With Advanced/Metastatic Solid Tumors (MK-4464-001)
Protocol Number:	611386539871345332
NCT Number:	NCT05514444
Sponsor Name:	Merck
Investigational Product:	pembrolizumab
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants Experiencing Dose-Limiting Toxicities (DLTs): A DLT is any toxicity assessed by the investigator to be possibly, probably, or definitely related to study intervention administration that results in a change to a given dose or a delay in initiating the next cycle. All toxicities will be graded using National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE). 2. Number of Participants Who Experience At Least One adverse event (AE): An AE is defined as any untoward medical occurrence in a participant, temporally associated with the use of study treatment, whether or not considered related to the study treatment. The number of participants who experience at least one AE will be presented. 3. Number of Participants Who Discontinue Study Treatment Due to an AE: An AE is defined as any untoward medical occurrence in a participant, temporally associated with the use of study treatment, whether or not considered related to the study treatment. The number of participants who discontinue study treatment due to an AE will be presented.
Secondary Obj.:	1. Minimum Plasma Concentration (Cmin) of MK-4464 2. Maximum Plasma Concentration (Cmax) of MK-4464 3. Area Under the Plasma Concentration-Time Curve (AUC) of MK-4464

2.2 Background & Rationale

The purpose of this study is to assess the safety, pharmacokinetics, and preliminary efficacy of MK-4464 as monotherapy and in combination with pembrolizumab in participants with advanced/metastatic solid tumors.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, Status: COMPLETED
-------------------	--

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Target N: 64

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The key Inclusion Criteria include but are not limited to the following:

- * Have a histologically or cytologically confirmed advanced/metastatic solid tumor by pathology report and have received, been intolerant to, been ineligible for, or refused all treatment known to confer clinical benefit
- * Must submit a baseline tumor sample for analysis
- * Have a performance status of 0 or 1 on the Eastern Cooperative Oncology Group (ECOG) Performance Scale
- * Human immunodeficiency virus (HIV) infected participants must have well controlled HIV on antiretroviral therapy (ART)
- * Participants who are HBsAg positive are eligible if they have received HBV antiviral therapy for at least 4 weeks, and have undetectable HBV viral load before randomization.

4.2 Exclusion Criteria

The key Exclusion Criteria include but are not limited to the following:

- * Has had chemotherapy, definitive radiation, or biological cancer therapy within 4 weeks (2 weeks for palliative radiation) before the first dose of study intervention or has not recovered to CTCAE Grade 1 or better from any AEs that were due to cancer therapeutics administered more than 4 weeks earlier
- * Has a history of a second malignancy, unless potentially curative treatment has been completed with no evidence of malignancy for 2 years
- * Has clinically active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Has an active infection requiring therapy
- * History of an allogenic stem cell transplant or a solid organ transplant
- * Has a history of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease
- * Has an active autoimmune disease that has required systemic treatment in the past 2 years
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease
- * Has known psychiatric or substance abuse disorders that would interfere with the participant's ability to cooperate with the requirements of the study
- * Has not fully recovered from any effects of major surgery without significant detectable infection
- * Has received radiation therapy to the lung that is ≥ 30 gray (Gy) within 6 months of the first dose of study treatment
- * Received a live or live-attenuated vaccine within 30 days before the first dose of study intervention
- * Is currently participating and receiving study intervention in a study of an investigational agent or has participated and received study intervention in a study of an investigational agent or has used an investigational device within 28 days

11. Study Identifiers & Governance

Primary ID: 611386539871345332

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Registry Number: NCT05514444

Sponsor: Merck

Study Title: Study of MK-4464 as Monotherapy and in Combination With Pembrolizumab in Participants With Advanced/Metastatic Solid Tumors (MK-4464-001)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____