

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1, Open-label, Non-comparative, Multicenter Clinical Study to Evaluate the Safety, Tolerability, and Pharmacokinetics of Ceftolozane/Tazobactam (MK-7625A) in Pediatric Participants With Nosocomial Pneumonia
Short Title:	Safety and Pharmacokinetics of Ceftolozane/Tazobactam in Pediatric Participants With Nosocomial Pneumonia (MK-7625A-036)
Protocol Number:	684063392650693124
NCT Number:	NCT04223752
Sponsor Name:	Merck
Investigational Product:	tazobactam
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of Participants With Any Adverse Events (AEs): An AE was defined as any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The percentage of participants experiencing any AE was reported for each arm. 2. Percentage of Participants With Any Serious AEs (SAEs): An SAE was defined as any untoward medical consequence that, at any dose, results in death, is life-threatening, requires inpatient hospitalization or prolongs existing hospitalization, results in persistent or significant disability/incapacity, is a congenital anomaly/birth defect, or any other important medical event. The percentage of participants with any SAE was reported for each arm. 3. Percentage of Participants With Any Drug-related AEs: A drug-related AE was defined as any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, and considered related to the study intervention. The percentage of participants with any drug related AEs was reported for each arm.
Secondary Obj.:	1. Plasma Concentrations of Ceftolozane 2. Area Under the Concentration-time Curve of an 8-hour Dosing Interval (AUC ₀₋₈) of Plasma Ceftolozane 3. Maximum Observed Concentration During a Dosage Interval (C _{max}) of Plasma Ceftolozane

2.2 Background & Rationale

This is a phase 1, open-label, non-comparative, multicenter clinical study to evaluate the safety, tolerability, and pharmacokinetics of ceftolozane/tazobactam (MK-7625A) in pediatric participants with nosocomial pneumonia (NP).

3. Study Design & Methodology

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, Status: COMPLETED
Target N: 41

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Is hospitalized and anticipated to receive a minimum of 8 days of concomitant standard-of-care \[SOC\] antibiotic therapy for proven or suspected NP.
- * If male, is abstinent from heterosexual intercourse, or agrees to use contraception during the intervention period and for ?30 days after the last dose of study intervention.
- * If female, is not pregnant or breastfeeding, or is not a woman of childbearing potential (WOCBP), or is a WOCBP using acceptable contraception, is a WOCBP with negative urine or serum pregnancy test within 48 hours of the first dose of study intervention, or is abstinent from heterosexual intercourse.

4.2 Exclusion Criteria

- * Has a documented history of any moderate or severe hypersensitivity (or allergic) reaction to any ?-lactam antibacterial.
- * Participants 3 months to \<18 years of age: has moderate to severe impairment of renal function, defined as an estimated creatinine clearance (CrCL) \<50 mL/min/1.73 m² based on the revised Schwartz equation or requirement for peritoneal dialysis, hemodialysis, or hemofiltration.
- * Participants \<3 months of age: has CrCL \<20 mL/min/1.73 m² based on the revised Schwartz equation or requirement for peritoneal dialysis, hemodialysis, or hemofiltration.
- * Is receiving or is anticipated to receive piperacillin/tazobactam while receiving ceftolozane/tazobactam or has received piperacillin/tazobactam within 24 hours prior to the first dose of ceftolozane/tazobactam.
- * Has participated in any clinical study of a therapeutic investigational product within 30 days prior to the first dose of ceftolozane/tazobactam.
- * Has previous participation in any study of ceftolozane or ceftolozane/tazobactam.
- * Has any condition or circumstance that, in the opinion of the investigator, would compromise the safety of the participant or the quality of study data.
- * Has any rapidly progressing disease or immediately life-threatening illness including acute hepatic failure or septic shock.
- * Has active immunosuppression.

11. Study Identifiers & Governance

Primary ID: 684063392650693124
Registry Number: NCT04223752
Sponsor: Merck
Study Title: Safety and Pharmacokinetics of Ceftolozane/Tazobactam in Pediatric Participants With Nosocomial Pneumonia (MK-7625A-036)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____