

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1, Open-label, Multicenter Study to Assess Safety, Tolerability, PK, and Efficacy of MK-1084 as Monotherapy and as Part of Various Combination Therapies in Participants With KRAS G12C Mutant Advanced Solid Tumors
Short Title:	A Study of MK-1084 in KRAS Mutant Advanced Solid Tumors (MK-1084-001)
Protocol Number:	854145671623249515
NCT Number:	NCT05067283
Sponsor Name:	Merck
Investigational Product:	fluorouracil
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants Who Experience a Dose-Limiting Toxicity (DLT): A DLT is defined as an event with toxicity including the type, severity, time of onset, time of resolution, and the probable association with study treatment that are not due to pre-existing conditions as defined by the Common Terminology Criteria for Adverse Events Version 5.0 (CTCAE 5.0). Number of participants who experience a DLT will be reported. 2. Number of Participants Who Experience an Adverse Event (AE): An AE is any unfavorable and unintended sign, symptom, or disease temporally associated with the use of a medicinal product or protocol-specified procedure, whether or not considered related to the medicinal product or protocol-specified procedure. Any worsening of a preexisting condition that is temporally associated with the use of the Sponsor's product, is also an AE. The number of participants who experience an AE will be reported. 3. Number of Participants Who Discontinue Study Treatment Due to an AE: An AE is any unfavorable and unintended sign, symptom, or disease temporally associated with the use of a medicinal product or protocol-specified procedure, whether or not considered related to the medicinal product or protocol-specified procedure. Any worsening of a preexisting condition that is temporally associated with the use of the Sponsor's product, is also an AE. The number of participants who discontinue study treatment due to an AE will be reported.
Secondary Obj.:	1. Objective Response Rate (ORR) 2. Duration of Response (DOR) 3. Mean Plasma Concentration of MK-1084

2.2 Background & Rationale

This is a study evaluating the safety, pharmacokinetics, and efficacy of MK-1084 alone, and MK-1084 plus other combination therapies in participants with advanced solid tumors with identified kirsten rat sarcoma viral oncogene

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

homolog G12C (KRAS G12C) mutation.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, Status: RECRUITING

Target N: 830

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

For all participants:

- * Has measurable disease by RECIST 1.1 criteria
- * Has adequate organ function
- * Male participants agree to protocol-specified contraception requirements including refraining from donating sperm and using protocol-specified contraceptives unless confirmed to be azoospermic
- * Female participants must not be pregnant or breastfeeding, and must agree to protocol-specified contraceptive requirements and must have a negative highly sensitive pregnancy test within 24 hours (for a urine test) or 72 hours (for a serum test) before the first dose of study intervention

For Arm 1 - Has locally advanced unresectable or metastatic solid-tumor malignancy with histologically OR blood-based confirmation of KRAS G12C mutation who has received at least 1 line of therapy for systemic disease

For Arm 2

Has an untreated metastatic non-small cell lung cancer (NSCLC) with histologically OR blood-based confirmation of KRAS G12C mutation and histologic confirmation of programmed cell death ligand 1 (PD-L1) tumor proportion score (TPS) ≥1%

For Arm 3

- * Has locally advanced unresectable or metastatic solid-tumor malignancy with histologically or blood-based confirmation of KRAS G12C mutation who has received at least 1 line of therapy for systemic disease Expansion Group A: 2L+NSCLC
- * Has histologically or cytologically confirmed diagnosis of unresectable or metastatic NSCLC with histological or blood-based confirmation of KRAS G12C mutation and submits archival tumor sample
- * Previous treatment failure of at least 1 line of systemic therapy Expansion Group B
- * Has locally advanced unresectable or metastatic solid-tumor malignancy, excluding NSCLC or CRC, with histologically or blood-based confirmation of KRAS G12C mutation who has received at least 1 line of therapy for systemic disease

Arm 4 only - Has an untreated advanced or metastatic nonsquamous NSCLC with histologically or blood-based confirmation of KRAS G12C mutation

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Arm 5 only

- * Histologically or cytologically confirmed diagnosis of locally advanced unresectable or metastatic colorectal adenocarcinoma and with histologically or blood-based confirmation of KRAS G12C mutation
- * Previous treatment failure of one or 2 previous line(s) of systemic therapy

Arm 6 only

\- Locally advanced unresectable or metastatic colorectal adenocarcinoma with histologically or blood-based confirmation of KRAS G12C mutation

4.2 Exclusion Criteria

- * Has received chemotherapy, definitive radiation, or biological cancer therapy within 4 weeks (2 weeks for palliative radiation) before first dose of study intervention
- * Has a history of second malignancy, unless potentially curative treatment has been completed with no evidence of malignancy for 5 years
- * Has clinically active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Has an active infection requiring systemic therapy
- * Known history of HIV infection or. has a known history of Hepatitis B virus or known active Hepatitis C virus infection
- * Has a history of interstitial lung disease, noninfectious pneumonitis requiring active steroid therapy, or ongoing pneumonitis
- * Has an active autoimmune disease requiring systemic therapy
- * Has not fully recovered from any effects of major surgical procedure without significant detectable infection
- * Has one or more of the following ophthalmological findings/conditions: intraocular pressure ≥ 21 mm Hg and/or any diagnosis of glaucoma; diagnosis of central serous retinopathy, retinal vein occlusion, or retinal artery occlusion and/or a diagnosis of retinal degenerative disease
- * Has received live or live-attenuated vaccine within 4 weeks of study start

Arm 4 Only

- * Is unable to interrupt aspirin or other nonsteroidal anti-inflammatories (NSAIDs), other than an aspirin dose ≥ 1.3 grams per day, for at least 2 days (5 days for long-acting agents \[for example, piroxicam\]) before, during, and for at least 2 days after administration of pemetrexed.
- * Is unable/unwilling to take folic acid, vitamin B12, and dexamethasone

11. Study Identifiers & Governance

Primary ID:	854145671623249515
Registry Number:	NCT05067283
Sponsor:	Merck
Study Title:	A Study of MK-1084 in KRAS Mutant Advanced Solid Tumors (MK-1084-001)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____