

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	Korean Post-marketing Surveillance for Xeljanz XR (Registered)
Short Title:	Korean Post-marketing Surveillance for Xeljanz XR
Protocol Number:	85643817693988424
NCT Number:	NCT04876781
Sponsor Name:	Pfizer
Investigational Product:	tofacitinib
Phase:	Not Applicable
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants With Treatment Emergent Adverse Events (AEs): AE is any untoward medical occurrence attributed to study drug in a participant who received study drug. Serious adverse event (SAE) is an AE resulting in any of the following outcomes or deemed significant for any other reason: death; initial or prolonged inpatient hospitalization; life-threatening experience (immediate risk of dying); persistent or significant disability/incapacity; congenital anomaly. Treatment-emergent are events between first dose of study drug and up to end of this PMS after last dose that were absent before treatment or that worsened relative to pretreatment state. Relatedness to Xeljanz XR was assessed by the investigator (Yes/No). Participants with multiple occurrences of an AE within a category were counted once within the category.
Secondary Obj.:	1. Change from baseline in Disease Activity Score based on 28-joints Count (DAS28) 2. Percentage of Participants With European League Against Rheumatism (EULAR) Response Based on DAS28 3. Number of Participants With an American College of Rheumatology 20% (ACR20) Response

2.2 Background & Rationale

Xeljanz XR extended-release tablets 11 mg (Tofacitinib citrate) is a drug subject to the risk management plan in accordance with Article 4-1-11 of the "Regulation on Safety of Medicinal Products, etc." in Korea. As part of additional pharmacovigilance activity, this Post-marketing Surveillance (PMS) was planned to evaluate safety and effectiveness of Xeljanz XR under routine clinical practice. At least 200 patients with Rheumatoid Arthritis, Psoriatic Arthritis, or Ankylosing Spondylitis who were treated with Xeljanz XR will be enrolled about four years.

3. Study Design & Methodology

Design Framework:	Type: OBSERVATIONAL, Status: RECRUITING
Target N:	200

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Patients must meet all of the following inclusion criteria to be eligible for inclusion in the study:

1. Xeljanz® XR is administered according to indications on the approved labeling

* Treatment of moderate to severe active RA in adult patients (18 years of age or older) who have had an inadequate response or are intolerant to methotrexate

* In combination with methotrexate, treatment of active psoriatic arthritis (PsA) in adult patients (18 years of age or older) who have had an inadequate response or who have been intolerant to a prior DMARD therapy.

* Treatment of active ankylosing spondylitis (AS) in adult patients (18 years of age or older) who have responded inadequately to conventional therapy

* In the following patients, Xeljanz® XR should be used only for patients who have had an inadequate response or are intolerant to the existing treatment.

A. Patients aged 65 or older B. Cardiovascular high-risk patients C. Patients at risk for malignancy

2. Patients who have previously been given Xeljanz 5mg, who have changed Xeljanz® XR, are also eligible for registration in the study

3. Evidence of a personally signed and dated informed consent document indicating that the patient (or a legally acceptable representative) has been informed of all pertinent aspects of the study

4.2 Exclusion Criteria

Patients meeting any of the following criteria will not be included in the study:

1. Patients with a history of hypersensitivity to any ingredients of this product.

2. Patients with serious infection (sepsis, etc.) or active infection including localized infection.

3. Patients with active tuberculosis.

4. Patients with severe hepatic function disorder.

5. Patients with an absolute neutrophil count (ANC) $<1,000$ cells/mm³. *

6. Patients with a lymphocyte count <500 cells/mm³. *

7. Patients with a hemoglobin level <9 g/dL. *

8. Pregnant or possibly pregnant women. * Do not initiate Xeljanz XR in the following cases: ANC; absolute neutrophil count $<1,000$ cells/mm³ ALC; absolute lymphocyte count <500 cells/mm³ Hemoglobin <9 g/dL

11. Study Identifiers & Governance

Primary ID: 85643817693988424

Registry Number: NCT04876781

Sponsor: Pfizer

Study Title: Korean Post-marketing Surveillance for Xeljanz XR

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____