

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1, Randomized, Single-blind, Placebo-controlled Study to Investigate the Pharmacokinetics, Safety, Tolerability, and Efficacy of AZD0780 in Combination With Ezetimibe, Ezetimibe/Rosuvastatin, or Ezetimibe/Bempedoic Acid in Healthy Male and Female Participants 18 to 75 Years of Age With Elevated LDL-C.
Short Title:	A Study to Investigate the Pharmacokinetics, Safety, Tolerability, and Efficacy of AZD0780 With Ezetimibe Combinations in Healthy Adults With Elevated LDL-C.
Protocol Number:	869235349604422440
NCT Number:	NCT06742853
Sponsor Name:	AstraZeneca
Investigational Product:	rosuvastatin
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Change from baseline in direct Low-density Lipoprotein Cholesterol (LDL-C): To evaluate the effect of AZD0780 on LDL-C levels versus placebo when dosed with ezetimibe or ezetimibe and rosuvastatin or ezetimibe and bempedoic acid. 2. Number of participants with adverse events: The safety and tolerability of AZD0780 when dosed with ezetimibe or ezetimibe and rosuvastatin or ezetimibe and bempedoic acid will be assessed.
Secondary Obj.:	1. Plasma concentration of AZD0780 2. Area under concentration-curve from time 0 to the last quantifiable concentration (AUClast) 3. Area under concentration-time curve in the dosing interval (AUCtau) (Day 15)

2.2 Background & Rationale

The main aim of this study is to assess the effects of AZD0780 when added on top of ezetimibe or ezetimibe and rosuvastatin or ezetimibe and bempedoic acid.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, Status: COMPLETED
Target N:	81

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

1. Signed the informed consent form before any study-related procedure.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

2. All females must have a negative serum pregnancy test at the Screening Visit and on admission to the Clinical Unit
3. Females of non-childbearing potential must be confirmed at the Screening Visit by fulfilling one of the following criteria: postmenopausal or surgically sterilized females.
4. Have a Body Mass Index (BMI) $\geq 18 \text{ kg/m}^2$ and weigh at least 50 kg.
5. Sexually active fertile male participants with partners of childbearing potential must adhere to the contraception methods.
6. Fasting LDL-C $\geq 100 \text{ mg/dL}$ but $\leq 190 \text{ mg/dL}$ ($\geq 2.6 \text{ mmol/L}$ but $\leq 4.9 \text{ mmol/L}$ for London EPCU) at the Screening Visit.
7. Fasting triglycerides $\leq 400 \text{ mg/dL}$ (or $\leq 10.3 \text{ mmol/L}$ for London EPCU) at the Screening Visit.

4.2 Exclusion Criteria

1. History of any clinically important disease or disorder.
2. History or presence of gastrointestinal, hepatic, or renal disease or any other condition known to interfere with absorption, distribution, metabolism, or excretion of drugs.
3. Any clinically important illness, medical/surgical procedure, or trauma within 4 weeks of the first administration of study intervention.
4. Any laboratory values with specific deviations in alanine transaminase (ALT), aspartate transaminase (AST), total bilirubin (TBL), estimated glomerular filtration rate, or hemoglobin at the Screening Visit or on Admission
5. Any clinically important abnormalities in clinical chemistry, hematology, or urinalysis results other than those described under exclusion criterion number 4, at Screening and/or Admission to the Clinical Unit
6. Any positive result on Screening for serum HBsAg, hepatitis B core antibody or human immunodeficiency virus.
7. History of severe allergy/hypersensitivity or ongoing clinically important allergy/hypersensitivity, as judged by the investigator or history of hypersensitivity to drugs with a similar chemical structure or class to AZD0780, ezetimibe, rosuvastatin, and bempedoic acid.
8. Treatment with any lipid-lowering therapy or AZD0780 within the 3 months prior to Screening.
9. Treatment with drugs for reduction or inhibition of Proprotein convertase subtilisin/kexin type 9 (PCSK9) within the last 12 months prior to Screening (approved or investigational and apart from AZD0780).

11. Study Identifiers & Governance

Primary ID:	869235349604422440
Registry Number:	NCT06742853
Sponsor:	AstraZeneca
Study Title:	A Study to Investigate the Pharmacokinetics, Safety, Tolerability, and Efficacy of AZD0780 With Ezetimibe Combinations in Healthy Adults With Elevated LDL-C.

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____