

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1 Clinical Study to Investigate the Safety and Pharmacokinetics of MK-5684 in China Participants With Metastatic Castration-Resistant Prostate Cancer
Short Title:	A Study of Opevesostat (MK-5684) in China Participants With Metastatic Castration-Resistant Prostate Cancer (mCRPC) (MK-5684-001)
Protocol Number:	892317133212947536
NCT Number:	NCT06136598
Sponsor Name:	Merck
Investigational Product:	dexamethasone
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants Who Experience an Adverse Event (AE): An AE is defined as any untoward medical occurrence associated with the use of a drug in a participant, whether or not considered drug related. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a medicinal product and does not imply any judgment about causality. 2. Number of Participants Who Discontinue Study Intervention Due to an AE: An AE is defined as any untoward medical occurrence associated with the use of a drug in a participant, whether or not considered drug related. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a medicinal product and does not imply any judgment about causality. 3. Maximum Plasma Concentration (Cmax) of opevesostat: Blood samples will be collected at pre-specified timepoints to determine the Cmax of opevesostat.
Secondary Obj.:	1. Prostate-specific Antigen (PSA) Response Rate 2. Radiographic Progression-free Survival (rPFS) Per Prostate Cancer Working Group (PCWG)-modified Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST 1.1) 3. Objective Response Rate (ORR) Per PCWG-modified RECIST 1.1

2.2 Background & Rationale

The primary objectives of this study are to evaluate the safety and tolerability of opevesostat in the treatment of male Chinese participants with metastatic castration-resistant prostate cancer (mCRPC) and to characterize the pharmacokinetic profile of opevesostat. There are no formal hypotheses to be tested in this study.

3. Study Design & Methodology

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, Status: ACTIVE_NOT_RECRUITING
Target N: 14

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has histologically or cytologically confirmed adenocarcinoma of the prostate without small cell histology.
- * Has prostate cancer while receiving androgen deprivation therapy (ADT), or post-bilateral orchiectomy, within 6 months before screening.
- * Has evidence of progression ≥ 4 weeks since last flutamide treatment or ≥ 6 weeks since last bicalutamide or nilutamide treatment.
- * Has evidence of metastatic disease documented by either bone lesions on bone scan and/or soft tissue shown by CT/MRI.
- * Has disease that progressed during or after treatment with at least 1 line of next-generation hormonal agents (NHAs) for hormone-sensitive prostate cancer (HSPC) or castration-resistant prostate cancer (CRPC) for at least 8 weeks (at least 14 weeks for participants with bone progression).
- * Has received at least 1 line of taxane-based chemotherapy for HSPC or CRPC and have had progressed disease during or on treatment, or refused or ineligible to receive chemotherapy.
- * Has a life expectancy of ≥ 3 months.

4.2 Exclusion Criteria

- * Has history of a second malignancy, unless potentially curative treatment has been completed with no evidence of malignancy for 3 years.
- * Has presence of gastrointestinal condition, e.g. malabsorption, that might affect the adsorption of study intervention.
- * Has a history of pituitary dysfunction.
- * Has poorly controlled diabetes mellitus.
- * Has active or unstable cardio/cerebro-vascular disease, including thromboembolic events.
- * Has undergone major surgery, including local prostate intervention (except prostate biopsy), within 4 weeks of the date of allocation.
- * Has received an anticancer monoclonal antibody (mAb) within 4 weeks of allocation, or has not recovered from adverse events (AEs) due to mAbs administered more than 4 weeks before the date of allocation.
- * Received prior systemic anticancer therapy including investigational agents within 4 weeks before the date of allocation.
- * Has a diagnosis of immunodeficiency or is receiving chronic systemic steroid therapy (in dosing exceeding 10 mg daily of prednisone equivalent) or any form of immunosuppressive therapy within 7 days prior to the start of study intervention.
- * Has a known active central nervous system (CNS) metastases and/or carcinomatous meningitis.
- * Has an active autoimmune disease that has required systemic treatment in the past 2 years.
- * Has a history of or current human immunodeficiency virus (HIV) infection.
- * Has a concurrent Hepatitis B or Hepatitis C virus infection.
- * Has a history of allogenic tissue or solid organ transplant.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

11. Study Identifiers & Governance

Primary ID:	892317133212947536
Registry Number:	NCT06136598
Sponsor:	Merck
Study Title:	A Study of Opevesostat (MK-5684) in China Participants With Metastatic Castration-Resistant Prostate Cancer (mCRPC) (MK-5684-001)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____