

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	EPIK-O: A Phase III, Multi-center, Randomized (1:1), Open-label, Active-controlled, Study to Assess the Efficacy and Safety of Alpelisib (BYL719) in Combination With Olaparib as Compared to Single Agent Cytotoxic Chemotherapy, in Participants With no Germline BRCA Mutation Detected, Platinum-resistant or Refractory, High-grade Serous Ovarian Cancer
Short Title:	Alpelisib Plus Olaparib in Platinum-resistant/Refractory, High-grade Serous Ovarian Cancer, With no Germline BRCA Mutation Detected
Protocol Number:	1051353345532287569
NCT Number:	NCT04729387
Sponsor Name:	Novartis
Investigational Product:	doxorubicin
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression Free Survival (PFS) based on Blinded Independent Review Committee (BIRC) assessment using RECIST 1.1 criteria: PFS is defined as the time from the date of randomization to the date of the first documented progression (based on RECIST 1.1 criteria) or death due to any cause. If a participant has not had an event, PFS will be censored at the date of the last adequate tumor assessment
Secondary Obj.:	1. Overall survival 2. Number of participants with dose interruptions and dose reductions 3. Dose intensity

2.2 Background & Rationale

The objective of this study is to assess the efficacy and safety of the combination of alpelisib and olaparib compared with single agent cytotoxic chemotherapy in patients with platinum resistant or refractory high-grade serous ovarian cancer, with no germline BRCA mutation detected.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	358

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

* Participant has histologically confirmed diagnosis of high-grade serous or high-grade endometrioid ovarian cancer,

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

fallopian tube cancer, or primary peritoneal cancer

- * Measurable disease, i.e., at least one measurable lesion per RECIST 1.1 criteria (a lesion at a previously irradiated site may only be counted as a target lesion if there is clear sign of progression since the irradiation)
- * If no measurable disease is present, the disease should be assessable by Gynecologic Cancer Intergroup criteria (GCIC) for CA-125
- * Participant has no germline BRCA1/2 mutation as determined by an FDA approved assay.
- * Participant has an Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1
- * Participant has platinum-resistant (progression within one to six months after completing platinum-based therapy) or platinum refractory disease (progression during treatment or within 4 weeks after the last dose), where platinum-based therapy is not an option, according to the GCIG 5th Ovarian Cancer Consensus Conference definitions. The platinum-based chemotherapy regimen does not necessarily need to be the last regimen the participant received prior to study entry.
- * Participant must have received at least one but no more than three prior systemic treatment regimens and for whom single-agent chemotherapy is appropriate as the next line of treatment.
- * Participant has adequate bone marrow and organ function

4.2 Exclusion Criteria

- * Participant has received prior treatment with any PI3K, mTOR or AKT inhibitor.
- * Participant is concurrently using other anti-cancer therapy
- * Participant is in a state of small or large bowel obstruction or has other impairment of gastrointestinal (GI) function or GI disease
- * Participant has had surgery within 14 days prior to starting study drug or has not recovered from major side effects
- * Participant has not recovered from all toxicities 5 related to prior anticancer therapies to baseline or NCI CTCAE Version 4.03 Grade ?1. Exception to this criterion: participants with any grade of alopecia are allowed to enter the study.
- * Participants with liver impairment and Child Pugh score B or C
- * Participant has received radiotherapy ? 4 weeks or limited field radiation for palliation ?2 weeks prior to randomization, and who has not recovered to baseline, grade 1 or better from related side effects of such therapy (with the exception of alopecia).
- * Participant has a known hypersensitivity to any of the study drugs or excipients

Other inclusion/exclusion criteria may apply

11. Study Identifiers & Governance

Primary ID:	1051353345532287569
Registry Number:	NCT04729387
Sponsor:	Novartis
Study Title:	Alpelisib Plus Olaparib in Platinum-resistant/Refractory, High-grade Serous Ovarian Cancer, With no Germline BRCA Mutation Detected

15. Sign-off & Approval

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____