

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A US Phase 3b, Multi-center, Randomized, Double-blind, Double-Dummy Study to Evaluate Efficacy of Remibrutinib Compared to Dupilumab at Early Timepoints in Adults With Chronic Spontaneous Urticaria Inadequately Controlled by Second Generation H1-Antihistamines
Short Title:	A Study to Evaluate Efficacy of Remibrutinib Compared to Dupilumab at Early Timepoints in Adults With Chronic Spontaneous Urticaria Inadequately Controlled by Second Generation H1-antihistamines
Protocol Number:	159497182413785272
NCT Number:	NCT06868212
Sponsor Name:	Novartis
Investigational Product:	ibrutinib
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Absolute change from baseline in Weekly Urticaria Activity Score (UAS7) at Week 4: The UAS7 is the sum of the Weekly Hives Severity (HSS7) score and the Weekly Itch Severity (ISS7) score, and ranges from 0-42. Weekly scores (HSS7 and ISS7 scores) will be derived by adding up the average daily scores of the 7 days preceding the visit.
Secondary Obj.:	1. Improvement of severity of hives, assessed as absolute change from baseline in HSS7 score at Week 4 2. Improvement of severity of itch, assessed as absolute change from baseline in ISS7 score at Week 4 3. Improvement of UAS7, assessed as absolute change from baseline in UAS7 score at Week 1

2.2 Background & Rationale

This is a US, multi-center, randomized, double-blind, double-dummy, Phase 3b study to evaluate efficacy of remibrutinib (25 mg twice daily [b.i.d.] by mouth [p.o.]) compared to dupilumab (600 mg loading dose administered subcutaneously (s.c.) followed by 300 mg every 2 weeks s.c.) at early timepoints (4 weeks and earlier), when administered as an add-on treatment to second generation H1-antihistamines (sgH1-AH) (standard label dose as background therapy) in adult US participants with moderate to severe chronic spontaneous urticaria (CSU) inadequately controlled by sgH1-AHs.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	400

4. Subject Selection (Eligibility Criteria)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.1 Inclusion Criteria

- * Adults ? 18 years of age at the time of signing the informed consent
- * CSU duration for ? 6 months prior to screening (defined as the onset of CSU determined by the Investigator based on all available supporting documentation)
- * Diagnosis of CSU inadequately controlled by sgH1-AH at the time of randomization, defined as:
- * The presence of itch and hives for ? 6 consecutive weeks prior to screening despite the use of sgH1-AH during the 7 days prior to randomization (Day 1):
- * UAS7 score (range, 0-42) ? 16, and
- * ISS7 score (range, 0-21) ? 6, and
- * HSS7 score (range, 0-21) ? 6
- * Documentation of hives within 3 months before randomization (either at screening and/or at randomization); or documented in the participants medical history
- * Willing and able to complete an Urticaria Patient Daily Diary (UPDD) for the duration of the study and adhere to the study protocol
- * Participants must not have had more than one missing UPDD entry (either morning or evening) in the 7 days prior to randomization (Day 1)

4.2 Exclusion Criteria

- * Previous use of remibrutinib or other bruton's tyrosine kinase (BTK) inhibitors
- * Previous use of dupilumab
- * Evidence of clinically significant cardiovascular (such as but not limited to myocardial infarction, unstable ischemic heart disease, NYHA Class III/IV left ventricular failure, arrhythmia and uncontrolled hypertension within 12 months prior to Visit 1), neurological, psychiatric, pulmonary, renal, hepatic (past history or current), endocrine or metabolic disorder, or immunodeficiency that, in the investigator's opinion, would compromise the safety of the participant, interfere with the interpretation of the study results or otherwise preclude participation or protocol adherence of the participant.
- * Evidence of hematological disorders (including coagulation disorders or significant bleeding risk)
- * History or evidence of gastrointestinal disease (including gastrointestinal bleeding, e.g., in association with use of nonsteroidal anti-inflammatory drugs (NSAID), that was clinically relevant (e.g., where intervention was indicated or requiring hospitalization or blood transfusion)
- * Requirement for anti-platelet medication, except for acetylsalicylic acid up to 100 mg/d or clopidogrel up to 75 mg/d. The use of dual anti-platelet therapy (e.g., acetylsalicylic acid + clopidogrel) is prohibited.
- * Requirement for anticoagulant medication (for example, warfarin or Novel Oral Anti-Coagulants \[NOAC\])
- * History or current hepatic disease including but not limited to acute or chronic hepatitis, cirrhosis or hepatic failure or hepatic parameters at screening: Aspartate Aminotransferase (AST)/ Alanine Aminotransferase (ALT) levels more than 1.5x ULN or International Normalized Ratio (INR) \> 1.5 at screening

11. Study Identifiers & Governance

Primary ID:	159497182413785272
Registry Number:	NCT06868212
Sponsor:	Novartis
Study Title:	A Study to Evaluate Efficacy of Remibrutinib Compared to Dupilumab at Early

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Timepoints in Adults With Chronic Spontaneous Urticaria Inadequately Controlled by
Second Generation H1-antihistamines

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____