

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A PHASE 3 STUDY TO EVALUATE THE EFFICACY, IMMUNOGENICITY, AND SAFETY OF RESPIRATORY SYNCYTIAL VIRUS (RSV) PREFUSION F SUBUNIT VACCINE IN ADULTS
Short Title:	Study to Evaluate the Efficacy, Immunogenicity, and Safety of RSVpreF in Adults.
Protocol Number:	204114652433121858
NCT Number:	NCT05035212
Sponsor Name:	Pfizer
Investigational Product:	RSVPreF3/AS01E RSV Vaccine
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Efficacy Study: Number of first episode of RSV-associated lower respiratory tract illness (LRTI-RSV) in the first RSV season: Reverse Transcriptase-Polymerase Chain Reaction (RT-PCR)-confirmed RSV A and/or RSV B- associated acute respiratory illness (ARI) is assessed. LRTI-RSV is defined as an ARI with 2 or more of the lower respiratory signs/symptoms lasting more than 1 day during the same illness, plus RT-PCR-confirmed RSV infection within 7 days of ARI symptom onset. Reverse Transcriptase-Polymerase Chain Reaction (RT-PCR)-confirmed RSV A and/or RSV B- associated acute respiratory illness (ARI) is assessed. LRTI-RSV is defined as an ARI with 3 or more of the lower respiratory signs/symptoms lasting more than 1 day during the same illness, plus RT-PCR-confirmed RSV infection within 7 days of ARI symptom onset. 2. Efficacy Study: Proportion of participants reporting prompted local reactions within 7-days after vaccination: Local reactions included pain at injection site, redness and swelling recorded by participants in an e-diary. Redness and swelling were measured and recorded in measuring device units. 1 measuring device unit =0.5 cm and graded as mild: 2.5 to 5.0 cm, moderate: > 5.0 to 10.0 cm and severe: >10 cm. Pain at injection site was graded as mild: did not interfere with daily activity, moderate: interfere with daily activity and severe: prevented daily activity 3. Efficacy Study: Proportion of participants reporting prompted systemic events within 7-days after vaccination: Systemic reactions:fever, fatigue/tiredness, headache, nausea, muscle pain, joint pain, vomiting, diarrhea and any systemic event recorded by participants in an e-diary. Fever: greater than equal to ($\geq$)38.0 degrees (deg) Celsius (C), mild ($\geq$38.0 to 38.4 deg C, >38.4 to 38.9 deg C), moderate ($\geq$38.9 to 40.0 deg C and >40.0 deg C), severe ($\geq$38.9 deg C to 40.0 deg C) and grade 4 ($\geq$40.0 deg C). Fatigue, headache, nausea,
--------------------	--

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

muscle pain and joint pain were graded as mild: did not interfere with activity, moderate: some interference with activity and severe: prevented daily routine activity. Vomiting was graded as mild: 1 to 2 times in 24 hours(h), moderate: \>2 times in 24h and severe: requires intravenous hydration. Diarrhea was graded as mild: 2 to 3 loose stools in 24h, moderate: 4 to 5 loose stools in 24h and severe: 6 or more loose stools in 24h.

Secondary Obj.:

1. Efficacy Study: Number of first episode of RSV-associated severe LRTI (sLRTI-RSV) in the first RSV season 2. Efficacy Study: Number of first episode of LRTI-RSV in the second RSV season 3. Efficacy Study: Number of first episode of LRTI-RSV across 2 RSV seasons

2.2 Background & Rationale

Efficacy Study: This randomized, double-blinded, placebo-controlled Phase 3 study is designed to assess the safety, immunogenicity, and efficacy of a single dose of RSVpreF in the prevention of LRTI-RSV in adults:

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N: 38861

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Participants who are willing and able to comply with all scheduled visits, vaccination plan, laboratory tests, lifestyle considerations, frequent symptom assessment by mobile device application, and other study procedures, including collection of nasal swabs by themselves and by study staff when indicated.
- * Healthy participants who are determined by medical history, physical examination (if required), and clinical judgment of the investigator to be eligible for inclusion in the study.

Note: Healthy participants with preexisting stable disease, defined as disease not requiring significant change in therapy or hospitalization for worsening disease during the 6 weeks before enrollment, can be included. Specific criteria for participants with known stable infection with HIV, HCV, or HBV can be found in the protocol.

- * Adults who are ambulatory and live in the community, or in assisted living or long-term care residential facilities that provide minimal assistance, such that the participant is primarily responsible for self-care and activities of daily living.
- * Capable of giving signed informed consent as described in the protocol, which includes compliance with the requirements and restrictions listed in the ICD and in this protocol.
- * Male or female participants ?60 years of age.
- * Male participants able to father children must agree to use a highly effective method of contraception from the time of informed consent through at least 28 days after study intervention administration.
- * Female participants must not be of childbearing potential.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.2 Exclusion Criteria

- * Bleeding diathesis or condition associated with prolonged bleeding that would, in the opinion of the investigator, contraindicate intramuscular injection.
- * History of severe adverse reaction associated with a vaccine and/or severe allergic reaction (eg, anaphylaxis) to any component of the study intervention(s) or any related vaccine.
- * Serious chronic disorder including metastatic malignancy, end-stage renal disease with or without dialysis, clinically unstable cardiac disease, or any other disorder that, in the investigator's opinion, excludes the participant from participating in the study.
- * Immunocompromised individuals with known or suspected immunodeficiency, as determined by history and/or laboratory/physical examination.
- * Other medical or psychiatric condition including recent (within the past year) or active suicidal ideation/behavior or laboratory abnormality that may increase the risk of study participation or, in the investigator's judgment, make the participant inappropriate for the study.
- * Participation in other studies involving an investigational product within 28 days prior to consent and/or through and including the 6-months follow-up visit (Visit 3).

Note: This criterion does not apply to participants who are participating in a follow-up period for another study involving a study intervention that is an investigational drug or vaccine, if receipt of the last dose was at least 6 months prior to consenting for this study and there is no further dosing anticipated from the previous study during the participant's participation in this study - Individuals who receive chronic systemic treatment with immunosuppressive therapy, including cytotoxic agents, monoclonal antibodies, systemic corticosteroids, or radiotherapy, eg, for cancer or an autoimmune disease, from 60 days before study intervention administration or planned receipt throughout the study. If systemic corticosteroids have been administered short term (<14 days) for treatment of an acute illness, participants should not be enrolled in the study until corticosteroid therapy has been discontinued for at least 28 days before study intervention administration. Inhaled/nebulized, intra articular, intrabursal, or topical (skin or eyes) corticosteroids are permitted.

Note: Participants with COPD or asthma can be enrolled if chronic corticosteroids do not exceed a dose equivalent to 10 mg/day of prednisone.

- * Receipt of blood/plasma products or immunoglobulin within 60 days before study intervention administration.
- * Previous vaccination with any licensed or investigational RSV vaccine or planned receipt during study participation.
- * Investigator site staff or Pfizer employees directly involved in the conduct of the study, site staff otherwise supervised by the investigator, and their respective family members.

Substudy A

Inclusion Criteria:

- * Participants who are willing and able to comply with all scheduled visits, vaccination plan, laboratory tests, lifestyle

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

considerations, frequent symptom assessment by mobile device application, and other study procedures, including collection of nasal swabs by themselves and by study staff when indicated.

* Healthy participants who are determined by medical history, physical examination (if required), and clinical judgment of the investigator to be eligible for inclusion in the study.

Note: Healthy participants with preexisting stable disease, defined as disease not requiring significant change in therapy or hospitalization for worsening disease during the 6 weeks before enrollment, can be included. Specific criteria for participants with known stable infection with HIV, HCV, or HBV can be found in the protocol.

* Adults who are ambulatory and live in the community, or in assisted living or long-term care residential facilities that provide minimal assistance, such that the participant is primarily responsible for self-care and activities of daily living.

* Capable of giving signed informed consent as described in the protocol, which includes compliance with the requirements and restrictions listed in the ICD and in this protocol.

* Male or female participants ≥60 years of age.

* Male participants able to father children must agree to use a highly effective method of contraception from the time of informed consent through at least 28 days after study intervention administration.

* Female participants must not be of childbearing potential.

* Participants who received RSVpreF in the efficacy study.

* Participants who have a Visit 2 serology sample available for testing from the efficacy study, completed the end-of-Season 2 visit (Visit 5), and did not meet exclusion criteria throughout the efficacy study duration.

Exclusion Criteria:

* Bleeding diathesis or condition associated with prolonged bleeding that would, in the opinion of the investigator, contraindicate intramuscular injection.

* History of severe adverse reaction associated with a vaccine and/or severe allergic reaction (eg, anaphylaxis) to any component of the study intervention(s) or any related vaccine.

* Serious chronic disorder including metastatic malignancy, end-stage renal disease with or without dialysis, clinically unstable cardiac disease, or any other disorder that, in the investigator's opinion, excludes the participant from participating in the study.

* Immunocompromised individuals with known or suspected immunodeficiency, as determined by history and/or laboratory/physical examination.

* Other medical or psychiatric condition including recent (within the past year) or active suicidal ideation/behavior or laboratory abnormality that may increase the risk of study participation or, in the investigator's judgment, make the participant inappropriate for the study.

* Participation in other studies involving an investigational product within 28 days prior to consent and/or through and including the 6-months follow-up visit (Visit 103).

Note: This criterion does not apply to participants who are participating in a follow-up period for another study involving a

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

study intervention that is an investigational drug or vaccine, if receipt of the last dose was at least 6 months prior to consenting for this study and there is no further dosing anticipated from the previous study during the participant's participation in this study

↳ Individuals who receive chronic systemic treatment with immunosuppressive therapy, including cytotoxic agents, monoclonal antibodies, systemic corticosteroids, or radiotherapy, eg, for cancer or an autoimmune disease, from 60 days before study intervention administration or planned receipt throughout the study. If systemic corticosteroids have been administered short term (<14 days) for treatment of an acute illness, participants should not be enrolled in the study until corticosteroid therapy has been discontinued for at least 28 days before study intervention administration. Inhaled/nebulized, intra articular, intrabursal, or topical (skin or eyes) corticosteroids are permitted.

Note: Participants with COPD or asthma can be enrolled if chronic corticosteroids do not exceed a dose equivalent to 10 mg/day of prednisone.

- * Receipt of blood/plasma products or immunoglobulin within 60 days before study intervention administration.
- * Previous vaccination with any licensed or investigational RSV vaccine or planned receipt during study participation.

Note: This criterion does not include the receipt of RSVpreF in the efficacy study. Per inclusion criterion #6, receipt of RSVpreF in the efficacy study is required to participate in Substudy A.

- * Investigator site staff or Pfizer employees directly involved in the conduct of the study, site staff otherwise supervised by the investigator, and their respective family members.
- * Participant was confirmed by the sponsor to have previously received the study intervention more than once.

Substudy B

Inclusion Criteria:

- * Participants who are willing and able to comply with all scheduled visits, vaccination plan, laboratory tests, lifestyle considerations, frequent symptom assessment by mobile device application, and other study procedures, including collection of nasal swabs by themselves and by study staff when indicated.
- * Healthy participants who are determined by medical history, physical examination (if required), and clinical judgment of the investigator to be eligible for inclusion in the study.

Note: Healthy participants with preexisting stable disease, defined as disease not requiring significant change in therapy or hospitalization for worsening disease during the 6 weeks before enrollment, can be included. Specific criteria for participants with known stable infection with HIV, HCV, or HBV can be found in the protocol.

- * Adults who are ambulatory and live in the community, or in assisted living or long-term care residential facilities that provide minimal assistance, such that the participant is primarily responsible for self-care and activities of daily living.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Capable of giving signed informed consent as described in the protocol, which includes compliance with the requirements and restrictions listed in the ICD and in this protocol.
- * Male or female participants ≥60 years of age.

* Male participants able to father children must agree to use a highly effective method of contraception from the time of informed consent through at least 28 days after study intervention administration.

- * Female participants must not be of childbearing potential. 6. Participants who received RSVpreF in the efficacy study.
- * Participants who received RSVpreF in the efficacy study.
- * Participants who have a Visit 2 serology sample available for testing from the efficacy study and did not meet exclusion criteria through Visit 4 of the efficacy study.

Exclusion Criteria:

- * Bleeding diathesis or condition associated with prolonged bleeding that would, in the opinion of the investigator, contraindicate intramuscular injection.
- * History of severe adverse reaction associated with a vaccine and/or severe allergic reaction (eg, anaphylaxis) to any component of the study intervention(s) or any related vaccine.
- * Serious chronic disorder including metastatic malignancy, end-stage renal disease with or without dialysis, clinically unstable cardiac disease, or any other disorder that, in the investigator's opinion, excludes the participant from participating in the study.
- * Immunocompromised individuals with known or suspected immunodeficiency, as determined by history and/or laboratory/physical examination.
- * Other medical or psychiatric condition including recent (within the past year) or active suicidal ideation/behavior or laboratory abnormality that may increase the risk of study participation or, in the investigator's judgment, make the participant inappropriate for the study.
- * Participation in other studies involving an investigational product within 28 days prior to consent and/or through and including the 6-months follow-up visit (Visit 203).

Note: This criterion does not apply to participants who are participating in a follow-up period for another study involving a study intervention that is an investigational drug or vaccine, if receipt of the last dose was at least 6 months prior to consenting for this study and there is no further dosing anticipated from the previous study during the participant's participation in this study - Individuals who receive chronic systemic treatment with immunosuppressive therapy, including cytotoxic agents, monoclonal antibodies, systemic corticosteroids, or radiotherapy, eg, for cancer or an autoimmune disease, from 60 days before study intervention administration or planned receipt throughout the study. If systemic corticosteroids have been administered short term (<14 days) for treatment of an acute illness, participants should not be enrolled in the study until corticosteroid therapy has been discontinued for at least 28 days before study intervention administration. Inhaled/nebulized, intra articular, intrabursal, or topical (skin or eyes) corticosteroids are permitted.

Note: Participants with COPD or asthma can be enrolled if chronic corticosteroids do not exceed a dose equivalent to 10

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

mg/day of prednisone.

- * Receipt of blood/plasma products or immunoglobulin within 60 days before study intervention administration.
- * Previous vaccination with any licensed or investigational RSV vaccine or planned receipt during study participation.

Note: This criterion does not include the receipt of RSVpreF in the efficacy study. Per inclusion criterion #6, receipt of RSVpreF in the efficacy study is required to participate in Substudy B.

- * Investigator site staff or Pfizer employees directly involved in the conduct of the study, site staff otherwise supervised by the investigator, and their respective family members.
- * Participant was confirmed by the sponsor to have previously received the study intervention more than once.
- * Participants who completed Vaccination 1 from the efficacy study less than 9 months or greater than 15 months prior to revaccination.

Substudy C

Inclusion Criteria:

- * Participants who are willing and able to comply with all scheduled visits, vaccination plan, laboratory tests, lifestyle considerations, frequent symptom assessment by mobile device application, and other study procedures, when indicated.
- * Healthy participants who are determined by medical history, physical examination (if required), and clinical judgment of the investigator to be eligible for inclusion in the study.

Note: Healthy participants with preexisting stable disease, defined as disease not requiring significant change in therapy or hospitalization for worsening disease during the 6 weeks before enrollment, can be included. Specific criteria for participants with known stable infection with HIV, HCV, or HBV can be found in the protocol.

- * Adults who are ambulatory and live in the community, or in assisted living or long-term care residential facilities that provide minimal assistance, such that the participant is primarily responsible for self-care and activities of daily living.
- * Capable of giving signed informed consent as described in the protocol, which includes compliance with the requirements and restrictions listed in the ICD and in this protocol.
- * Male or female participants ≥60 years of age.
- * Participants who received RSVpreF in the efficacy study.
- * Participants who have a Visit 2 serology sample available for testing from the efficacy study, completed the end-of-Season 2 visit (Visit 5), and did not meet exclusion criteria throughout the efficacy study duration.

Exclusion Criteria:

- * Bleeding diathesis or condition associated with prolonged bleeding that would, in the opinion of the investigator,

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

contraindicate intramuscular injection.

- * History of severe adverse reaction associated with a vaccine and/or severe allergic reaction (eg, anaphylaxis) to any component of the study intervention(s) or any related vaccine.
- * Serious chronic disorder including metastatic malignancy, end-stage renal disease with or without dialysis, clinically unstable cardiac disease, or any other disorder that, in the investigator's opinion, excludes the participant from participating in the study.
- * Immunocompromised individuals with known or suspected immunodeficiency, as determined by history and/or laboratory/physical examination.
- * Other medical or psychiatric condition including recent (within the past year) or active suicidal ideation/behavior or laboratory abnormality that may increase the risk of study participation or, in the investigator's judgment, make the participant inappropriate for the study.
- * Participation in other studies involving an investigational product within 28 days prior to consent and/or through and including the 6-months follow-up visit (Visit 303 or Visit 306).

Note: This criterion does not apply to participants who are participating in a follow-up period for another study involving a study intervention that is an investigational drug or vaccine, if receipt of the last dose was at least 6 months prior to consenting for this study and there is no further dosing anticipated from the previous study during the participant's participation in this study

↳ Individuals who receive chronic systemic treatment with immunosuppressive therapy, including cytotoxic agents, monoclonal antibodies, systemic corticosteroids, or radiotherapy, eg, for cancer or an autoimmune disease, from 60 days before study intervention administration or planned receipt throughout the study. If systemic corticosteroids have been administered short term (<14 days) for treatment of an acute illness, participants should not be enrolled in the study until corticosteroid therapy has been discontinued for at least 28 days before study intervention administration. Inhaled/nebulized, intra articular, intrabursal, or topical (skin or eyes) corticosteroids are permitted.

Note: Participants with COPD or asthma can be enrolled if chronic corticosteroids do not exceed a dose equivalent to 10 mg/day of prednisone.

- * Receipt of blood/plasma products or immunoglobulin within 60 days before study intervention administration.
- * Previous vaccination with any licensed or investigational RSV vaccine or planned receipt during study participation.

Note: This criterion does not include the receipt of RSVpreF in the efficacy study. Per inclusion criterion #6, receipt of RSVpreF in the efficacy study is required to participate in Substudy C.

- * Investigator site staff or Pfizer employees directly involved in the conduct of the study, site staff otherwise supervised by the investigator, and their respective family members.
- * Participant was confirmed by the sponsor to have previously received the study intervention more than once.
- * Prior history of any subtype of Guillain-Barré syndrome of any etiology.
- * Current or prior participation in Substudy A or Substudy B.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* Participants who completed Vaccination 1 from the efficacy study less than 32 months or greater than 40 months prior to revaccination.

11. Study Identifiers & Governance

Primary ID: 204114652433121858
Registry Number: NCT05035212
Sponsor: Pfizer
Study Title: Study to Evaluate the Efficacy, Immunogenicity, and Safety of RSVpreF in Adults.

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____