

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	PROSPER: A MULTINATIONAL, PHASE 3, RANDOMIZED, DOUBLE-BLIND, PLACEBO-CONTROLLED, EFFICACY AND SAFETY STUDY OF ENZALUTAMIDE IN PATIENTS WITH NONMETASTATIC CASTRATION-RESISTANT PROSTATE CANCER
Short Title:	Safety and Efficacy Study of Enzalutamide in Patients With Nonmetastatic Castration-Resistant Prostate Cancer
Protocol Number:	217472311191024771
NCT Number:	NCT02003924
Sponsor Name:	Pfizer
Investigational Product:	enzalutamide
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Metastasis Free Survival (MFS): MFS:time from randomization to first date of radiographic progression (RP) (by Blinded independent central radiology review \[BICR\]) at any time or death within 112 days of treatment discontinuation without evidence of RP.RP for bone disease:appearance of 1 or more metastatic lesions on bone scan.RP for soft tissue disease:per Response Evaluation Criteria in Solid Tumors,\[RECIST 1.1\])-at least a 20 percent (%) increase in the sum of diameters of target lesions,taking as reference the smallest sum on study (includes the baseline sum if smallest on study).Participants who did not have MFS event at the time of analysis data cut-off (28 June 2017) were censored at date of last assessment showing no objective evidence of RP prior to skeletal-related event or two or more consecutive missed tumor assessments. Participants who were randomized but later confirmed to have metastatic disease before randomization were censored on date of randomization. Analysis was based on Kaplan-Meier estimates.
Secondary Obj.:	1. Time to Prostate-Specific Antigen (PSA) Progression 2. Time to First Use of New Antineoplastic Therapy 3. Overall Survival

2.2 Background & Rationale

The purpose of this study is to assess the safety and efficacy of enzalutamide in patients with non metastatic prostate cancer.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
-------------------	--

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Target N: 1401

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Histologically or cytologically confirmed adenocarcinoma of the prostate without neuroendocrine differentiation, signet cell, or small cell features;
- * Ongoing androgen deprivation therapy with a gonadotropin-releasing hormone (GnRH) agonist/antagonist or prior bilateral orchiectomy (medical or surgical castration);
- * Testosterone ≥ 50 ng/dL (≥ 1.73 nmol/L) at screening;
- * Progressive disease on androgen deprivation therapy at enrollment;
- * PSA and the screening PSA assessed by the central laboratory (central PSA) should be ≥ 2 μ g/L (2 ng/mL);
- * PSA doubling time ≥ 10 months;
- * No prior or present evidence of metastatic disease;
- * Asymptomatic prostate cancer;
- * Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1;
- * Estimated life expectancy ≥ 12 months.

4.2 Exclusion Criteria

- * Prior cytotoxic chemotherapy;
- * Use of hormonal therapy or biologic therapy for prostate cancer (other than approved bone targeting agents and GnRH agonist/antagonist therapy) or use of an investigational agent within 4 weeks of randomization;
- * Known or suspected brain metastasis or active leptomeningeal disease;
- * History of another invasive cancer within 3 years of randomization;
- * Absolute neutrophil count $< 1000/\mu$ L, platelet count $< 100,000/\mu$ L, or hemoglobin < 10 g/dL (6.2 mmol/L) at screening;
- * Total bilirubin ≥ 1.5 times the upper limit of normal;
- * Creatinine > 2 mg/dL (177 μ mol/L) at screening;
- * Albumin < 3.0 g/dL (30 g/L) at screening;
- * History of seizure or any condition that may predispose to seizure;
- * Clinically significant cardiovascular disease;
- * Gastrointestinal disorder affecting absorption;
- * Major surgery within 4 weeks of randomization;
- * Hypersensitivity reaction to the active pharmaceutical ingredient or any of the capsule components, including Labrasol, butylated hydroxyanisole, and butylated hydroxytoluene;
- * Any concurrent disease, infection, or comorbid condition that interferes with the ability of the patient to participate in the trial, which places the patient at undue risk, or complicates the interpretation of data, in the opinion of the investigator or medical monitor.

11. Study Identifiers & Governance

Primary ID: 217472311191024771

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Registry Number: NCT02003924

Sponsor: Pfizer

Study Title: Safety and Efficacy Study of Enzalutamide in Patients With Nonmetastatic Castration-Resistant Prostate Cancer

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____