

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Multicenter, Randomized, Double-Masked, Active Comparator-Controlled Study to Evaluate the Efficacy and Safety of Faricimab in Patients With Choroidal Neovascularization Secondary to Pathologic Myopia
Short Title:	A Study to Evaluate the Efficacy and Safety of Faricimab in Patients With Choroidal Neovascularization Secondary to Pathologic Myopia
Protocol Number:	219075524383025253
NCT Number:	NCT06176352
Sponsor Name:	Roche
Investigational Product:	ranibizumab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Change from Baseline in Best-Corrected Visual Acuity (BCVA) Averaged Over Weeks 4, 8, and 12
Secondary Obj.:	1. Change from Baseline in BCVA Over Time 2. Percentage of Participants Gaining ≥15 Letters in BCVA from Baseline Averaged Over Weeks 4, 8, and 12 3. Percentage of Participants Gaining ≥15 Letters in BCVA from Baseline Over Time

2.2 Background & Rationale

This is a Phase III, multicenter, randomized, double-masked, active comparator-controlled study evaluating the efficacy and safety of faricimab in patients with myopic choroidal neovascularization (CNV). This non-inferiority study will compare 6.0 mg faricimab versus 0.5 mg ranibizumab administered at a pro-re-nata (PRN) dosing regimen after an initial active IVT treatment administration at randomization (Day 1).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	280

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- Treatment-naïve choroidal neovascularization (CNV) secondary to myopia
- Diagnosis of active myopic CNV in the study eye:
 - Presence of high myopia, worse than -6 diopters of spherical equivalence

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

2. Antero-posterior elongation measurement greater than or equal to 26.0 mm
3. Presence of posterior changes compatible with pathologic myopia (e.g., tessellated fundus, lacquer cracks, etc.)
4. Presence of active leakage from CNV on FFA (determined by Central Reading Centre \[CRC\])
5. Presence of intraretinal or subretinal fluid or increase of CST on OCT (determined by CRC)
3. BCVA of 78 to 24 letters, inclusive (20/32 to 20/320 approximate Snellen equivalent), using the Early Treatment Diabetic Retinopathy Study (ETDRS) protocol on Day 1
4. Overtly healthy as determined by medical evaluation that includes medical history, physical examination, and laboratory tests
5. Ability to comply with the study protocol, in the Investigator's judgment
6. Other protocol-defined inclusion criteria apply

4.2 Exclusion Criteria

1. Any major illness or major surgical procedure within 1 month before screening
2. Pregnancy or breastfeeding, or intention to become pregnant during the study or within 3 months after the final study treatment administration
3. Uncontrolled blood pressure (systolic ≥ 180 millimetres of mercury \[mmHg\], diastolic ≥ 100 mmHg)
4. Stroke (cerebral vascular accident) or myocardial infarction within 6 months prior to Day 1
5. History of systemic or ocular disease that would contraindicate treatment with the investigational drug or comparator
6. Uncontrolled glaucoma in study eye
7. Any prior or concomitant treatment for CNV or vitreomacular-interface abnormalities, including, but not restricted to, intravitreal, periocular or laser interventions in study eye
8. Prior or concomitant periocular or intravitreal pharmacological treatment, including anti-VEGF medication, for other retinal diseases (e.g. geography atrophy, nAMD, DME etc.) in study eye
9. Other protocol-defined exclusion criteria apply

11. Study Identifiers & Governance

Primary ID:	219075524383025253
Registry Number:	NCT06176352
Sponsor:	Roche
Study Title:	A Study to Evaluate the Efficacy and Safety of Faricimab in Patients With Choroidal Neovascularization Secondary to Pathologic Myopia

15. Sign-off & Approval

Principal Investigator: _____	Date: _____
Clinical Operations Lead: _____	Date: _____
Lead Statistician: _____	Date: _____