

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 3, Randomized, Double-Blind, Placebo-Controlled, 3-Arm, 3-Period Study to Assess the Efficacy and Safety of a New Formulation of Oral Cladribine Compared With Placebo in Participants With Generalized Myasthenia Gravis (MyClad)
Short Title:	Efficacy and Safety of a New Formulation of Oral Cladribine Compared With Placebo in Participants With Generalized Myasthenia Gravis (MyClad)
Protocol Number:	276832736207126425
NCT Number:	NCT06463587
Sponsor Name:	Merck
Investigational Product:	cladribine
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Change from Baseline in Myasthenia Gravis - Activities of Daily Living (MG-ADL) Scale Score at Week 24 During the Double-Blind Placebo Controlled (DBPC) Period
Secondary Obj.:	1. Change from Baseline in Quantitative Myasthenia Gravis (QMG) Scale Score at Week 24 During the Double-Blind Placebo Controlled (DBPC) Period 2. Percentage of MG-ADL Responders at Week 24 During the Double-Blind Placebo Controlled (DBPC) Period 3. Change from Baseline in Myasthenia Gravis Composite (MGC) Scale Score at Week 24 During the Double-Blind Placebo Controlled (DBPC) Period

2.2 Background & Rationale

The purpose of this clinical study is to determine the efficacy and safety of a new oral cladribine formulation in participants with Generalized Myasthenia Gravis (gMG) in comparison to placebo. It will also investigate the sustained efficacy, the need for retreatment, and the long-term safety of oral cladribine in gMG. An additional component is included to characterize the Pharmacokinetics (PK) of the new cladribine formulation in gMG participants. This study is divided into 3 periods: the double-blind placebo control (DBPC) pivotal period, and 2 extensions, the blinded extension (BE) and the retreatment (RT) period.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	264

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Adults of ? 18 years of age at the time of signing the informed consent.
- * Diagnosis of Myasthenia Gravis with generalized muscle weakness, meeting clinical criteria for Myasthenia Gravis Foundation of America Class II to IVa classification.
- * In participants positive for Acetylcholine receptor antibody (anti-AChR) or muscle-specific kinase antibody(anti-MuSK)
- * In participants that are autoantibody seronegative and participants who are positive for anti-low-density lipoprotein receptor-related protein 4 antibodies (anti-LRP4)
- * Has a Screening and Baseline MG-ADL score more than or equal to ($\geq$) 6 with $\geq$ 50 percentage (%) of the total score due to non-ocular symptoms. Screening and Baseline MG-ADL scores must be stable. The difference between the Screening and Baseline scores should not be more than 2 and there should be no reported MG exacerbation during the Screening period
- * If treated with oral corticosteroids: should be on a stable daily dose for at least 3 months prior to and during screening. In such case, the daily dose of oral steroids should not exceed 20 milligrams(mg)/day for prednisone/ prednisolone or 16 mg/day for methylprednisolone
- * If treated with acetylcholinesterase inhibitor should be on a stable daily dose (pyridostigmine dose ? 480 mg/day) for at least 3 months prior to and during screening
- * Have a body weight $\geq$ 40 kilograms
- * Other protocol defined inclusion criteria could apply

4.2 Exclusion Criteria

- * Immunologic disorder other than MG or any other condition requiring chronic oral, intravenous, intramuscular, or intraarticular corticosteroid therapy. Well-controlled thyroid disease, as per the Treating Investigator or the participants regular treating physician recorded in the source documents, is not exclusionary
- * Molecularly characterized or suspected congenital myasthenic syndrome, Lambert-Eaton myasthenic syndrome, inherited myopathy, muscular dystrophy, acquired myopathy or any other neurologic or systematic disease that mimics MG muscular weakness
- * Active, clinically significant viral, bacterial, or fungal infection, including brain MRI findings consistent with signs of infection such as PML, or any major episode of infection requiring hospitalization or treatment with parenteral anti-infectives within 8 weeks prior or during Screening, or completion of oral anti-infectives within 8 weeks prior or during Screening. Vaginal candidiasis, onychomycosis, and genital or oral herpes simplex virus considered by the Investigator to be sufficiently controlled would not be exclusionary
- * Has a history of or current diagnosis of active tuberculosis (TB)
- * Active malignancy, or history of cancer
- * Treatment with nonsteroidal immunosuppressants, used in gMG, such as azathioprine, mycophenolate mofetil, methotrexate, cyclosporine, tacrolimus within 4 weeks prior to randomization
- * Treatment with eculizumab, rozanolixizumab efgartigimod, ravulizumab, or zilucoplan within 8 weeks prior to randomization
- * History of thymectomy within 6 months prior to Screening.
- * History of generalized seizures (except for history of infantile febrile seizures)
- * Negative for Varicella Zoster Virus antibodies at screening
- * History of myasthenic crisis in the last 12 months prior to and during screening

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * History of recurrent infections (that is 3 or more infections per year) within the last 2 years
- * Discontinuation of treatment with any non-steroidal immunosuppressants used in gMG, such as azathioprine, mycophenolate mofetil, methotrexate, cyclosporine, tacrolimus within the last 6 months prior to Screening
- * If treated with non-steroidal immunosuppressants for gMG, the dose at Screening should not exceed 50 mg/day for azathioprine, 500 mg/day for mycophenolate mofetil, 1 mg/day for tacrolimus, 50 mg/day for cyclosporine, or 7.5 mg/week for methotrexate
- * Participation in clinical study of any investigational drug within 6 months, or 5 half-lives of the investigational drug used in the previous clinical study prior to randomization, whichever is longer. However, participants with any prior exposure to cladribine may not enter the study regardless of timing of exposure
- * Other protocol defined exclusion criteria could apply

11. Study Identifiers & Governance

Primary ID: 276832736207126425

Registry Number: NCT06463587

Sponsor: Merck

Study Title: Efficacy and Safety of a New Formulation of Oral Cladribine Compared With Placebo in Participants With Generalized Myasthenia Gravis (MyClad)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____