

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	Two Part (Double-blind Inclisiran Versus Placebo [Year 1] Followed by Open-label Inclisiran [Year 2]) Randomized Multicenter Study to Evaluate Safety, Tolerability, and Efficacy of Inclisiran in Adolescents (12 to Less Than 18 Years) With Heterozygous Familial Hypercholesterolemia and Elevated LDL-cholesterol (ORION-16)
Short Title:	Study to Evaluate Efficacy and Safety of Inclisiran in Adolescents With Heterozygous Familial Hypercholesterolemia
Protocol Number:	276960662917842551
NCT Number:	NCT04652726
Sponsor Name:	Novartis
Investigational Product:	inclisiran
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage Change in LDL-C From Baseline to Day 330 (Part 1/Year 1): Percentage change in low-density lipoprotein cholesterol (LDL-C) from baseline to Day 330 (Year 1)
Secondary Obj.:	1. Time-adjusted Percent Change in LDL-C From Baseline After Day 90 and up to Day 330 (Part 1/Year 1) 2. Absolute Change in LDL-C From Baseline to up Day 330 (Part 1/Year 1) 3. Percent Change in Apo B From Baseline up to Day 330 (Part 1/Year 1)

2.2 Background & Rationale

This was a pivotal phase III study designed to evaluate safety, tolerability, and efficacy of inclisiran in adolescents with heterozygous familial hypercholesterolemia (HeFH) and elevated low density lipoprotein cholesterol (LDL-C).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: COMPLETED
Target N:	141

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Heterozygous Familial Hypercholesterolemia (HeFH) diagnosed either by genetic testing or on phenotypic criteria
- * Fasting LDL-C ≥ 130 mg/dL (3.4 mmol/L) at screening
- * Fasting triglycerides < 400 mg/dL (4.5 mmol/L) at screening
- * On maximally tolerated dose of statin (investigator's discretion) with or without other lipid-lowering therapy; stable for ? 30 days before screening

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.2 Exclusion Criteria

- * Homozygous familial hypercholesterolemia (HoFH)
- * Active liver disease
- * Secondary hypercholesterolemia, e.g. hypothyroidism or nephrotic syndrome
- * Previous treatment with monoclonal antibodies directed towards PCSK9 (within 90 days of screening)
- * Recent and/or planned use of other investigational medicinal products or devices

11. Study Identifiers & Governance

Primary ID:	276960662917842551
Registry Number:	NCT04652726
Sponsor:	Novartis
Study Title:	Study to Evaluate Efficacy and Safety of Inclisiran in Adolescents With Heterozygous Familial Hypercholesterolemia

15. Sign-off & Approval

Principal Investigator:	_____	Date:	_____
Clinical Operations Lead:	_____	Date:	_____
Lead Statistician:	_____	Date:	_____