

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Prospective, Open-Label, Multicenter Randomized Phase III Trial to Compare The Efficacy and Safety of A Combined Regimen of Obinutuzumab and Venetoclax Versus Obinutuzumab and Chlorambucil in Previously Untreated Patients With CLL and Coexisting Medical Conditions
Short Title:	Comparison of the Treatments of Obinutuzumab + Venetoclax Versus Obinutuzumab + Chlorambucil in Patients With Chronic Lymphocytic Leukemia
Protocol Number:	551814129434349837
NCT Number:	NCT02242942
Sponsor Name:	Roche
Investigational Product:	chlorambucil
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression Free Survival (PFS) Based on Investigator Assessment According to IWCLL Criteria: PFS was determined according to IWCLL 2008 criteria and defined as the time from randomization to the first occurrence of PD or death from any cause. Disease progression was characterized by at least one of the following: 1) $\geq 50\%$ increase in the absolute number of circulating lymphocytes to at least $5 \times 10^9/L$, 2) Appearance of new palpable lymph nodes (> 15 mm in longest diameter) or any new extra-nodal lesion; 3) $\geq 50\%$ increase in the longest diameter of any previous site of lymphadenopathy; 4) $\geq 50\%$ increase in the enlargement of the liver and/or spleen; 5) Transformation to a more aggressive histology.
Secondary Obj.:	1. Progression Free Survival (PFS) Based on Institutional Review Committee (IRC)-Assessments According to International Workshop on Chronic Lymphocytic Leukemia (IWCLL) Criteria 2. Percentage of Participants With an Overall Response (OR) at Completion of Treatment, as Determined by the Investigator According to IWCLL Criteria 3. Percentage of Participants With a Complete Response Rate (CRR) at the Completion of Treatment Assessment as Determined by the Investigator According to IWCLL Criteria

2.2 Background & Rationale

This open-label, multicenter, randomized Phase III study is designed to compare the efficacy and safety of a combined regimen of obinutuzumab and venetoclax versus obinutuzumab + chlorambucil in participants with chronic lymphocytic leukemia (CLL) and coexisting medical conditions. The time on study treatment was approximately one year and the follow-up period will be up to 9 years

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE3, Status: COMPLETED

Target N: 445

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Documented previously untreated CLL according to the International Workshop on Chronic Lymphocytic Leukemia (IWCLL) criteria
- * CLL requiring treatment according to IWCLL criteria
- * Total Cumulative Illness Rating Scale (CIRS score) greater than ($\geq$) 6
- * Adequate marrow function independent of growth factor or transfusion support within 2 weeks of screening as per protocol, unless cytopenia is due to marrow involvement of CLL
- * Adequate liver function
- * Life expectancy $\geq$ 6 months
- * Agreement to use highly effective contraceptive methods per protocol

4.2 Exclusion Criteria

- * Transformation of CLL to aggressive Non-Hodgkin's lymphoma (Richter's transformation or pro-lymphocytic leukemia)
- * Known central nervous system involvement
- * Participants with a history of confirmed progressive multifocal leukoencephalopathy (PML)
- * An individual organ/ system impairment score of 4 as assessed by the CIRS definition limiting the ability to receive the treatment regimen of this trial with the exception of eyes, ears, nose, throat organ system
- * Participants with uncontrolled autoimmune hemolytic anemia or immune thrombocytopenia
- * Inadequate renal function
- * History of prior malignancy, except for conditions as listed in the protocol if participants have recovered from the acute side effects incurred as a result of previous therapy
- * Use of investigational agents or concurrent anti-cancer treatment within the last 4 weeks of registration
- * Participants with active bacterial, viral, or fungal infection requiring systemic treatment within the last two months prior to registration
- * History of severe allergic or anaphylactic reactions to humanized or murine monoclonal antibodies or known sensitivity or allergy to murine products
- * Hypersensitivity to chlorambucil, obinutuzumab, or venetoclax or to any of the excipients
- * Pregnant women and nursing mothers
- * Positive test results for chronic hepatitis B virus (HBV) infection (defined as positive hepatitis B surface antigen [HBsAg] serology) or positive test result for hepatitis C (hepatitis C virus [HCV] antibody serology testing)
- * Participants with known infection with human immunodeficiency virus (HIV) or human T-cell leukemia virus-1 (HTLV-1)
- * Requires the use of warfarin, marcumar, or phenprocoumon
- * Received agents known to be strong and moderate Cytochrome P450 3A inhibitors or inducers within 7 days prior to the first dose of study drug

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

11. Study Identifiers & Governance

Primary ID: 551814129434349837

Registry Number: NCT02242942

Sponsor: Roche

Study Title: Comparison of the Treatments of Obinutuzumab + Venetoclax Versus Obinutuzumab + Chlorambucil in Patients With Chronic Lymphocytic Leukemia

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____