

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

## 1. Administrative & Study Identification

|                          |                                                                                                                                                                                                                                                                                                                                             |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Full Study Title:        | AN OPEN-LABEL, RANDOMIZED, CONTROLLED PHASE 3 STUDY OF SIGVOTATUG VEDOTIN IN COMBINATION WITH PEMBROLIZUMAB COMPARED WITH PEMBROLIZUMAB MONOTHERAPY AS FIRST-LINE TREATMENT IN PARTICIPANTS WITH PD-L1 HIGH (≥50% OF TUMOR CELLS EXPRESSING PD-L1), LOCALLY ADVANCED, UNRESECTABLE, OR METASTATIC NON-SMALL CELL LUNG CANCER (BE6A LUNG-02) |
| Short Title:             | This is a Study to Learn About How the Combination of the Study Medicines Sigvotatug Vedotin Plus Pembrolizumab Works in People With Non-small Cell Lung Cancer With High Levels of PD-L1.                                                                                                                                                  |
| Protocol Number:         | 590872609063007348                                                                                                                                                                                                                                                                                                                          |
| NCT Number:              | NCT06758401                                                                                                                                                                                                                                                                                                                                 |
| Sponsor Name:            | Pfizer                                                                                                                                                                                                                                                                                                                                      |
| Investigational Product: | pembrolizumab                                                                                                                                                                                                                                                                                                                               |
| Phase:                   | PHASE3                                                                                                                                                                                                                                                                                                                                      |
| Medical Monitor:         | [Not Provided in Posting]                                                                                                                                                                                                                                                                                                                   |

## 2. Study Synopsis & Rationale

### 2.1 Study Objective

|                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Primary Objective: | 1. Overall Survival: Overall survival defined as the duration from enrollment to death. 2. Progression Free Survival (PFS) assessed by blinded independent central review (BICR): Progression-free survival is defined as the time interval from the date of randomization to the date of first documented tumor progression determined by blinded independent central review (BICR) assessment as per Response Evaluation Criteria in Solid Tumors (RECIST 1.1) or death due to any cause, whichever come first. |
| Secondary Obj.:    | 1. Progression Free Survival as assessed by Investigator 2. Objective Response Rate as assessed by BICR 3. Objective Response Rate as assessed by Investigator                                                                                                                                                                                                                                                                                                                                                    |

### 2.2 Background & Rationale

The purpose of the study is to compare how the new combination treatment (Sigvotatug Vedotin plus pembrolizumab) works compared to pembrolizumab alone in patients with non-small cell lung cancer (NSCLC) with high levels of PD-L1. This is a protein that acts as a kind of "brake" to keep the body's immune responses under control.

## 3. Study Design & Methodology

|                   |                                                         |
|-------------------|---------------------------------------------------------|
| Design Framework: | Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING |
| Target N:         | 714                                                     |

## 4. Subject Selection (Eligibility Criteria)

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

## 4.1 Inclusion Criteria

1. Participants must meet the following criteria:

1. Have pathologically confirmed Stage IIIB or IIIC NSCLC and not be a candidate for surgical resection or definitive chemoradiation, or Stage IV NSCLC per the AJCC Staging Manual (Version 8.0) and the UICC Staging System (Eighth edition).
2. Participants with non-squamous histology must have documented negative test results for EGFR, ALK, and ROS1 AGAs and no known AGAs in NTRK, BRAF, RET, MET, or other AGAs with approved front-line therapies per local standard of care.
3. Large cell neuroendocrine carcinoma is excluded.
4. Candidate for treatment with pembrolizumab monotherapy per local guidelines.
2. Tumor has PD-L1 expression in ≥50% of tumor cells (TPS ≥50%) as determined by local testing
3. Measurable disease based on RECIST v1.1 per investigator.
4. Resolution of acute effects of any prior therapy to either baseline severity or NCI CTCAE Grade 1 or less (except for AEs not constituting a safety risk in the investigator's judgment), unless otherwise excluded.

## 4.2 Exclusion Criteria

1. Life expectancy of <3 months in the opinion of the investigator.
2. Any medical or psychiatric condition including recent (within the past year) or active suicidal ideation/behavior or laboratory abnormality that may increase the risk of study participation or make the participant inappropriate for the study.
3. Participants with any history of another malignancy within 3 years before the first dose of study intervention, or any evidence of residual disease from a previously diagnosed malignancy.
4. Known or suspected hypersensitivity, intolerance, or contraindication to any excipient contained in the drug formulation of sigvatug vedotin or pembrolizumab.
5. Participants with any of the following respiratory conditions:
  1. Evidence of noninfectious or drug-induced ILD or pneumonitis
  2. Known DLCO (adjusted for hemoglobin) <50% predicted.
  3. Grade ≥3 pulmonary disease unrelated to underlying malignancy
6. Known active CNS lesions are excluded. Participants with definitively treated brain metastases (surgery and/or radiotherapy) may be eligible. Clinically inactive brain metastases of longest diameter <0.5 cm are permitted.
7. Major surgery (defined as a surgery requiring inpatient hospitalization of at least 48 hours) within 21 days or minor surgery within 7 days prior to first dose of study intervention.
8. Receipt of a live vaccine within 30 days prior to first dose of study intervention.
9. Pre-existing peripheral neuropathy Grade ≥2 per NCI CTCAE v5.0.
10. Uncontrolled diabetes mellitus, defined as HbA1c ≥8.0% or HbA1c between 7.0% and 8.0% with associated diabetes symptoms (polyuria or polydipsia) that are not otherwise explained.
11. Prior immune-related AE that led to anti-PD-(L)1 treatment discontinuation, required a high-dose steroid taper (≥0.5 mg/kg prednisone or equivalent per day) for >2 weeks, or required treatment with systemic immunosuppressive therapy.

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

12. History of autoimmune disease that has required systemic treatment in the past 2 years
13. Participants with prior solid organ or bone marrow transplantation.
14. Currently receiving a high-dose steroid ( $\geq 10$  mg prednisone or equivalent per day) or other immune suppressant or has a condition requiring a chronic high-dose steroid or immune suppressant.
15. Prior and concomitant therapy:
  1. Any prior treatment with MMAE-derived drugs or IB6 targeting agents.
  2. Prior systemic therapy, including anti-PD-(L)1 therapy, for locally advanced, unresectable, or metastatic NSCLC.
    - \* (Neo)adjuvant anti-PD-(L)1 is allowed if recurrence or progression occurred  $\geq 9$  months after the last dose.
    - \* Other (neo)adjuvant or definitive therapy is allowed if recurrence or progression occurred  $\geq 6$  months after the last dose.
  3. Prior radiotherapy to the lung within 6 months of first dose of study intervention, referencing the last date radiotherapy was received.
  4. Chemotherapy, biologics, and/or other antitumor treatment with immunotherapy not specifically prohibited that is completed less than 4 weeks prior to first dose of study intervention, or 2 weeks for palliative radiotherapy.
  5. Any prior therapy with an immune-oncology agent directed to a stimulatory or co-inhibitory T-cell receptor
16. History of or current ongoing infection, including participants positive for active HIV, HBV, or HCV.
17. Severe uncontrolled cardiac or cerebrovascular condition within the previous 6 months

## 11. Study Identifiers & Governance

|                  |                                                                                                                                                                                            |
|------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Primary ID:      | 590872609063007348                                                                                                                                                                         |
| Registry Number: | NCT06758401                                                                                                                                                                                |
| Sponsor:         | Pfizer                                                                                                                                                                                     |
| Study Title:     | This is a Study to Learn About How the Combination of the Study Medicines Sigvotatug Vedotin Plus Pembrolizumab Works in People With Non-small Cell Lung Cancer With High Levels of PD-L1. |

## 15. Sign-off & Approval

Principal Investigator: \_\_\_\_\_ Date: \_\_\_\_\_  
Clinical Operations Lead: \_\_\_\_\_ Date: \_\_\_\_\_  
Lead Statistician: \_\_\_\_\_ Date: \_\_\_\_\_