

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A PHASE 3, EXTERNAL AND SYNTHETIC PLACEBO-CONTROLLED RANDOMIZED STUDY WITH DOSE-UP FOR NON-RESPONDERS TO INVESTIGATE SAFETY AND EFFICACY OF RITLECITINIB 50 MG AND 100 MG ONCE DAILY IN ADULT AND ADOLESCENT PARTICIPANTS 12 YEARS OF AGE AND OLDER WITH ALOPECIA AREATA
Short Title:	A Study of 2 Doses of Ritlecitinib in People 12 Years of Age and Older With Alopecia Areata
Protocol Number:	674030387309581743
NCT Number:	NCT06873945
Sponsor Name:	Pfizer
Investigational Product:	ritlecitinib
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of participants with absolute Severity of Alopecia Tool (SALT) score less than or equal to 20: Difference in the percentage of participants with SALT score less than or equal to 20 between ritlecitinib 100 mg once-daily (QD) versus placebo
Secondary Obj.:	1. Percentage of participants with absolute SALT score less than or equal to 20 2. Change from baseline in SALT score 3. EU Only: Percentage of participants with Patient Global Impression of Change (PGI-C) response, defined as a score of "moderately improved" or "greatly improved"

2.2 Background & Rationale

The purpose of the study is to learn about the safety and effects of the study medicine (called ritlecitinib) for the treatment of alopecia areata. Alopecia areata is a disease that causes hair loss on the scalp, face, and areas of the body.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	550

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Age:

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. 18 years of age or older at screening. Adolescents (12 to <18 years of age at screening) are also eligible for this study, but only if permitted by the local IRB/EC and local regulatory health authority (if applicable). Where these approvals have not been granted, only participants 18 years of age and older at screening will be enrolled.

Disease Characteristics:

2. Must meet the following alopecia areata criteria at both Screening and Baseline:

1. Have a clinical diagnosis of alopecia areata with no other etiology of hair loss.
2. ≥50% hair loss of the scalp, as measured by SALT, without evidence of terminal hair regrowth within the previous 6 months.
3. Current episode of hair loss ≥10 years.

4.2 Exclusion Criteria

Medical Conditions:

1. Diseases or conditions other than alopecia areata which affect hair loss, including other types of alopecia, other scalp disease that may impact the alopecia areata assessment, or active systemic diseases that may cause hair loss.
2. History of severe allergic or anaphylactoid reaction to any kinase inhibitor or a known allergy/hypersensitivity to any component (including excipients) of the study intervention.
3. Any psychiatric condition including recent or active suicidal ideation or behavior that meets protocol-defined criteria.
4. General Infection History:

- * Have a history of systemic infection requiring hospitalization or parenteral therapy (antimicrobial, antiviral, antiparasitic, antiprotozoal, or antifungal), or as otherwise judged clinically significant by the investigator, within 3 months prior to Day 1.

- * Have active acute or chronic infection requiring treatment with oral antibiotics, antivirals, antiparasitics, antiprotozoals, or antifungals within 4 weeks prior to Day 1. NOTE: participants may be rescreened after the infection resolves.

- * Evidence or history of untreated, currently treated or inadequately treated active or latent infection with *Mycobacterium tuberculosis*.

5. Specific Viral Infection History:

- * History (single episode) of disseminated herpes zoster or disseminated herpes simplex, or a recurrent (more than one episode of) localized, dermatomal herpes zoster.

- * Infected with hepatitis B or hepatitis C viruses: all participants will undergo screening for hepatitis B and C for eligibility.

6. Other Medical Conditions:

- * Have hearing loss with progression over the previous 5 years, sudden hearing loss, or middle or inner ear disease such as otitis media, cholesteatoma, Meniere's disease, labyrinthitis, or other auditory condition that is considered acute,

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

fluctuating or progressive.

- * Abnormal findings on the screening chest imaging (eg, chest x-ray) including, but not limited to, presence of active TB or other infections, cardiomyopathy, or malignancy. Chest imaging may be performed up to 12 weeks prior to Screening.

- * Have any malignancies or have a history of malignancies with the exception of adequately treated or excised nonmetastatic basal cell or squamous cell cancer of the skin or cervical carcinoma in situ.

- * Have a history of any lymphoproliferative disorder such as EBV-related lymphoproliferative disorder, history of lymphoma, history of leukemia, or signs and symptoms suggestive of current lymphatic or lymphoid disease.

- * Significant trauma or major surgery within 1 month of the first dose of study drug or considered in imminent need for surgery.

7. Adolescent participants 12 to <18 years of age without one of the following:

- * Documented evidence from a health professional of having received varicella vaccination (2 doses); or

- * Evidence of prior exposure to varicella zoster virus (VZV) based on serological testing (ie, a positive VZV IgG Ab result) at Screening.

8. Any medical or laboratory abnormality that may increase the risk of study participation or, in the investigator's judgment, make the participant inappropriate for the study.

Prior/Concomitant Therapy:

9. Current or prior use of any prohibited medication(s), vaccine(s), or treatment(s) within the protocol-defined timelines.

Prior/Concurrent Clinical Study Experience:

10. Previous administration with an investigational drug or vaccine within 8 weeks (or longer as determined by the local requirement) or 5 half-lives (whichever is longer) before the first dose of study intervention in this study. Participation in studies of other investigational products (drug or vaccine) at any time during their participation in this study.

Diagnostic Assessments:

11. Any exclusionary abnormalities in laboratory values at Screening, as assessed by the study-specific laboratory and, if deemed necessary, confirmed by a single repeat.

12. Screening standard 12-lead ECG that demonstrates clinically relevant abnormalities.

Other Exclusion Criteria:

13. Investigator site staff directly involved in the conduct of the study and their family members, site staff otherwise supervised by the investigator, and sponsor and sponsor delegate employees directly involved in the conduct of the study and their family members.

11. Study Identifiers & Governance

Primary ID: 674030387309581743

Registry Number: NCT06873945

Sponsor: Pfizer

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Study Title:

A Study of 2 Doses of Ritlecitinib in People 12 Years of Age and Older With Alopecia Areata

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____