

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Open-Label, Multicenter Randomized Study Evaluating Glofitamab as a Single Agent Versus Investigator's Choice in Patients With Relapsed/Refractory Mantle Cell Lymphoma
Short Title:	A Study to Evaluate Glofitamab as a Single Agent vs. Investigator's Choice in Participants With Relapsed/Refractory Mantle Cell Lymphoma
Protocol Number:	687385381493064446
NCT Number:	NCT06084936
Sponsor Name:	Roche
Investigational Product:	rituximab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-free survival (PFS)
Secondary Obj.:	1. Complete response (CR) rate 2. Objective response rate (ORR) 3. Overall survival (OS)

2.2 Background & Rationale

The purpose of this study is to evaluate the efficacy of glofitamab monotherapy compared with an investigator's choice of either rituximab plus bendamustine (BR), or lenalidomide with rituximab (R-Len) in patients with relapsed or refractory (R/R) mantle cell lymphoma (MCL).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	182

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Life expectancy at least 12 weeks
- * Histologically-confirmed MCL, with documentation of either overexpression of cyclin D1 or the presence of t(11:14)
- * Relapsed (disease progression after the last treatment regimen) or refractory (failure to achieve a partial or complete response from the last treatment regimen) disease
- * At least 1 line of prior systemic therapy including a BTK inhibitor and additional systemic therapy option
- * Confirmed availability of tumor tissue, unless deemed unsafe per investigator assessment
- * At least one bi-dimensionally measurable (defined as at least 1.5 cm) nodal lesion, or one bi-dimensionally measurable

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

(at least 1 cm) extranodal lesion, as measured on CT scan

- * Eastern Cooperative Oncology Group (ECOG) performance status of 0, 1, or 2
- * Negative HIV test at screening
- * Adequate hematological function

4.2 Exclusion Criteria

- * Pregnancy or breastfeeding, or intention of becoming pregnant during the study or within 3 months after the final dose of tocilizumab, 2 months after the final dose of glofitamab, whichever is longer
- * Leukemic, non-nodal MCL
- * History of severe allergic or anaphylactic reactions to humanized or murine monoclonal antibodies (or recombinant antibody-related fusion proteins) or known sensitivity or allergy to murine products
- * Contraindication to obinutuzumab or rituximab, and either bendamustine or lenalidomide
- * Prior treatment with glofitamab or other bispecific antibodies targeting both CD20 and CD3
- * Prior treatment with CAR-T cell therapy
- * Treatment with systemic therapy or BTK inhibitors, or any investigational agent for the purposes of treating cancer within 2 weeks or 5 half-lives (whichever is shorter) prior to first study treatment
- * Primary or secondary CNS lymphoma at the time of recruitment or history of CNS lymphoma
- * Current or history of CNS disease, such as stroke, epilepsy, CNS vasculitis, or neurodegenerative disease
- * History of other malignancy that could affect compliance with the protocol or interpretation of results
- * Significant or extensive cardiovascular disease
- * Known active bacterial, viral, fungal, mycobacterial, parasitic, or other infection at study enrollment or any major episode of infection within 4 weeks prior to the first study treatment
- * Suspected or latent tuberculosis
- * Positive test for hepatitis B virus (HBV) or hepatitis C virus (HCV)
- * Known or suspected chronic active Epstein-Barr viral infection (EBV)
- * Known or suspected history of hemophagocytic lymphohistiocytosis (HLH)
- * Known history of progressive multifocal leukoencephalopathy (PML)
- * Adverse events from prior anti-cancer therapy that have not resolved to Grade 1 or better
- * Administration of a live, attenuated vaccine within 4 weeks before first study treatment administration or anticipation that such a live, attenuated vaccine will be required during the study
- * Prior solid organ transplantation or allogenic stem cell transplant
- * Eligibility for stem cell transplantation (SCT)
- * Active autoimmune disease requiring treatment
- * Prior treatment with systemic immunosuppressive medications within 2 weeks or five half-lives (whichever is shorter) prior to the first dose of study treatment
- * Corticosteroid therapy within 2 weeks prior to first dose of study treatment
- * Recent major surgery (within 4 weeks before the first study treatment) other than for diagnosis
- * Clinically significant history of cirrhotic liver disease

11. Study Identifiers & Governance

Primary ID: 687385381493064446

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Registry Number: NCT06084936

Sponsor: Roche

Study Title: A Study to Evaluate Glofitamab as a Single Agent vs. Investigator's Choice in Participants With Relapsed/Refractory Mantle Cell Lymphoma

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____