

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Open-label, Phase 3 Study of MK-2870 vs. Platinum Doublets in Participants With EGFR-mutated, Advanced Nonsquamous Non-small Cell Lung Cancer (NSCLC) Who Have Progressed on Prior EGFR Tyrosine Kinase Inhibitors
Short Title:	Sacituzumab Tirumotecan (MK-2870) Versus Pemetrexed and Carboplatin Combination Therapy in Participants With Epidermal Growth Factor (EGFR)-Mutated, Advanced Nonsquamous Non-small Cell Lung Cancer (NSCLC) Who Have Progressed on Prior EGFR Tyrosine Kinase Inhibitors (MK-2870-009)
Protocol Number:	774835801660864471
NCT Number:	NCT06305754
Sponsor Name:	Merck
Investigational Product:	paracetamol
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-Free Survival (PFS): Progression-Free Survival is defined as the time from randomization to the first documented disease progression per Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST 1.1) based on blinded independent central review (BICR) or death due to any cause, whichever occurs first. PFS for all randomized participants will be reported. 2. Overall Survival (OS): Overall survival is defined as the time from randomization to death due to any cause. Overall survival for all randomized participants will be reported.
Secondary Obj.:	1. Objective Response Rate (ORR) 2. Duration of Response (DOR) 3. Change from Baseline in European Organization for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30 (EORTC QLQ-C30) Global Health Status (Item 29) and Quality of Life (Item 30) Combined Score

2.2 Background & Rationale

The purpose of this study is to evaluate sacituzumab tirumotecan versus pemetrexed in combination with carboplatin for the treatment of epidermal growth factor receptor (EGFR)-mutated advanced non-squamous non-small cell lung cancer (NSCLC). Participants in this study have NSCLC that has continued to progress on prior treatment with EGFR tyrosine kinase inhibitors (TKIs).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	520

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Histologically or cytologically confirmed diagnosis of advanced-stage nonsquamous non-small cell lung cancer (NSCLC).
- * Participants who have adverse events (AEs) due to previous anticancer therapies must have recovered to Grade ?1 or baseline.
- * Participants who are Hepatitis B surface antigen (HBsAg) positive are eligible if they have received Hepatitis B virus (HBV) antiviral therapy for at least 4 weeks and have undetectable HBV viral load.
- * Participants with history of Hepatitis C virus (HCV) infection are eligible if HCV viral load is undetectable.
- * Human immunodeficiency virus (HIV)-infected participants must have well controlled HIV on antiretroviral therapy.
- * Life expectancy of at least 3 months.

4.2 Exclusion Criteria

- * Predominantly squamous cell histology NSCLC.
- * History of second malignancy, unless potentially curative treatment has been completed with no evidence of malignancy for 3 years.
- * Grade ?2 peripheral neuropathy.
- * History of documented severe dry eye syndrome, severe Meibomian gland disease and/or blepharitis, or severe corneal disease that prevents/delays corneal healing.
- * Active inflammatory bowel disease requiring immunosuppressive medication or previous history of inflammatory bowel disease.
- * Uncontrolled, or significant cardiovascular disease or cerebrovascular disease.
- * Received prior radiotherapy within 2 weeks of start of study intervention, or radiation-related toxicities, requiring corticosteroids.
- * Received a live or live-attenuated vaccine within 30 days before the first dose of study intervention. Administration of killed vaccines is allowed.
- * Received radiation therapy to the lung that is >30 Gray within 6 months of the first dose of study intervention.
- * Known active central nervous system metastases and/or carcinomatous meningitis.
- * Active infection requiring systemic therapy.
- * History of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease.
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease.
- * Concurrent active HBV and HCV infection.
- * History of allogeneic tissue/solid organ transplant.
- * Participants who have not adequately recovered from major surgery or have ongoing surgical complications.

11. Study Identifiers & Governance

Primary ID: 774835801660864471
Registry Number: NCT06305754
Sponsor: Merck

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Study Title:

Sacituzumab Tirumotecan (MK-2870) Versus Pemetrexed and Carboplatin Combination Therapy in Participants With Epidermal Growth Factor (EGFR)-Mutated, Advanced Nonsquamous Non-small Cell Lung Cancer (NSCLC) Who Have Progressed on Prior EGFR Tyrosine Kinase Inhibitors (MK-2870-009)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____