

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A 96-week, Two-arm, Randomized, Single-masked, Multi-center, Phase III Study Assessing the Efficacy and Safety of Brolucizumab 6 mg Compared to Panretinal Photocoagulation Laser in Patients With Proliferative Diabetic Retinopathy
Short Title:	Study of Efficacy and Safety of Brolucizumab Versus Panretinal Photocoagulation Laser in Patients With Proliferative Diabetic Retinopathy
Protocol Number:	807690768812278036
NCT Number:	NCT04278417
Sponsor Name:	Novartis
Investigational Product:	brolucizumab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective: 1. Change From Baseline in Best-corrected Visual Acuity (BCVA) at Week 54 for the Study Eye: BCVA was assessed using Early Treatment Diabetic Retinopathy Study (ETDRS) visual acuity testing charts.

Visual function of the study eye was assessed using the ETDRS protocol. Participants with a BCVA ETDRS letter score of ≥ 34 ETDRS letters (Snellen equivalent 20/200) at Screening / Baseline in the study eye were included.

Min and max possible scores are 0-100 respectively. A higher score represents better functioning.

Last observation carried forward (LOCF) was used for the imputation of missing values.

Secondary Obj.: 1. Number and Percentage of Subjects With no Proliferative Diabetic Retinopathy (PDR) at Week 54 for the Study Eye 2. Number and Percentage of Subjects With no Proliferative Diabetic Retinopathy (PDR) at Week 96 for the Study Eye 3. Number and Percentage of Subjects With Center-involved Diabetic Macular Edema (CI- DME) up to Week 54 for the Study Eye

2.2 Background & Rationale

The purpose of this study was to evaluate the efficacy and safety of brolucizumab compared to panretinal photocoagulation laser (PRP) in patients with proliferative diabetic retinopathy (PDR). This evaluation will provide information that brolucizumab is non-inferior to PRP with respect to the change in best corrected visual acuity at Week 54.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE3, Status: COMPLETED

Target N: 689

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Signed informed consent must be obtained prior to participation
- * Able to complete adequate fundus photographs and retinal images
- * Diagnosis of type 1 or 2 Diabetes Mellitus (DM) and HbA1c less than or equal to 12% at screening
- * DM treatment stable for at least 3 months
- * PDR diagnosis with no previous PRP treatment in the study eye

4.2 Exclusion Criteria

- * Concomitant conditions or ocular disorders in the study eye at Screening or Baseline that could compromise a response to study treatment.
- * Presence of diabetic macular edema in the study eye
- * Active infection or inflammation in the study eye
- * Uncontrolled glaucoma (IOP greater than 25 mmHg)
- * Intravitreal anti-VEGF treatment within 6 months
- * Treatment with intraocular corticosteroids
- * End stage renal disease requiring dialysis or kidney transplant
- * Uncontrolled blood pressure
- * Systemic anti-VEGF therapy at any time

Other protocol-defined inclusion/exclusion criteria may apply.

11. Study Identifiers & Governance

Primary ID: 807690768812278036

Registry Number: NCT04278417

Sponsor: Novartis

Study Title: Study of Efficacy and Safety of Brolucizumab Versus Panretinal Photocoagulation Laser in Patients With Proliferative Diabetic Retinopathy

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____