

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase IIIb Multicenter, Randomized, Double-blind, Controlled Study to Evaluate the Efficacy, Safety and Pharmacokinetics of a Higher Dose of Ocrelizumab in Adults With Relapsing Multiple Sclerosis
Short Title:	A Study to Evaluate the Efficacy, Safety and Pharmacokinetics (PK) of a Higher Dose of Ocrelizumab in Adults With Relapsing Multiple Sclerosis (RMS)
Protocol Number:	909857978771445983
NCT Number:	NCT04544436
Sponsor Name:	Roche
Investigational Product:	chlorcyclizine
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Time to Onset of 12-week Composite Confirmed Disability Progression (cCDP12): Time to onset of cCDP is defined as the first occurrence of a predefined confirmed progression event measured by Expanded Disability Status Scale (EDSS), Timed 25-foot Walk Test (T25FWT) or 9-hole Peg Test (9-HPT).
Secondary Obj.:	1. Time to Onset of 24-week cCDP (cCDP24) 2. Time to Onset of 48-week cCDP (cCDP48) 3. Time to Onset of cCDP12 Independent of Protocol-defined Relapses (PDR)

2.2 Background & Rationale

This is a randomized, double-blind, controlled, parallel group, multicenter study to evaluate efficacy, safety and PK of a higher dose of ocrelizumab per intravenous (IV) infusion every 24 weeks (Q24W) in participants with RMS, in comparison to the approved 600 milligrams (mg) dose of ocrelizumab.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	864

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Diagnosis of RMS
- * At least two documented clinical relapses within the last 2 years prior to screening, or one clinical relapse in the year prior to screening. No relapse 30 days prior to screening and at baseline

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Participants must be neurologically stable for at least 30 days prior to randomization and baseline
- * EDSS score, at screening and baseline, from 0 to 5.5 inclusive
- * Average T25FWT score over two trials at screening and over two trials at baseline respectively, up to 150 (inclusive) seconds
- * Average 9HPT score over four trials at screening and over four trials at baseline respectively, up to 250 (inclusive) seconds
- * Documented magnetic resonance imaging (MRI) of brain with abnormalities consistent with MS at screening
- * Participants requiring symptomatic treatment for MS and/or physiotherapy must be treated at a stable dose. No initiation of symptomatic treatment for MS or physiotherapy within 4 weeks of randomization
- * Females of childbearing potential, agreement to remain abstinent or use adequate contraceptive methods
- * Female participants without reproductive potential may be enrolled e.g. if post-menopausal or if surgically sterile

4.2 Exclusion Criteria

- * History of primary progressive MS at screening
- * Any known or suspected active infection at screening or baseline (except nailbed infections), or any major episode of infection requiring hospitalization or treatment with IV antimicrobials within 8 weeks or treatment with oral antimicrobials within 2 weeks, prior to and during screening
- * History of confirmed or suspected progressive multifocal leukoencephalopathy
- * History of cancer, including hematologic malignancy and solid tumors, within 10 years of screening
- * Immunocompromised state
- * Receipt of a live or live-attenuated vaccine within 6 weeks prior to randomization
- * Inability to complete an MRI or contraindication to gadolinium administration
- * Contraindications to mandatory pre-medications for infusion-related reaction (IRRs)
- * Known presence of other neurologic disorders that could interfere with the diagnosis of MS or assessments of efficacy and/or safety during the study
- * Any concomitant disease that may require chronic treatment with systemic corticosteroids or immunosuppressants during the course of the study
- * Significant, uncontrolled disease that may preclude participant from participating in the study
- * History of or currently active primary or secondary, non-drug-related, immunodeficiency
- * Pregnant or breastfeeding or intending to become pregnant
- * Lack of peripheral venous access
- * History of alcohol or other drug abuse within 12 months prior to screening
- * Treatment with any investigational agent within 24 weeks prior to screening or treatment with any experimental procedure for MS
- * Previous use of anti- cluster of differentiation 20 (CD20s) (including ocrelizumab), unless the last infusion was more than 2 years before screening, B-cell count is normal, and the stop of the treatment was not motivated by safety reasons or lack of efficacy
- * Previous treatment with fingolimod, siponimod, or ozanimod within 6 weeks of baseline
- * Previous treatment with natalizumab within 4.5 months of baseline
- * Previous treatment with interferons beta (1a or 1b), or glatiramer acetate within 2 weeks of baseline
- * Any previous treatment with mitoxantrone, cladribine, atacept, alemtuzumab, and daclizumab - Previous treatment

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

with any other immunomodulatory or immunosuppressive medication not already listed above without appropriate washout as described in the applicable local label. If the washout requirements are not described in the applicable local label, then the wash out period must be five times the half-life of the medication

- * Any previous treatment with bone marrow transplantation and hematopoietic stem cell transplantation
- * Any previous history of transplantation or anti-rejection therapy
- * Treatment with IV immunoglobulin (Ig) or plasmapheresis within 12 weeks prior to randomization
- * Systemic corticosteroid therapy within 4 weeks prior to screening
- * Positive screening tests for active, latent, or inadequately treated hepatitis B
- * Sensitivity or intolerance to any ingredient (including excipients) of ocrelizumab
- * Any additional exclusionary criterion as per ocrelizumab local label, if more stringent than the above

11. Study Identifiers & Governance

Primary ID: 909857978771445983

Registry Number: NCT04544436

Sponsor: Roche

Study Title: A Study to Evaluate the Efficacy, Safety and Pharmacokinetics (PK) of a Higher Dose of Ocrelizumab in Adults With Relapsing Multiple Sclerosis (RMS)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____