

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 3 Randomized, Open-label Clinical Study to Evaluate the Pharmacokinetics and Safety of Subcutaneous Pembrolizumab Coformulated With Hyaluronidase (MK-3475A) Versus Intravenous Pembrolizumab, Administered With Chemotherapy, in the First-line Treatment of Participants With Metastatic Non-small Cell Lung Cancer
Short Title:	A Study of Subcutaneous (SC) Pembrolizumab Coformulated With Berahyaluronidase Alfa (MK-3475A) vs Intravenous Pembrolizumab in Adult Participants With Metastatic Non-small Cell Lung Cancer (NSCLC) (MK-3475A-D77)-Japan Extension
Protocol Number:	324789051674853911
NCT Number:	NCT06212752
Sponsor Name:	Merck
Investigational Product:	cisplatin
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Objective Response Rate (ORR): ORR was defined as the percentage of participants who have a confirmed complete response (CR: disappearance of all target lesions) or partial response (PR: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters) per Response Evaluation Criteria In Solid Tumors 1.1 (RECIST 1.1) as assessed by blinded independent central review (BICR). The percentage of participants with CR or PR were reported.
Secondary Obj.:	1. Cycle 1: Area Under the Curve From Time 0 to 6 Weeks (AUC0-6 Weeks) of Pembrolizumab After the First Dose 2. Cycle 3: Trough Serum Concentration (C _{trough}) of Pembrolizumab at Steady State 3. Cycle 1: Maximum Serum Concentration (C _{max}) of Pembrolizumab After the First Dose

2.2 Background & Rationale

This study is to assess the pharmacokinetics (PK) and safety of SC pembrolizumab (+) berahyaluronidase alfa vs intravenous (IV) pembrolizumab, administered with chemotherapy in first line treatment of adult Japanese participants with metastatic non-small cell lung cancer. The primary hypotheses of this study are pembrolizumab (+) berahyaluronidase alfa subcutaneous (SC) is noninferior to pembrolizumab IV with respect to PK parameters.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	39

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has histologically or cytologically confirmed diagnosis of squamous or non-squamous Non-small Cell Lung Cancer (NSCLC)
- * Must provide archival tumor tissue sample or newly obtained core, incisional, or excisional biopsy of a tumor lesion not previously irradiated
- * Has a life expectancy of at least 3 months

4.2 Exclusion Criteria

- * Has a diagnosis of small cell lung cancer or, for mixed tumors, presence of small cell elements
- * Has received prior systemic anticancer therapy for metastatic NSCLC
- * Has received prior systemic anticancer therapy including investigational agents within 4 weeks before randomization
- * Has received prior radiotherapy within 2 weeks of start of study intervention or has radiation-related toxicity requiring corticosteroids
- * Has received radiation therapy to the lung (≥ 30 Gray) within 6 months of start of study intervention
- * Has received a live or live-attenuated vaccine within 30 days before the first dose of study intervention
- * Has a diagnosis of immunodeficiency or is receiving chronic systemic steroid therapy
- * Has a known additional malignancy that is progressing or has required active treatment within the past 3 years
- * Has an active autoimmune disease that has required systemic treatment in past 2 years
- * Has an active infection requiring systemic therapy
- * Has a history of human immunodeficiency virus (HIV) infection
- * Has a history of Hepatitis B or C
- * Has not adequately recovered from major surgery or has ongoing surgical complications
- * Has a history of allogenic tissue/solid organ transplant

11. Study Identifiers & Governance

Primary ID:	324789051674853911
Registry Number:	NCT06212752
Sponsor:	Merck
Study Title:	A Study of Subcutaneous (SC) Pembrolizumab Coformulated With Berahyaluronidase Alfa (MK-3475A) vs Intravenous Pembrolizumab in Adult Participants With Metastatic Non-small Cell Lung Cancer (NSCLC) (MK-3475A-D77)-Japan Extension

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____