

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	An Open-label, Randomized Phase 3 Study to Evaluate Efficacy and Safety of Pembrolizumab (MK-3475) in Combination With Belzutifan (MK-6482) and Lenvatinib (MK-7902), or MK-1308A in Combination With Lenvatinib, Versus Pembrolizumab and Lenvatinib, as First-Line Treatment in Participants With Advanced Clear Cell Renal Cell Carcinoma (ccRCC)
Short Title:	A Study of Pembrolizumab (MK-3475) in Combination With Belzutifan (MK-6482) and Lenvatinib (MK-7902), or Pembrolizumab/Quavonlimab (MK-1308A) in Combination With Lenvatinib, vs Pembrolizumab and Lenvatinib, for Treatment of Advanced Clear Cell Renal Cell Carcinoma (MK-6482-012)-China Extension Study
Protocol Number:	528694434473181104
NCT Number:	NCT05899049
Sponsor Name:	Merck
Investigational Product:	lenvatinib
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression Free Survival (PFS) According to Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST 1.1) as Assessed by Blinded Independent Central Review (BICR): PFS is defined as the time from randomization to the first documented progressive disease (PD) or death due to any cause, whichever occurs first. Per RECIST 1.1, PD is defined as ≥20% increase in the sum of diameters of target lesions. In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of ≥5 mm. The appearance of one or more new lesions is also considered PD. PFS as assessed by BICR based on RECIST 1.1 will be presented. 2. Overall Survival (OS): OS is defined as the time from randomization to death due to any cause.
Secondary Obj.:	1. Objective Response Rate (ORR) Per RECIST 1.1 as Assessed by BICR 2. Duration of Response (DOR) Per RECIST 1.1 as Assessed by BICR 3. Number of Participants Who Experienced At least One Adverse Event (AE)

2.2 Background & Rationale

The goal of this China extension study is to evaluate the efficacy and safety of pembrolizumab plus belzutifan plus lenvatinib or pembrolizumab/quavonlimab plus lenvatinib versus pembrolizumab plus lenvatinib as first-line treatment in Chinese participants with advanced clear cell renal cell carcinoma (ccRCC).

3. Study Design & Methodology

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Design Framework: Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N: 249

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has histologically confirmed diagnosis of RCC with clear cell component.
- * Has received no prior systemic therapy for advanced ccRCC
- * Male participants are abstinent from heterosexual intercourse or agree to use contraception during and for at least 7 days after last dose of study intervention with belzutifan and lenvatinib.
- * Female participants are not pregnant or breastfeeding and are either not a woman of child-bearing potential (WOCBP) or use a contraceptive method that is highly effective or are abstinent from heterosexual intercourse during the intervention period and for at least 120 days after pembrolizumab or pembrolizumab/quavonlimab or for at least 30 days after last dose of lenvatinib or belzutifan, whichever occurs last
- * Has adequately controlled blood pressure with or without antihypertensive medications
- * Has adequate organ function.
- * Participants receiving bone resorptive therapy must have therapy initiated at least 2 weeks prior to randomization/allocation

4.2 Exclusion Criteria

- * Has a known additional malignancy that is progressing or has required active treatment within the past 3 years
- * Has had major surgery, other than nephrectomy within 4 weeks prior to randomization
- * Has known central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Has received prior radiotherapy within 2 weeks prior to first dose of study intervention
- * Has hypoxia or requires intermittent supplemental oxygen or requires chronic supplemental oxygen
- * Has clinically significant cardiac disease within 12 months from first dose of study intervention
- * Has a history of interstitial lung disease
- * Has symptomatic pleural effusion; a participant who is clinically stable following treatment of this condition is eligible
- * Has preexisting gastrointestinal or non-gastrointestinal fistula
- * Has a diagnosis of immunodeficiency or is receiving chronic systemic steroid therapy or any other form of immunosuppressive therapy within 7 days prior to the first dose of study treatment
- * Has a known psychiatric or substance abuse disorder that would interfere with requirements of the study
- * Has received a live or live-attenuated vaccine within 30 days before the first dose of study drug; killed vaccines are allowed
- * Has an active autoimmune disease that has required systemic treatment in the past 2 years
- * Has a history of noninfectious pneumonitis that required steroids or has current pneumonitis
- * Has an active infection requiring systemic therapy
- * Has a known history of human immunodeficiency virus (HIV) infection
- * Has a known history of Hepatitis B
- * Has radiographic evidence of intratumoral cavitation, encasement or invasion of a major blood vessel
- * Has clinically significant history of bleeding within 3 months prior to randomization

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* Has had an allogenic tissue/solid organ transplant

11. Study Identifiers & Governance

Primary ID: 528694434473181104

Registry Number: NCT05899049

Sponsor: Merck

Study Title: A Study of Pembrolizumab (MK-3475) in Combination With Belzutifan (MK-6482) and Lenvatinib (MK-7902), or Pembrolizumab/Quavonlimab (MK-1308A) in Combination With Lenvatinib, vs Pembrolizumab and Lenvatinib, for Treatment of Advanced Clear Cell Renal Cell Carcinoma (MK-6482-012)-China Extension Study

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____