

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 3, Multicenter, Randomized, Double-blind, Placebo-controlled Study to Evaluate the Efficacy and Safety of Suvorexant for the Treatment of Insomnia in Participants With Opioid Use Disorder
Short Title:	A Study of Suvorexant (MK-4305) for the Treatment of Insomnia Disorder in Participants With Opioid Use Disorder (MK-4305-098)
Protocol Number:	976173145632929565
NCT Number:	NCT06655883
Sponsor Name:	Merck
Investigational Product:	suvorexant
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Change from Baseline in Total Sleep Time at Week 8: Total sleep time will be measured in a sleep laboratory by polysomnography at the participant's habitual bedtime. Change from baseline in total sleep time at Week 8 will be reported. 2. Number of Participants Who Experience One or More Adverse Events (AEs): An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. 3. Number of Participants Who Experience One or More Serious Adverse Events (SAEs): An SAE is any untoward medical occurrence that at any dose results in death, is life threatening, requires or prolongs inpatient hospitalization, results in persistent or significant disability or incapacity, is a congenital anomaly or birth defect or is another important medical event deemed such by medical or scientific judgement.
Secondary Obj.:	1. Change from Baseline in Wakefulness after Persistent Sleep Onset at Week 8 2. Number of Participants Who are Positive for Substance Abuse at Day 1 3. Number of Participants Who are Positive for Substance Abuse at Day 4

2.2 Background & Rationale

People with opioid use disorder (OUD) can have trouble falling or staying asleep. Researchers want to know if suvorexant will help people with OUD fall asleep and stay asleep. The goal of this study is to learn about the safety of suvorexant and how well people tolerate it. Researchers also want to learn if suvorexant helps people sleep longer compared to people who take placebo. A placebo looks like the study medicine but has no actual study medicine in it.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
-------------------	---

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Target N: 300

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Has a primary diagnosis of OUD according to Diagnostic and Statistical Manual of Mental Disorder, 5th Edition (DSM-5), and confirmed through the Mini International Neuropsychiatric Interview (MINI).
- * Is on a verified, stable dose of medications for opioid use disorder (MOUD) treatment.
- * Meets DSM-5 criteria for the diagnosis of Insomnia Disorder
- * Has a regular bedtime between 8 PM (20:00) and 1 AM (01:00) and is willing to maintain it for the duration of the study.
- * Has not used opioids for a period of at least 4 weeks before entering the study.

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * Has current uncontrolled major co-morbid psychiatric illness including major depressive disorder, bipolar disorder, schizophrenia, or any psychiatric condition with psychotic features.
- * Has current diagnosis or history within 5 years of any of the following: narcolepsy, sleep paralysis, severe periodic limb movement disorder, restless leg syndrome, cataplexy, circadian rhythm sleep disorder, parasomnia including nightmare disorder, sleep terror disorder, sleepwalking disorder, rapid eye movement (REM) behavior disorder, significant degree of sleep-related breathing disorder, excessive daytime sleepiness (EDS), or primary hypersomnia.
- * Is at imminent risk of self-harm.
- * Has a known history of stroke that may confound the diagnosis of insomnia.
- * Has a clinically significant movement disorder such as akinesia.
- * Has a history of hepatitis or liver disease.
- * Has habitual use of central nervous system (CNS)-depressants or stimulants that may be responsible for the participant's disturbed sleep.
- * Has a history of malignancy, ≥3 years prior to start of study, with the exception of nonmelanoma skin cancer, prostate cancer or localized carcinoma in situ of the cervix.
- * Has a history of hypersensitivity to more than 3 chemical classes of drugs, including prescription and over-the-counter medications.
- * Has donated blood products or had phlebotomy within 8 weeks prior to start of study.
- * Has a history of transmeridian travel within 2 weeks prior to start of study.

11. Study Identifiers & Governance

Primary ID: 976173145632929565

Registry Number: NCT06655883

Sponsor: Merck

Study Title: A Study of Suvorexant (MK-4305) for the Treatment of Insomnia Disorder in Participants

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

With Opioid Use Disorder (MK-4305-098)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____