

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 2, Randomized, Double-Blind, Placebo- and Active-Comparator-Controlled Clinical Study of V940 (mRNA-4157) Plus Pembrolizumab Versus Placebo Plus Pembrolizumab in Participants With First-Line Advanced Melanoma (INTerpath-012)
Short Title:	A Clinical Study of V940 and Pembrolizumab (MK-3475) in People With Melanoma (V940-012/INTerpath-012)
Protocol Number:	1030324190384348278
NCT Number:	NCT06961006
Sponsor Name:	Merck
Investigational Product:	pembrolizumab
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-free Survival (PFS): PFS is defined as the time from randomization to the first documented disease progression per Response Criteria in Solid Tumors Version 1.1 (RECIST 1.1) by investigator assessment or death due to any cause, whichever occurs first. Progressive disease (PD) is defined as ≥20% increase in the sum of diameters of target lesions. In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of ≥5 mm. The appearance of one or more new lesions is also considered PD. PFS per RECIST 1.1 as assessed by investigator will be presented.
Secondary Obj.:	1. Objective Response Rate (ORR) 2. Duration of Response (DOR) 3. Overall Survival (OS)

2.2 Background & Rationale

Researchers want to learn if V940 with pembrolizumab can stop advanced melanoma from growing or spreading. Melanoma is a type of skin cancer. Advanced means the cancer has spread to other parts of the body and cannot be removed with surgery. A standard (or usual) treatment for advanced melanoma is immunotherapy. Immunotherapy is a treatment that helps the immune system fight cancer. V940 is a study treatment designed to help a person's immune system attack their specific cancer. Pembrolizumab is an immunotherapy.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING
Target N:	160

4. Subject Selection (Eligibility Criteria)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Has unresectable and histologically confirmed Stage III or IV cutaneous melanoma per American Joint Committee on Cancer (AJCC) Eighth Edition guidelines.
- * Has been untreated for melanoma except if participant received prior adjuvant or neoadjuvant therapy with targeted therapy or immunotherapy (such as anti-cytotoxic T-lymphocyte-associated protein 4 [CTLA-4], anti-programmed cell death 1 protein [PD-1] therapy or interferon), and only if relapse did not occur within 12 months after treatment discontinuation.
- * Have documentation of serine/threonine-protein kinase B-raf (BRAF) V600-activating mutation status or had BRAF V600 mutation testing per local institutional standards during the screening period (participants with BRAF mutation positive melanoma as well as BRAF wild-type or unknown are eligible).
- * Have the presence of at least 1 measurable lesion by computed tomography (CT) or magnetic resonance imaging (MRI) per Response Criteria in Solid Tumors Version 1.1 (RECIST 1.1) as determined by the local site investigator/radiology assessment.
- * Provides tumor tissue (preferably from a metastatic site and, if not available, from the primary tumor) that is suitable for next generation sequencing and biomarker analysis as required for this study.
- * Participants with human immunodeficiency virus (HIV) must have well controlled HIV on antiretroviral therapy (ART).
- * Participants who are hepatitis B surface antigen (HBsAg) positive are eligible if they have received hepatitis B virus (HBV) antiviral therapy for at least 4 weeks, and have undetectable HBV viral load prior to randomization.
- * Participants with history of hepatitis C virus (HCV) infection are eligible if HCV viral load is undetectable at screening.

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * Has clinically significant heart failure, defined as New York Heart Association class III or IV, within the past 6 months, unless the disease is well controlled in the opinion of the investigator.
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease.
- * Has ocular or mucosal melanoma.
- * Received transfusion of blood products (including platelets or red blood cells) or administration of colony-stimulating factors (including granulocyte colony-stimulating factor, granulocyte macrophage colony-stimulating factor, or recombinant erythropoietin) within 2 weeks of the Screening blood sample (including the blood sample for V940 generation).
- * Received prior therapy with an anti-PD-1, anti-PD-L1, or anti-PD-L2 agent, or with an agent directed to another stimulatory or coinhibitory T-cell receptor (eg, CTLA-4, lymphocyte activation gene 3 [LAG-3], tumor necrosis factor receptors [OX-40 or CD137]), with some exceptions.
- * Received prior systemic anticancer therapy for melanoma before randomization, with some exceptions.
- * Received prior radiotherapy within 2 weeks of start of study intervention or has ongoing radiation related toxicities.
- * Received a live or live-attenuated vaccine within 30 days before the first dose of study intervention.
- * Received prior treatment with another universal or personalized cancer vaccine.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

11. Study Identifiers & Governance

Primary ID: 1030324190384348278

Registry Number: NCT06961006

Sponsor: Merck

Study Title: A Clinical Study of V940 and Pembrolizumab (MK-3475) in People With Melanoma (V940-012/INTerpath-012)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____