

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Open-label, Multi-arm, Two-part, Phase II Study to Assess Efficacy and Safety of Multiple LXH254 Combinations in Patients With Previously Treated Unresectable or Metastatic BRAFV600 or NRAS Mutant Melanoma
Short Title:	Study of Efficacy and Safety of LXH254 Combinations in Patients With Previously Treated Unresectable or Metastatic Melanoma
Protocol Number:	1035438760563737702
NCT Number:	NCT04417621
Sponsor Name:	Novartis
Investigational Product:	trametinib
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Overall Response Rate: Confirmed ORR using RECIST v1.1, per local assessment
Secondary Obj.:	1. Duration of Reposnse (DOR) 2. Progression Free Survival (PFS) 3. Disease Control Rate (DCR)

2.2 Background & Rationale

The primary purpose of this study is to evaluate the efficacy of LXH254 combinations in previously treated unresectable or metastatic melanoma

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: ACTIVE_NOT_RECRUITING
Target N:	134

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Male or female must be ≥ 12 years For adolescents only (12-17 years): body weight ≥ 40kg Histologically confirmed unresectable or metastatic cutaneous melanoma

Previously treated for unresectable or metastatic melanoma:

* Participants with NRAS mutation:

* Participants must have received prior systemic therapy for unresectable or metastatic melanoma with checkpoint inhibitors (CPI), either an anti-PD-1/PD-L1 as a single agent or in combination with anti-CTLA-4, investigational agents,

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

chemotherapy or locally directed anti-neoplastic agents.

- * A maximum of two prior lines of systemic CPI-containing immunotherapy for unresectable or metastatic melanoma are allowed. Additional agents administered with CPI are permitted.
- * To rule out pseudo-progression, participants must have documented confirmed progressive disease as per RECIST v1.1 while on/after treatment with checkpoint inhibitor therapy. Confirmation is not required for patients who remained on treatment for >6 months.
- * Participants with BRAFV600 mutant disease:
 - * Participants must have received prior systemic therapy for unresectable or metastatic melanoma with checkpoint inhibitors (CPI), either an anti-PD-1/PD-L1 as a single agent or in combination with anti-CTLA-4, investigational agents, chemotherapy or locally directed anti-neoplastic agents. Additionally, participants must have received targeted therapy with a RAFi as a single agent or in combination with a MEKi (+/- CPI allowed) as the last prior therapy.
- * A maximum of two prior lines of systemic CPI-containing immunotherapy for unresectable or metastatic melanoma are allowed. Additional agents with CPI are permitted.
- * A maximum of one line of targeted therapy is allowed, and it must be the most recent line of therapy.
- * Participants must have documented progressive disease as per RECIST v1.1 while on/after treatment with targeted therapy.

Other protocol-defined inclusion criteria may apply.

4.2 Exclusion Criteria

Treatment with any of the following anti-cancer therapies prior to the first dose of study treatment within the stated timeframes:

- * ? 4 weeks for radiation therapy or ? 2 weeks for limited field radiation for palliation prior to the first dose of study treatment.
- * ? 2 weeks for small molecule therapeutics.
- * ? 4 weeks for any immunotherapy treatment including immune checkpoint inhibitors.
- * ? 4 weeks for chemotherapy agents, locally directed anti-neoplastic agents, or other investigational agents.
- * ? 6 weeks for cytotoxic agents with major delayed toxicities, such as nitrosourea and mitomycin c.

Participants participating in additional parallel investigational drug or medical device studies.

All primary central nervous system (CNS) tumors or symptomatic CNS metastases that are neurologically unstable
History or current evidence of retinal vein occlusion (RVO) or current risk factors for RVO (e.g. uncontrolled glaucoma or ocular hypertension, history of hyperviscosity or hypercoagulability syndromes).

Any medical condition that would, in the investigator's judgment, prevent the patient's participation in the clinical study due to safety concerns or compliance with clinical study procedures.

Other protocol-defined exclusion criteria may apply

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

11. Study Identifiers & Governance

Primary ID: 1035438760563737702

Registry Number: NCT04417621

Sponsor: Novartis

Study Title: Study of Efficacy and Safety of LXH254 Combinations in Patients With Previously Treated Unresectable or Metastatic Melanoma

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____