

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase II, Multicenter, Randomized, Double Masked, Active Comparator-Controlled Study to Investigate the Efficacy, Safety, Tolerability, Pharmacokinetics, and Pharmacodynamics of RO7200220 in Combination With Ranibizumab Administered Intravitreally in Patients With Diabetic Macular Edema
Short Title:	A Study to Investigate Vamikibart (RO7200220) in Combination With Ranibizumab in Diabetic Macular Edema
Protocol Number:	535366253228634946
NCT Number:	NCT05151744
Sponsor Name:	Roche
Investigational Product:	ranibizumab
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Change From Baseline in Best Corrected Visual Acuity (BCVA) Averaged Over Week 44 and Week 48 in Treatment-naïve Participants: BCVA was measured via Early Treatment Diabetic Retinopathy Study (ETDRS) chart at a starting distance of 4 meters using a set of three Precision vision TM or Lighthouse distance acuity charts (modified ETDRS Charts 1, 2, and R) prior to dilating eyes by a trained and certified visual acuity (VA) examiner. The BCVA letter score ranges from 0 to 100 letters. Higher scores and gain in BCVA from baseline indicate improvement in VA. This analysis used a Mixed Model for Repeated Measurements (MMRM) model.
Secondary Obj.:	1. Number of Participants With Systemic and Ocular Adverse Events (AEs) 2. Change From Baseline in BCVA Averaged Over Week 44 and Week 48 in Previously Treated Participants 3. Change From Baseline in BCVA Averaged Over Week 44 and Week 48 in Overall ITT Population

2.2 Background & Rationale

Study BP43464 is a phase II, multicenter, randomized, double-masked active comparator-controlled study designed to assess the efficacy, safety, tolerability, pharmacokinetics (PK) and pharmacodynamics (PD) of vamikibart in combination with, anti-vascular endothelial growth factor (VEGF) inhibitor, ranibizumab compared with ranibizumab alone in participants with diabetic macular edema. Only one eye will be chosen as the study eye. The duration of the study will be 76 weeks.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: COMPLETED
-------------------	--

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Target N: 187

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Diagnosis of diabetes mellitus (Type 1 or Type 2)
- * Macular thickening secondary to diabetic macular edema (DME) involving the center of the macula
- * Decreased visual acuity attributable primarily to DME
- * Ability and willingness to provide written informed consent and to comply with the study protocol
- * Willingness to allow Aqueous Humor collection
- * For women of childbearing potential: agreement to remain abstinent or use at least one highly effective contraceptive method that results in a failure rate of $\leq 1\%$ per year during the treatment period and for at least 12 weeks after the final dose of study treatment

4.2 Exclusion Criteria

- * Hemoglobin A1c (HbA1c) of greater than ($>$) 12%
- * Uncontrolled blood pressure, defined as a systolic value greater than ($>$) 180 millimeters of mercury (mmHg) and/or a diastolic value > 100 mmHg while a patient is at rest
- * Currently pregnant or breastfeeding, or intend to become pregnant during the study
- * Prior treatment with panretinal photocoagulation or macular laser to the study eye
- * Any intraocular or periocular corticosteroid treatment within the past 16 weeks prior to Day 1 to the study eye
- * Prior Iluvien or Retisert implants within 3 years prior to Day 1 to the study eye
- * Prior or concomitant treatment with anti-VEGF therapy within 8 weeks prior to Day 1 to the study eye; VabysmoTM within 16 weeks prior to Day 1, prior Beovu[®] is not permitted
- * Prior administration of IVT brodalumab (Beovu[®]): ever; vabysmo: ≤ 24 weeks prior to Day 1) in either eye
- * Any proliferative diabetic retinopathy
- * Active intraocular or periocular infection or active intraocular inflammation in the study eye
- * Any current or history of ocular disease other than DME that may confound assessment of the macula or affect central vision in the study eye
- * Any current ocular condition which, in the opinion of the investigator, is currently causing or could be expected to contribute to irreversible vision loss due to a cause other than DME in the study eye
- * Other protocol-specified inclusion/exclusion criteria may apply

11. Study Identifiers & Governance

Primary ID:	535366253228634946
Registry Number:	NCT05151744
Sponsor:	Roche
Study Title:	A Study to Investigate Vamikibart (RO7200220) in Combination With Ranibizumab in Diabetic Macular Edema

15. Sign-off & Approval

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____