

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase II, Open Label, Randomized, Parallel Arm Study of NIS793 (With and Without Spartalizumab) in Combination With SOC Chemotherapy Gemcitabine/Nab-paclitaxel, and Gemcitabine/Nab-paclitaxel Alone in First-line Metastatic Pancreatic Ductal Adenocarcinoma (mPDAC)
Short Title:	Study of Efficacy and Safety of NIS793 (With and Without Spartalizumab) in Combination With SOC Chemotherapy in First-line Metastatic Pancreatic Ductal Adenocarcinoma (mPDAC)
Protocol Number:	63827394175507268
NCT Number:	NCT04390763
Sponsor Name:	Novartis
Investigational Product:	gemcitabine
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Safety run-in Part: Number of Participants With Dose-Limiting Toxicities (DLTs): A dose-limiting toxicity (DLT) is defined as an adverse event or abnormal laboratory value of Common Terminology Criteria for Adverse Events (CTCAE) grade $\geq$ 3 where the relationship to study treatment cannot be ruled out and is not clearly related solely to disease, disease progression, inter-current illness or concomitant medications, which occurs within the DLT evaluation period. The DLT evaluation period is the first 28 days of treatment with NIS793 with spartalizumab in combination with gemcitabine/nab-paclitaxel. Other clinically significant toxicities may be considered to be DLTs, even if not CTCAE grade 3 or higher. 2. Safety run-in Part: Number of Participants With AEs and SAEs During the On-treatment Period: Number of participants with AEs (any adverse events regardless of seriousness) and serious adverse events (SAEs), including changes from baseline in vital signs, electrocardiograms and laboratory results qualifying and reported as AEs.
--------------------	--

AE grades to characterize the severity of the AEs were based on the Common Terminology Criteria for Adverse Events (CTCAE) version 5.0. For CTCAE v5.0, Grade 1 = mild; Grade 2 = moderate; Grade 3 = severe; Grade 4 = life-threatening; Grade 5 = death related to AE.

The on-treatment period is defined from the day of first administration of any study treatment up to 30 days after the date of the last actual administration of any study drug.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. Safety run-in Part: Number of Participants With Dose Reductions and Dose Interruptions of NIS793, Spartalizumab, Gemcitabine and Nab-paclitaxel: Number of participants with at least one dose reduction and at least one dose interruption of study drugs. Dose adjustments were permitted for patients who did not tolerate the protocol-specified dosing schedule.

No dose reductions were allowed for NIS793 and spartalizumab beyond the first 28 days period of Safety run-in part.

Secondary Obj.:

1. Randomized Part: Number of Participants With AEs and SAEs During the On-treatment Period 2. Randomized Part: Number of Participants With Dose Reductions and Dose Interruptions of NIS793, Spartalizumab, Gemcitabine and Nab-paclitaxel 3. Randomized Part: Dose Intensity of NIS973 and Spartalizumab

2.2 Background & Rationale

The purpose of this Phase II study is to assess the efficacy and safety of NIS793 with and without spartalizumab in combination with gemcitabine/nab-paclitaxel versus gemcitabine/nab-paclitaxel in previously untreated mPDAC.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: TERMINATED

Target N: 164

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

1. Signed informed consent must be obtained prior to participation in the study.
2. Male or female ? 18 years of age at the time of informed consent.
3. Participants with histologically or cytologically confirmed treatment-naïve metastatic adenocarcinoma of the pancreas with measurable disease as per RECIST 1.1.
4. Participants must have a site of disease amenable to biopsy, and be candidate for tumor biopsy according to the treating institution's guidelines. Participants must be willing to undergo a tumor biopsy at screening and during therapy on the study. In the event a new biopsy cannot be safely performed at study entry, an archival sample (collected \<6 months prior) may be substituted following documented discussion with Novartis.
5. ECOG performance status ? 1.

4.2 Exclusion Criteria

1. Previous radiotherapy, surgery (with exception of placement of biliary stent, which is allowed), chemotherapy or any other investigational therapy for the treatment of metastatic pancreatic cancer. Participants having received previous chemotherapy in the adjuvant setting.
2. Participants amenable to potentially curative resection.
3. Participants with a diagnosis of pancreatic neuroendocrine tumors (NETs), acinar, or islet cell tumors.
4. Having out of range laboratory values as pre-defined in the protocol.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

5. Participants with MSI-H pancreatic adenocarcinoma.
6. Presence of symptomatic CNS metastases, or CNS metastases that require local CNS directed therapy (such as radiotherapy or surgery), or increasing doses of corticosteroids 2 weeks prior to study entry.
7. History of severe hypersensitivity reactions to any ingredient of study drug(s) and other mAbs and/or their excipients.
8. The participant exhibits any of the events outlined in the contra-indications or special warnings and precautions sections of gemcitabine and nab-paclitaxel as per locally approved labels.
9. Impaired cardiac function or clinically significant cardiac disease.
10. Known history of testing positive HIV infection.
11. Active HBV or HCV infection. Participants whose disease is controlled under antiviral therapy should not be excluded.
12. History of or current interstitial lung disease or pneumonitis grade ≥ 2
13. High risk of clinically significant gastrointestinal tract bleeding or any other condition associated with or history of significant bleeding.

11. Study Identifiers & Governance

Primary ID:	63827394175507268
Registry Number:	NCT04390763
Sponsor:	Novartis
Study Title:	Study of Efficacy and Safety of NIS793 (With and Without Spartalizumab) in Combination With SOC Chemotherapy in First-line Metastatic Pancreatic Ductal Adenocarcinoma (mPDAC)

15. Sign-off & Approval

Principal Investigator: _____	Date: _____
Clinical Operations Lead: _____	Date: _____
Lead Statistician: _____	Date: _____