

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Phase 2 Study of Pembrolizumab And Chemotherapy With or Without MK-4830 as Neoadjuvant Treatment for High-Grade Serous Ovarian Cancer
Short Title:	Pembrolizumab With Chemotherapy and MK-4830 for Treating Participants With Ovarian Cancer (MK-4830-002)
Protocol Number:	890374987997240463
NCT Number:	NCT05446870
Sponsor Name:	Merck
Investigational Product:	carboplatin
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Change From Baseline in Circulating Tumor Deoxyribonucleic Acid (ctDNA): Blood samples were collected to determine levels of ctDNA. The fold change in the mean mutant/tumor molecules per mL (MTM/mL) at Cycle 3 from baseline is presented.
Secondary Obj.:	1. Participants With Surgery and Pathological Complete Response (pCR): Change From Baseline in ctDNA 2. Association of Change From Baseline in ctDNA With pCR 3. Participants With Surgery and Chemotherapy Response Score (CRS): Change From Baseline in ctDNA

2.2 Background & Rationale

The primary objective is to evaluate in participants with high-grade serous ovarian cancer (HGSOC), whether the reduction from baseline in circulating tumor deoxyribonucleic acid (ctDNA) at Cycle 3 (?ctDNA) is larger in participants receiving MK-4830 + pembrolizumab in combination with standard of care (SOC) therapy than in those receiving pembrolizumab + SOC therapy.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: COMPLETED
Target N:	160

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has histologically-confirmed International Federation of Gynecology and Obstetrics (FIGO) Stage III or Stage IV HGSOC, primary peritoneal cancer, or fallopian tube cancer.
- * Is a candidate for carboplatin and paclitaxel chemotherapy, to be administered in the neoadjuvant and adjuvant setting.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Is a candidate for interval debulking surgery.
- * Is able to provide archival tissue or newly obtained core, incisional, or excisional biopsy of a tumor lesion.
- * Has adequate organ functions.

4.2 Exclusion Criteria

- * Has a non-HGSOC histology.
- * Has a history of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease.
- * Has a known additional malignancy that is progressing or has required active treatment within the past 3 years.
- * Has received prior treatment for any stage of ovarian cancer (OC), including radiation or systemic anticancer therapy.
- * Planned or has been administered intraperitoneal chemotherapy as first-line therapy.
- * Has received prior therapy with an anti-programmed cell death 1 protein (PD-1), anti-programmed cell death 1 ligand 1 (PD-L1), anti-programmed cell death 1 ligand 2 (PD-L2), anti-immunoglobulin-like transcript 4 (ILT4), or anti-human leukocyte antigen (HLA)-G agent or with an agent directed to another stimulatory or coinhibitory T-cell receptor.
- * Has received a live or live-attenuated vaccine within 30 days before the first dose of study intervention.
- * Is currently participating in or has participated in a study of an investigational agent or has used an investigational device within 4 weeks before the first dose of study intervention.
- * Has a diagnosis of immunodeficiency or is receiving chronic systemic steroid therapy (in dosing exceeding 10 mg daily of prednisone equivalent) or any other form of immunosuppressive therapy within 7 days prior the first dose of study medication.
- * Has known active central nervous system (CNS) metastases and/or carcinomatous meningitis.
- * Has severe hypersensitivity to pembrolizumab, carboplatin, paclitaxel (or docetaxel, if applicable), Avastin or biosimilar (if using) and/or any of their excipients.
- * Has an active autoimmune disease that has required systemic treatment in past 2 years.
- * Has an active infection requiring systemic therapy.
- * Has a known history of human immunodeficiency virus (HIV) infection.
- * Has a known history of hepatitis B or known active hepatitis C virus infection.
- * Has received colony-stimulating factors within 4 weeks prior to receiving study intervention on Day 1 of Cycle 1.
- * Has had surgery <6 months prior to Screening to treat borderline ovarian tumors, early-stage OC, or early-stage fallopian tube cancer.
- * Has a known psychiatric or substance abuse disorder that would interfere with the participant's ability to cooperate with the requirements of the study.
- * Has current, clinically relevant bowel obstruction.
- * Has a history of hemorrhage, hemoptysis, or active gastrointestinal (GI) bleeding within 6 months prior to randomization.
- * Has uncontrolled hypertension.
- * Has had an allogenic tissue/solid organ transplant.
- * Has either had major surgery within 3 weeks of randomization or has not recovered from any effects of any major surgery.

11. Study Identifiers & Governance

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Primary ID: 890374987997240463
Registry Number: NCT05446870
Sponsor: Merck
Study Title: Pembrolizumab With Chemotherapy and MK-4830 for Treating Participants With Ovarian Cancer (MK-4830-002)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____